

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
30 August 2001 (30.08.2001)

PCT

(10) International Publication Number
WO 01/62728 A1

(51) International Patent Classification⁷: **C07D 207/04**,
211/06, 295/08, A61K 31/40, 31/435, A61P 11/00, 37/00,
19/00, 31/00

& D Lund, S-221 87 Lund (SE). **KLINGSTEDT, Tomas**
[SE/SE]; AstraZeneca R & D Lund, S-221 87 Lund (SE).
PETTERSSON, Lars [SE/SE]; Näktergalsvägen 33,
S-247 36 Södra Sandby (SE).

(21) International Application Number: PCT/SE01/00403

(22) International Filing Date: 23 February 2001 (23.02.2001)

(74) Agent: **ASTRAZENECA AB**; Global Intellectual Prop-
erty, S-151 85 Södertälje (SE).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

0000620-5	25 February 2000 (25.02.2000)	SE
0002234-3	14 June 2000 (14.06.2000)	SE
0003979-2	31 October 2000 (31.10.2000)	SE

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(71) Applicant (*for all designated States except US*): **AS-
TRAZENECA AB** [SE/SE]; S-151 85 Södertälje (SE).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **BODKIN, Michael**
[GB/GB]; Lilly Research Centre, Erl Wood Manor Sun-
ninghill, Windlesham, Surrey GU20 6PH (GB). **ERIKS-
SON, Tomas** [SE/SE]; AstraZeneca R & D Lund, S-221
87 Lund (SE). **HANSEN, Peter** [SE/SE]; AstraZeneca R
& D Lund, S-221 87 Lund (SE). **HEMMERLING, Mar-
tin** [DE/SE]; AstraZeneca R & D Lund, S-221 87 Lund
(SE). **HENRIKSSON, Krister** [SE/SE]; AstraZeneca R


(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: NOVEL COMPOUNDS



(I)

(57) Abstract: The invention provides compounds of general formula (I) wherein Q, R, R², R⁴, R⁵, R⁶, R⁷ and R⁸ are as defined in the specification, processes for their preparation, pharmaceutical compositions containing them and their use in therapy.

BEST AVAILABLE COPY

WO 01/62728 A1

NOVEL COMPOUNDS

The present invention relates to novel compounds, processes for their preparation, pharmaceutical compositions containing them and their use in therapy.

5

US 5789402 describes certain indole derivatives which are said to be useful for the treatment of diseases which are caused or affected by disorders of the serotonin-affected neurological systems, particularly those relating to the serotonin 1_A receptor and those relating to the uptake of serotonin.

10

Chemokines play an important role in immune and inflammatory responses in various diseases and disorders, including asthma and allergic diseases, as well as autoimmune pathologies such as rheumatoid arthritis and atherosclerosis. These small secreted molecules are a growing superfamily of 8-14 kDa proteins characterised by a conserved
15 four cysteine motif. The chemokine superfamily can be divided into two main groups exhibiting characteristic structural motifs, the Cys-X-Cys (C-X-C) and Cys-Cys (C-C) families. These are distinguished on the basis of a single amino acid insertion between the NH-proximal pair of cysteine residues and sequence similarity.


20 The C-X-C chemokines include several potent chemoattractants and activators of neutrophils such as interleukin-8 (IL-8) and neutrophil-activating peptide 2 (NAP-2).

The C-C chemokines include potent chemoattractants of monocytes and lymphocytes but not neutrophils such as human monocyte chemotactic proteins 1-3 (MCP-1, MCP-2 and
25 MCP-3), RANTES (Regulated on Activation, Normal T Expressed and Secreted), eotaxin and the macrophage inflammatory proteins 1 α and 1 β (MIP-1 α and MIP-1 β).

Studies have demonstrated that the actions of the chemokines are mediated by subfamilies of G protein-coupled receptors, among which are the receptors designated CCR1, CCR2,
30 CCR2A, CCR2B, CCR3, CCR4, CCR5, CCR6, CCR7, CCR8, CCR9, CCR10, CXCR1,


CXCR2, CXCR3 and CXCR4. These receptors represent good targets for drug development since agents which modulate these receptors would be useful in the treatment of disorders and diseases such as those previously mentioned.

- 5 In accordance with the present invention, there is therefore provided a compound of general formula




- 10 wherein,

R represents either a group



or a group



- 15

m is 0, 1, 2 or 3;

each R¹ independently represents halogen, cyano, nitro, carboxyl, hydroxyl,

C₃-C₆ cycloalkyl, C₁-C₆ alkoxy, C₁-C₆ alkoxy carbonyl, C₁-C₆ haloalkyl,

C₁-C₆ haloalkoxy, -NR⁹R¹⁰, C₃-C₆ cycloalkylamino, C₁-C₆ alkylthio,

- 20 C₁-C₆ alkylcarbonyl, C₁-C₆ alkylcarbonylamino, sulphonamido (-SO₂NH₂),

C₁-C₆ alkylsulphonyl, -C(O)NR¹¹R¹², -NR¹³C(O)-(NH)_pR¹⁴, phenyl, or C₁-C₆ alkyl optionally substituted by carboxyl or C₁-C₆ alkoxy carbonyl;

p is 0 or 1;


X represents an oxygen or sulphur atom or a CH_2 , $\text{CH}(\text{CH}_3)$, OCH_2 , CH_2O , CH_2NH , NH or carbonyl group and Y represents a nitrogen atom or a CH or $\text{C}(\text{OH})$ group, provided that when X represents an oxygen or sulphur atom or a CH_2O , CH_2NH or NH group, then Y represents a CH group;

Z^1 represents a bond or a group $(\text{CH}_2)_q$ where q is 1 or 2;

Z^2 represents a bond or a group CH_2 , with the proviso that Z^1 and Z^2 do not both simultaneously represent a bond;

Q represents an oxygen or sulphur atom or a group CH_2 or NH ;

R^2 represents a group



n is 0, 1 or 2;

each R^3 independently represents a C_1 - C_6 alkyl, C_1 - C_6 alkoxy carbonyl, $-\text{CH}_2\text{OH}$ or

carboxyl group;

R^4 , R^5 , R^6 and R^7 each independently represent a hydrogen atom or a C_1 - C_6 alkyl group, or R^4 , R^5 , R^6 and R^7 together represent a C_1 - C_4 alkylene chain linking the two carbon atoms to which they are attached to form a 4- to 7-membered saturated carbocycle,

or R^5 , R^6 and R^7 each represent a hydrogen atom and R^4 and R^8 together with the carbon atoms to which they are attached form a 5- to 6-membered saturated carbocycle;

R^8 represents a hydrogen atom, a C_1 - C_6 alkyl group or is linked to R^4 as defined above;

R^9 and R^{10} each independently represent a hydrogen atom or a C_1 - C_6 alkyl group, or R^9 and R^{10} together with the nitrogen atom to which they are attached form a 4- to 7-membered saturated heterocycle;

R^{11} and R^{12} each independently represent a hydrogen atom or a C_1 - C_6 alkyl group optionally substituted by C_1 - C_6 alkoxycarbonyl;

R^{13} represents a hydrogen atom or a C_1 - C_6 alkyl group;

R^{14} represents a hydrogen atom, or a C_1 - C_6 alkyl group optionally substituted by carboxyl, C_1 - C_6 alkoxy or C_1 - C_6 alkoxycarbonyl;

R^{15} represents carboxyl, C_1 - C_6 alkylcarbonyl, C_1 - C_6 alkoxycarbonyl, C_1 - C_6 alkoxycarbonyl- C_1 - C_6 alkyl or a group $-NR^{17}R^{18}$, $-NHSO_2CH_3$, $-NHC(O)CH_3$, $-C(O)NR^{17}R^{18}$, $-NHC(O)NR^{17}R^{18}$, $-OC(O)NR^{17}R^{18}$, $-OCH_2C(O)NR^{17}R^{18}$, $-NHC(O)OR^{17'}$ or $-OR^{17''}$;

t is 0, 1, 2 or 3;

each R^{16} independently represents halogen, cyano, nitro, carboxyl, hydroxyl, C_3 - C_6 cycloalkyl, C_1 - C_6 alkoxy, C_1 - C_6 alkoxycarbonyl, C_1 - C_6 haloalkyl, C_1 - C_6 haloalkoxy, $-NR^{19}R^{20}$, C_3 - C_6 cycloalkylamino, C_1 - C_6 alkylthio, C_1 - C_6 alkylcarbonyl, C_1 - C_6 alkylcarbonylamino, sulphonamido ($-SO_2NH_2$), C_1 - C_6 alkylsulphonyl, $-C(O)NR^{21}R^{22}$, $-NR^{23}C(O)(NH)_vR^{24}$, phenyl, or C_1 - C_6 alkyl optionally substituted by carboxyl or C_1 - C_6 alkoxycarbonyl;

R^{17} and R^{18} each independently represent (i) a hydrogen atom, (ii) a 5- to 6-membered saturated or unsaturated ring which may comprise at least one heteroatom chosen from nitrogen, oxygen and sulphur, the ring being optionally substituted with at least one substituent selected from halogen, methyl and trifluoromethyl, or (iii) a C_1 - C_6 alkyl group optionally substituted by at least one substituent selected from halogen, trifluoromethyl, carboxyl, C_1 - C_6 alkoxycarbonyl and a 5- to 6-membered saturated or unsaturated ring which may comprise at least one heteroatom chosen from

nitrogen, oxygen and sulphur, the ring being optionally substituted with at least one substituent selected from halogen, methyl and trifluoromethyl, or

R^{17} and R^{18} together with the nitrogen atom to which they are attached form a 4- to 7-membered saturated heterocycle;

5 $R^{17'}$ represents a hydrogen atom, or a C_1 - C_6 alkyl group optionally substituted by carboxyl or C_1 - C_6 alkoxy carbonyl;

$R^{17''}$ is defined as for R^{17} above except that $R^{17''}$ does not represent a hydrogen atom;

R^{19} and R^{20} each independently represent a hydrogen atom or a C_1 - C_6 alkyl group, or R^{19} and R^{20} together with the nitrogen atom to which they are attached form a 4- to 7-

10 membered saturated heterocycle;

R^{21} and R^{22} each independently represent a hydrogen atom or a C_1 - C_6 alkyl group optionally substituted by C_1 - C_6 alkoxy carbonyl;

v is 0 or 1;

R^{23} represents a hydrogen atom or a C_1 - C_6 alkyl group; and


15 R^{24} represents a hydrogen atom, or a C_1 - C_6 alkyl group optionally substituted by carboxyl, C_1 - C_6 alkoxy or C_1 - C_6 alkoxy carbonyl;

provided that when X is an oxygen atom or a group CH_2 , Y is CH , Z^1 and Z^2 each represent a group CH_2 and Q is an oxygen atom, then R^2 is other than an unsubstituted indolyl group;

20 or a pharmaceutically acceptable salt or solvate thereof.


In the context of the present specification, an alkyl substituent group or an alkyl moiety in a substituent group may be linear or branched.

25 In one aspect of the present invention, there is provided a compound of general formula



wherein,

R represents a group



m is 0, 1, 2 or 3;

each R^1 independently represents halogen, cyano, nitro, carboxyl, hydroxyl, C_3 - C_6 cycloalkyl, C_1 - C_6 alkoxy, C_1 - C_6 alkoxycarbonyl, C_1 - C_6 haloalkyl, C_1 - C_6 haloalkoxy, $-NR^9R^{10}$, C_3 - C_6 cycloalkylamino, C_1 - C_6 alkylthio, C_1 - C_6 alkylcarbonyl, C_1 - C_6 alkylcarbonylamino, sulphonamido, C_1 - C_6 alkylsulphonyl, $-C(O)NR^{11}R^{12}$, $-NR^{13}C(O)-(NH)_pR^{14}$, phenyl, or C_1 - C_6 alkyl optionally substituted by carboxyl or C_1 - C_6 alkoxycarbonyl;

p is 0 or 1;


X represents an oxygen or sulphur atom or a CH_2 , $CH(CH_3)$, OCH_2 , CH_2O , CH_2NH , NH or carbonyl group and Y represents a nitrogen atom or a CH or $C(OH)$ group, provided that when X represents an oxygen or sulphur atom or a CH_2O , CH_2NH or NH group, then Y represents a CH group;

Z^1 represents a bond or a group $(CH_2)_q$ where q is 1 or 2;

Z^2 represents a bond or a group CH_2 , with the proviso that Z^1 and Z^2 do not both simultaneously represent a bond;

Q represents an oxygen or sulphur atom or a group CH_2 or NH ;

R^2 represents a group



n is 0, 1 or 2;

each R³ independently represents a C₁-C₆ alkyl, C₁-C₆ alkoxy carbonyl, -CH₂OH or carboxyl group;

R⁴, R⁵, R⁶ and R⁷ each independently represent a hydrogen atom or a C₁-C₆ alkyl group, or R⁴, R⁵, R⁶ and R⁷ together represent a C₁-C₄ alkylene chain linking the two carbon atoms to which they are attached to form a 4- to 7-membered saturated carbocycle, or R⁵, R⁶ and R⁷ each represent a hydrogen atom and R⁴ and R⁸ together with the carbon atoms to which they are attached form a 5- to 6-membered saturated carbocycle;

R⁸ represents a hydrogen atom, a C₁-C₆ alkyl group or is linked to R⁴ as defined above;

R⁹ and R¹⁰ each independently represent a hydrogen atom or a C₁-C₆ alkyl group, or R⁹ and R¹⁰ together with the nitrogen atom to which they are attached form a 4- to 7-membered saturated heterocycle;

R¹¹ and R¹² each independently represent a hydrogen atom or a C₁-C₆ alkyl group optionally substituted by C₁-C₆ alkoxy carbonyl;

R¹³ represents a hydrogen atom or a C₁-C₆ alkyl group;

R¹⁴ represents a hydrogen atom, or a C₁-C₆ alkyl group optionally substituted by carboxyl, C₁-C₆ alkoxy or C₁-C₆ alkoxy carbonyl;

R¹⁵ represents carboxyl, C₁-C₆ alkyl carbonyl, C₁-C₆ alkoxy carbonyl, C₁-C₆ alkoxy carbonyl C₁-C₆ alkyl or a group -NR¹⁷R¹⁸, -NHSO₂CH₃, -NHC(O)CH₃, -C(O)NR¹⁷R¹⁸, -NHC(O)NR¹⁷R¹⁸, -OC(O)NR¹⁷R¹⁸, -OCH₂C(O)NR¹⁷R¹⁸, -NHC(O)OR^{17'} or -OR^{17''};

t is 0, 1, 2 or 3;

each R¹⁶ independently represents halogen, cyano, nitro, carboxyl, hydroxyl, C₃-C₆ cycloalkyl, C₁-C₆ alkoxy, C₁-C₆ alkoxy carbonyl, C₁-C₆ haloalkyl, C₁-C₆ haloalkoxy, -NR¹⁹R²⁰, C₃-C₆ cycloalkylamino, C₁-C₆ alkylthio, C₁-C₆ alkyl carbonyl, C₁-C₆ alkyl carbonylamino, sulphonamido (-SO₂NH₂), C₁-C₆ alkylsulphonyl, -C(O)NR²¹R²², -NR²³C(O)(NH)_vR²⁴, phenyl, or C₁-C₆ alkyl optionally substituted by carboxyl or C₁-C₆ alkoxy carbonyl;

R^{17} and R^{18} each independently represent (i) a hydrogen atom, (ii) a 5- to 6-membered saturated or unsaturated ring which may comprise at least one heteroatom chosen from nitrogen, oxygen and sulphur, the ring being optionally substituted with at least one substituent selected from halogen, methyl and trifluoromethyl, or

(iii) a C_1 - C_6 alkyl group optionally substituted by at least one substituent selected from halogen, trifluoromethyl, carboxyl, C_1 - C_6 alkoxy carbonyl and a 5- to 6-membered saturated or unsaturated ring which may comprise at least one heteroatom chosen from nitrogen, oxygen and sulphur, the ring being optionally substituted with at least one substituent selected from halogen, methyl and trifluoromethyl, or

R^{17} and R^{18} together with the nitrogen atom to which they are attached form a 4- to 7-membered saturated heterocycle;

R^{17} represents a hydrogen atom, or a C_1 - C_6 alkyl group optionally substituted by carboxyl or C_1 - C_6 alkoxy carbonyl;

$R^{17''}$ is defined as for R^{17} above except that $R^{17''}$ does not represent a hydrogen atom;

R^{19} and R^{20} each independently represent a hydrogen atom or a C_1 - C_6 alkyl group, or R^{19} and R^{20} together with the nitrogen atom to which they are attached form a 4- to 7-membered saturated heterocycle;

R^{21} and R^{22} each independently represent a hydrogen atom or a C_1 - C_6 alkyl group optionally substituted by C_1 - C_6 alkoxy carbonyl;


v is 0 or 1;

R^{23} represents a hydrogen atom or a C_1 - C_6 alkyl group; and

R^{24} represents a hydrogen atom, or a C_1 - C_6 alkyl group optionally substituted by carboxyl, C_1 - C_6 alkoxy or C_1 - C_6 alkoxy carbonyl;


or a pharmaceutically acceptable salt or solvate thereof.

In another aspect of the invention, there is provided a compound of general formula



wherein,

R represents a group



m is 0, 1, 2 or 3;

5 each R¹ independently represents halogen, cyano, nitro, carboxyl, hydroxyl, C₃-C₆ cycloalkyl, C₁-C₆ alkoxy, C₁-C₆ alkoxycarbonyl, C₁-C₆ haloalkyl, C₁-C₆ haloalkoxy, -NR⁹R¹⁰, C₃-C₆ cycloalkylamino, C₁-C₆ alkylthio, C₁-C₆ alkylcarbonyl, C₁-C₆ alkylcarbonylamino, sulphonamido, C₁-C₆ alkylsulphonyl, -C(O)NR¹¹R¹², -NR¹³C(O)-(NH)_pR¹⁴, phenyl, or C₁-C₆ alkyl optionally substituted by
10 carboxyl or C₁-C₆ alkoxycarbonyl;

p is 0 or 1;

X represents an oxygen or sulphur atom or a CH₂, CH(CH₃), OCH₂, CH₂O, CH₂NH, NH or carbonyl group and Y represents a nitrogen atom or a CH or C(OH) group, provided that when X represents an oxygen or sulphur atom or a CH₂O, CH₂NH or NH group, then


15 Y represents a CH group;

Z¹ represents a bond or a group (CH₂)_q where q is 1 or 2;

Z² represents a bond or a group CH₂, with the proviso that Z¹ and Z² do not both simultaneously represent a bond;

Q represents an oxygen or sulphur atom or a group CH₂ or NH;

20 R² represents a group



n is 0, 1 or 2;

each R^3 independently represents a C_1 - C_6 alkyl, C_1 - C_6 alkoxy carbonyl, $-CH_2OH$ or
 5 carboxyl group;

R^4 , R^5 , R^6 and R^7 each independently represent a hydrogen atom or a C_1 - C_6 alkyl
 group, or R^4 , R^5 , R^6 and R^7 together represent a C_1 - C_4 alkylene chain linking the two
 carbon atoms to which they are attached to form a 4- to 7-membered saturated carbocycle,
 or R^5 , R^6 and R^7 each represent a hydrogen atom and R^4 and R^8 together with the carbon
 10 atoms to which they are attached form a 5- to 6-membered saturated carbocycle;

R^8 represents a hydrogen atom, a C_1 - C_6 alkyl group or is linked to R^4 as defined
 above;

R^9 and R^{10} each independently represent a hydrogen atom or a C_1 - C_6 alkyl group, or
 R^9 and R^{10} together with the nitrogen atom to which they are attached form a 4- to 7-
 15 membered saturated heterocycle;

R^{11} and R^{12} each independently represent a hydrogen atom or a C_1 - C_6 alkyl group
 optionally substituted by C_1 - C_6 alkoxy carbonyl;

R^{13} represents a hydrogen atom or a C_1 - C_6 alkyl group; and


R^{14} represents a hydrogen atom, or a C_1 - C_6 alkyl group optionally substituted by carboxyl, C_1 - C_6 alkoxy or C_1 - C_6 alkoxycarbonyl;

provided that when X is an oxygen atom or a group CH_2 , Y is CH, Z^1 and Z^2 each represent a group CH_2 and Q is an oxygen atom, then R^2 is other than an unsubstituted

5 indolyl group;


or a pharmaceutically acceptable salt or solvate thereof.

In a further aspect of the invention, there is provided a compound of general formula



wherein,

R represents a group



15 m is 0, 1, 2 or 3;

each R^1 independently represents halogen, cyano, nitro, carboxyl, hydroxyl,

C_3 - C_6 cycloalkyl, C_1 - C_6 alkoxy, C_1 - C_6 alkoxycarbonyl, C_1 - C_6 haloalkyl,

C_1 - C_6 haloalkoxy, $-NR^9R^{10}$, C_3 - C_6 cycloalkylamino, C_1 - C_6 alkylthio,


C_1 - C_6 alkylcarbonyl, C_1 - C_6 alkylcarbonylamino, sulphonamido, C_1 - C_6 alkylsulphonyl,

20 $-C(O)NR^{11}R^{12}$, $-NR^{13}C(O)-(NH)_pR^{14}$, phenyl, or C_1 - C_6 alkyl optionally substituted by carboxyl or C_1 - C_6 alkoxycarbonyl;

p is 0 or 1;

Q represents an oxygen or sulphur atom or a group CH_2 or NH;

R^2 represents a group



R^4 , R^5 , R^6 and R^7 each independently represent a hydrogen atom or a C_1 - C_6 alkyl group, or R^4 , R^5 , R^6 and R^7 together represent a C_1 - C_4 alkylene chain linking the two carbon atoms to which they are attached to form a 4- to 7-membered saturated carbocycle, or R^5 , R^6 and R^7 each represent a hydrogen atom and R^4 and R^8 together with the carbon atoms to which they are attached form a 5- to 6-membered saturated carbocycle;

R^8 represents a hydrogen atom, a C_1 - C_6 alkyl group or is linked to R^4 as defined above;

R^9 and R^{10} each independently represent a hydrogen atom or a C_1 - C_6 alkyl group, or R^9 and R^{10} together with the nitrogen atom to which they are attached form a 4- to 7-membered saturated heterocycle;

R^{11} and R^{12} each independently represent a hydrogen atom or a C_1 - C_6 alkyl group optionally substituted by C_1 - C_6 alkoxy carbonyl;

R^{13} represents a hydrogen atom or a C_1 - C_6 alkyl group;

R^{14} represents a hydrogen atom, or a C_1 - C_6 alkyl group optionally substituted by carboxyl, C_1 - C_6 alkoxy or C_1 - C_6 alkoxy carbonyl;

R^{15} represents carboxyl, C_1 - C_6 alkyl carbonyl, C_1 - C_6 alkoxy carbonyl, C_1 - C_6 alkoxy carbonyl C_1 - C_6 alkyl or a group $-NR^{17}R^{18}$, $-NHSO_2CH_3$, $-NHC(O)CH_3$,

$-C(O)NR^{17}R^{18}$, $-NHC(O)NR^{17}R^{18}$, $-OC(O)NR^{17}R^{18}$, $-OCH_2C(O)NR^{17}R^{18}$,
 $-NHC(O)OR^{17'}$ or $-OR^{17''}$;

t is 0, 1, 2 or 3;

each R^{16} independently represents halogen, cyano, nitro, carboxyl, hydroxyl,
 5 C_3 - C_6 cycloalkyl, C_1 - C_6 alkoxy, C_1 - C_6 alkoxycarbonyl, C_1 - C_6 haloalkyl,
 C_1 - C_6 haloalkoxy, $-NR^{19}R^{20}$, C_3 - C_6 cycloalkylamino, C_1 - C_6 alkylthio,
 C_1 - C_6 alkylcarbonyl, C_1 - C_6 alkylcarbonylamino, sulphonamido ($-SO_2NH_2$),
 C_1 - C_6 alkylsulphonyl, $-C(O)NR^{21}R^{22}$, $-NR^{23}C(O)(NH)_vR^{24}$, phenyl, or C_1 - C_6 alkyl
 optionally substituted by carboxyl or C_1 - C_6 alkoxycarbonyl;

10 R^{17} and R^{18} each independently represent (i) a hydrogen atom, (ii) a 5- to 6-
 membered saturated or unsaturated ring which may comprise at least one heteroatom
 chosen from nitrogen, oxygen and sulphur, the ring being optionally substituted with at
 least one substituent selected from halogen, methyl and trifluoromethyl, or
 (iii) a C_1 - C_6 alkyl group optionally substituted by at least one substituent selected from
 15 halogen, trifluoromethyl, carboxyl, C_1 - C_6 alkoxycarbonyl and a 5- to 6-membered
 saturated or unsaturated ring which may comprise at least one heteroatom chosen from
 nitrogen, oxygen and sulphur, the ring being optionally substituted with at least one
 substituent selected from halogen, methyl and trifluoromethyl, or

R^{17} and R^{18} together with the nitrogen atom to which they are attached form a 4- to 7-
 20 membered saturated heterocycle;

$R^{17'}$ represents a hydrogen atom, or a C_1 - C_6 alkyl group optionally substituted by
 carboxyl or C_1 - C_6 alkoxycarbonyl;

$R^{17''}$ is defined as for R^{17} above except that $R^{17''}$ does not represent a hydrogen atom;

R^{19} and R^{20} each independently represent a hydrogen atom or a C_1 - C_6 alkyl group, or
 25 R^{19} and R^{20} together with the nitrogen atom to which they are attached form a 4- to 7-
 membered saturated heterocycle;

R^{21} and R^{22} each independently represent a hydrogen atom or a C_1 - C_6 alkyl group
 optionally substituted by C_1 - C_6 alkoxycarbonyl;

v is 0 or 1;

30 R^{23} represents a hydrogen atom or a C_1 - C_6 alkyl group; and

R^{24} represents a hydrogen atom, or a C_1 - C_6 alkyl group optionally substituted by carboxyl, C_1 - C_6 alkoxy or C_1 - C_6 alkoxycarbonyl; or a pharmaceutically acceptable salt or solvate thereof.

5 The integer m is preferably 0, 1 or 2.

Each R^1 independently represents halogen (e.g. chlorine, fluorine, bromine or iodine), cyano, nitro, carboxyl, hydroxyl, C_3 - C_6 cycloalkyl (cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl), C_1 - C_6 , preferably C_1 - C_4 , alkoxy (e.g. methoxy, ethoxy, n-propoxy or
10 n-butoxy), C_1 - C_6 , preferably C_1 - C_4 , alkoxycarbonyl (e.g. methoxycarbonyl or ethoxycarbonyl), C_1 - C_6 , preferably C_1 - C_4 , haloalkyl (e.g. trifluoromethyl), C_1 - C_6 , preferably C_1 - C_4 , haloalkoxy (e.g. trifluoromethoxy), $-NR^9R^{10}$, C_3 - C_6 cycloalkylamino (e.g. cyclopropylamino, cyclobutylamino, cyclopentylamino or cyclohexylamino), C_1 - C_6 , preferably C_1 - C_4 , alkylthio (e.g. methylthio or ethylthio),
15 C_1 - C_6 , preferably C_1 - C_4 , alkylcarbonyl (e.g. methylcarbonyl, ethylcarbonyl, n-propylcarbonyl, isopropylcarbonyl, n-butylcarbonyl, n-pentylcarbonyl or n-hexylcarbonyl), C_1 - C_6 , preferably C_1 - C_4 , alkylcarbonylamino (e.g. methylcarbonylamino or ethylcarbonylamino), sulphonamido, C_1 - C_6 , preferably C_1 - C_4 , alkylsulphonyl (e.g. methylsulphonyl, ethylsulphonyl,
20 n-propylsulphonyl, isopropylsulphonyl, n-butylsulphonyl, n-pentylsulphonyl or n-hexylsulphonyl), $-C(O)NR^{11}R^{12}$, $-NR^{13}C(O)-(NH)_pR^{14}$, phenyl, or C_1 - C_6 , preferably C_1 - C_4 , alkyl (e.g. methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, n-pentyl or n-hexyl) optionally substituted by carboxyl or C_1 - C_6 , preferably C_1 - C_4 , alkoxycarbonyl (e.g. methoxycarbonyl or ethoxycarbonyl).

25

Most preferably, each R^1 independently represents halogen (particularly chlorine or fluorine), cyano, nitro, C_1 - C_6 alkoxy (especially methoxy), C_1 - C_6 alkylcarbonyl (especially methylcarbonyl) or C_1 - C_6 alkylcarbonylamino (particularly methylcarbonylamino).

30

Preferably X represents an oxygen atom or a CH₂, OCH₂, CH₂O, NH or carbonyl group.

Preferably Y represents a nitrogen atom or CH group.

- 5 Preferred combinations of X - Y include O - CH, OCH₂ - CH, NH - CH, CH₂O - CH, CH₂ - N, C(O) - N and CH₂ - CH.

Preferred combinations of Y, Z¹ and Z² include:

Y	Z ¹	Z ²
CH	CH ₂	bond
CH	bond	CH ₂
CH	CH ₂	CH ₂
CH	(CH ₂) ₂	bond
N	CH ₂	CH ₂

10

Q preferably represents an oxygen atom.

- Each R³ independently represents a C₁-C₆, preferably C₁-C₄, alkyl (e.g. methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, n-pentyl or n-hexyl), C₁-C₆, preferably
 15 C₁-C₄, alkoxycarbonyl (e.g. methoxycarbonyl or ethoxycarbonyl), -CH₂OH or carboxyl group. It is preferred that R³ represents a methyl, methoxycarbonyl, ethoxycarbonyl, -CH₂OH or carboxyl group.

- R⁴, R⁵, R⁶ and R⁷ each independently represent a hydrogen atom or a C₁-C₆, preferably
 20 C₁-C₄, alkyl (e.g. methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, n-pentyl or n-hexyl), or R⁴, R⁵, R⁶ and R⁷ together represent a C₁-C₄ alkylene chain linking the two carbon atoms to which they are attached to form a 4- to 7-membered saturated carbocycle (e.g. cyclohexyl or preferably cyclopentyl), or R⁵, R⁶ and R⁷ each represent a hydrogen

atom and R⁴ and R⁸ together with the carbon atoms to which they are attached form a 5- to 6-membered saturated carbocycle (preferably cyclopentyl).

R⁸ represents a hydrogen atom, a C₁-C₆, preferably C₁-C₄, alkyl group (e.g. methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, n-pentyl or n-hexyl) or is linked to R⁴ as defined above.

R⁹ and R¹⁰ each independently represent a hydrogen atom or a C₁-C₆, preferably C₁-C₄, alkyl group (e.g. methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, n-pentyl or n-hexyl), or R⁹ and R¹⁰ together with the nitrogen atom to which they are attached form a 4- to 7-membered saturated heterocycle (preferably pyrrolidinyl or piperidinyl).

R¹¹ and R¹² each independently represent a hydrogen atom or a C₁-C₆, preferably C₁-C₄, alkyl group (e.g. methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, n-pentyl or n-hexyl) optionally substituted by a C₁-C₆, preferably C₁-C₄, alkoxycarbonyl substituent group.

R¹³ represents a hydrogen atom or a C₁-C₆, preferably C₁-C₄, alkyl group (e.g. methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, n-pentyl or n-hexyl).

R¹⁴ represents a hydrogen atom, or a C₁-C₆, preferably C₁-C₄, alkyl group (e.g. methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, n-pentyl or n-hexyl) optionally substituted by carboxyl, C₁-C₆, preferably C₁-C₄, alkoxy or C₁-C₆, preferably C₁-C₄, alkoxycarbonyl.

R¹⁵ represents carboxyl, C₁-C₆, preferably C₁-C₄, alkylcarbonyl (e.g. methylcarbonyl, ethylcarbonyl, n-propylcarbonyl, isopropylcarbonyl, n-butylcarbonyl, n-pentylcarbonyl or n-hexylcarbonyl), C₁-C₆, preferably C₁-C₄, alkoxycarbonyl (e.g. methoxycarbonyl or ethoxycarbonyl), C₁-C₆ alkoxycarbonylC₁-C₆ alkyl, preferably

C₁-C₄ alkoxycarbonylC₁-C₄ alkyl (e.g. methoxycarbonylmethyl or methoxycarbonylethyl), or a group -NR¹⁷R¹⁸, -NHSO₂CH₃, -NHC(O)CH₃, -C(O)NR¹⁷R¹⁸, -NHC(O)NR¹⁷R¹⁸, -OC(O)NR¹⁷R¹⁸, -OCH₂C(O)NR¹⁷R¹⁸, -NHC(O)OR^{17'} or -OR^{17''}.

- 5 It is preferred that R¹⁵ represents C₁-C₄ alkoxy (especially methoxy), C₁-C₄ alkylcarbonyl (especially methylcarbonyl or ethylcarbonyl), C₁-C₄ alkoxycarbonylC₁-C₄ alkyl (particularly methoxycarbonylmethyl or methoxycarbonylethyl), -NHC(O)CH₃, -C(O)NR¹⁷R¹⁸, -NHSO₂CH₃ or -NHC(O)NR¹⁷R¹⁸.
- 10 Each R¹⁶ independently represents halogen (e.g. chlorine, fluorine, bromine or iodine), cyano, nitro, carboxyl, hydroxyl, C₃-C₆ cycloalkyl (cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl), C₁-C₆, preferably C₁-C₄, alkoxy (e.g. methoxy, ethoxy, n-propoxy or n-butoxy), C₁-C₆, preferably C₁-C₄, alkoxycarbonyl (e.g. methoxycarbonyl or ethoxycarbonyl), C₁-C₆, preferably C₁-C₄, haloalkyl (e.g. trifluoromethyl),
- 15 C₁-C₆, preferably C₁-C₄, haloalkoxy (e.g. trifluoromethoxy), -NR¹⁹R²⁰, C₃-C₆ cycloalkylamino (e.g. cyclopropylamino, cyclobutylamino, cyclopentylamino or cyclohexylamino), C₁-C₆, preferably C₁-C₄, alkylthio (e.g. methylthio or ethylthio), C₁-C₆, preferably C₁-C₄, alkylcarbonyl (e.g. methylcarbonyl, ethylcarbonyl, n-propylcarbonyl, isopropylcarbonyl, n-butylcarbonyl, n-pentylcarbonyl or
- 20 n-hexylcarbonyl), C₁-C₆, preferably C₁-C₄, alkylcarbonylamino (e.g. methylcarbonylamino or ethylcarbonylamino), sulphonamido, C₁-C₆, preferably C₁-C₄, alkylsulphonyl (e.g. methylsulphonyl, ethylsulphonyl, n-propylsulphonyl, isopropylsulphonyl, n-butylsulphonyl, n-pentylsulphonyl or n-hexylsulphonyl), -C(O)NR²¹R²², -NR²³C(O)-(NH)_vR²⁴, phenyl, or
- 25 C₁-C₆, preferably C₁-C₄, alkyl (e.g. methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, n-pentyl or n-hexyl) optionally substituted by carboxyl or C₁-C₆, preferably C₁-C₄, alkoxycarbonyl (e.g. methoxycarbonyl or ethoxycarbonyl).

Preferably, each R¹⁶ independently represents halogen (particularly chlorine or fluorine), hydroxyl, cyano, C₁-C₄ alkoxy (especially methoxy), C₁-C₄ alkoxycarbonyl (especially methoxycarbonyl), C₁-C₄ haloalkyl (especially trifluoromethyl), C₁-C₄ alkylcarbonyl (particularly methylcarbonyl), phenyl or C₁-C₄ alkyl (e.g. methyl or tert-butyl).

5

R¹⁷ and R¹⁸ each independently represent (i) a hydrogen atom, (ii) a 5- to 6-membered saturated or unsaturated ring which may comprise at least one heteroatom (e.g. one, two or three heteroatoms independently) chosen from nitrogen, oxygen and sulphur (such as cyclopentyl, cyclohexyl, pyrrolyl, imidazolyl, pyridinyl, pyrazinyl, pyridazinyl,

10

pyrimidinyl, thienyl or furanyl), the ring being optionally substituted with at least one substituent (e.g. one, two or three substituents independently) selected from halogen (e.g. fluorine, chlorine, bromine or iodine), methyl and trifluoromethyl, or

(iii) a C₁-C₆, preferably C₁-C₄, alkyl group (e.g. methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, n-pentyl or n-hexyl) optionally substituted by at least one

15

substituent (e.g. one, two or three substituents independently) selected from

halogen (e.g. fluorine, chlorine, bromine or iodine), trifluoromethyl, carboxyl,

C₁-C₆, preferably C₁-C₄, alkoxycarbonyl, especially methoxycarbonyl, and

a 5- to 6-membered saturated or unsaturated ring which may comprise at least one

heteroatom (e.g. one, two or three heteroatoms independently) chosen from nitrogen,

20

oxygen and sulphur (such as cyclopentyl, cyclohexyl, pyrrolyl, imidazolyl, pyridinyl,

pyrazinyl, pyridazinyl, pyrimidinyl, thienyl or furanyl), the ring being optionally

substituted with at least one substituent (e.g. one, two or three substituents independently)

selected from halogen (e.g. fluorine, chlorine, bromine or iodine), methyl and

trifluoromethyl, or

25

R¹⁷ and R¹⁸ together with the nitrogen atom to which they are attached form a 4- to 7-membered saturated heterocycle (preferably pyrrolidinyl or piperidinyl).

R¹⁷ represents a hydrogen atom or a C₁-C₆, preferably C₁-C₄, alkyl group (e.g. methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, n-pentyl or n-hexyl) optionally

substituted by carboxyl or, more preferably, C₁-C₆, preferably C₁-C₄, alkoxycarbonyl, especially methoxycarbonyl,

R¹⁹ and R²⁰ each independently represent a hydrogen atom or a C₁-C₆, preferably C₁-C₄, alkyl group (e.g. methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, n-pentyl or n-hexyl), or R¹⁹ and R²⁰ together with the nitrogen atom to which they are attached form a 4- to 7-membered saturated heterocycle (preferably pyrrolidinyl or piperidinyl).

R²¹ and R²² each independently represent a hydrogen atom or a C₁-C₆, preferably C₁-C₄, alkyl group (e.g. methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, n-pentyl or n-hexyl) optionally substituted by a C₁-C₆, preferably C₁-C₄, alkoxycarbonyl substituent group.

R²³ represents a hydrogen atom or a C₁-C₆, preferably C₁-C₄, alkyl group (e.g. methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, n-pentyl or n-hexyl).

R²⁴ represents a hydrogen atom, or a C₁-C₆, preferably C₁-C₄, alkyl group (e.g. methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, n-pentyl or n-hexyl) optionally substituted by carboxyl, C₁-C₆, preferably C₁-C₄, alkoxy or C₁-C₆, preferably C₁-C₄, alkoxycarbonyl.

Preferred compounds of the invention include:

N-(2-{3-[3R,S-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2R,S-hydroxy-propoxy}-phenyl)-acetamide hydrochloride,

N-(5-Chloro-2-{3-[3R,S-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2R,S-hydroxy-propoxy}-phenyl)-acetamide hydrochloride,

N-(2-{3-[4-(3,4-Dichlorophenoxy)-1-piperidinyl]-2-hydroxypropoxy}phenyl)-acetamide,

1-(2-Aminophenoxy)-3-[4-(3,4-dichlorophenoxy)-1-piperidinyl]-2-propanol dihydrochloride,

N-(2-{3-[3-(3,4-dichlorophenoxy)-1-pyrrolidinyl]-2-hydroxypropoxy} phenyl)-acetamide hydrochloride,

5 2-{3-[4-(4-Fluoro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-benzoic acid methyl ester,

2-(2-{3-[4-(3,4-Difluoro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-benzoylamino)-2-methyl-propionic acid methyl ester,

10 N-[2-(((1*R*,2*S*,3*R*)*-3-[4-(3,4-dichlorophenoxy)-1-piperidinyl]-2-hydroxycyclopentyl}oxy)phenyl]acetamide,

N-[2-(((1*S*,2*S*,3*R*)*-3-[4-(3,4-dichlorophenoxy)-1-piperidinyl]-2-hydroxycyclopentyl}oxy)phenyl]acetamide,

N-[2-(((2,3-*trans*)-3-[4-(3,4-dichlorophenoxy)-1-piperidinyl]-2-hydroxycyclohexyl}oxy)phenyl]acetamide,

15 N-(5-Chloro-2-{3-[3-(3,4-dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

N-(3-Acetyl-2-{3-[3-(3,4-dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-5-methyl-phenyl)-acetamide,

20 N-(2-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-4-methyl-phenyl)-acetamide,

N-(2-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-5-fluoro-phenyl)-acetamide,

1-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-3-(1*H*-indol-7-yloxy)-propan-2-ol,

25 1-(7-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-indol-1-yl)-ethanone,

N-(4-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-biphenyl-3-yl)-acetamide,

N-(2-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-4-fluoro-phenyl)-acetamide,

N-(2-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-5-methyl-phenyl)-acetamide,

N-(2-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

5 N-(5-Chloro-2-{3-[3-(4-chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

N-(3-Acetyl-2-{3-[3-(4-chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-5-methyl-phenyl)-acetamide,

10 N-(2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-4-methyl-phenyl)-acetamide,

N-(2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-5-fluoro-phenyl)-acetamide,

1-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-3-(1H-indol-7-yloxy)-propan-2-ol,

15 1-(7-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-indol-1-yl)-ethanone,

N-(4-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-biphenyl-3-yl)-acetamide,

N-(2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-4-fluoro-phenyl)-acetamide,

20 N-(2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-5-methyl-phenyl)-acetamide,

N-(2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide

25 N-(5-Chloro-2-{3-[3-(4-fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

N-(3-Acetyl-2-{3-[3-(4-fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-5-methyl-phenyl)-acetamide,

N-(2-{3-[3-(4-Fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-4-methyl-phenyl)-acetamide,

N-(5-Fluoro-2-{3-[3-(4-fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

1-[3-(4-Fluoro-phenoxy)-pyrrolidin-1-yl]-3-(1H-indol-7-yloxy)-propan-2-ol,

1-(7-{3-[3-(4-Fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-indol-1-yl)-

5 ethanone,

N-(4-{3-[3-(4-Fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-biphenyl-3-yl)-acetamide,

N-(4-Fluoro-2-{3-[3-(4-fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

10 N-(2-{3-[3-(4-Fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-5-methyl-phenyl)-acetamide,

N-(2-{3-[3-(4-Fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

15 N-(5-Chloro-2-{3-[3-(3,4-difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

N-(3-Acetyl-2-{3-[3-(3,4-difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-5-methyl-phenyl)-acetamide,

N-(2-{3-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-4-methyl-phenyl)-acetamide,

20 N-(2-{3-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-5-fluoro-phenyl)-acetamide,

1-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-3-(1H-indol-7-yloxy)-propan-2-ol,

1-(7-{3-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-indol-1-yl)-ethanone,

25 N-(4-{3-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-biphenyl-3-yl)-acetamide,

N-(2-{3-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-4-fluoro-phenyl)-acetamide,

30 N-(2-{3-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-5-methyl-phenyl)-acetamide,

N-(2-{3-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

N-(5-Chloro-2-{3-[4-(3,4-dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

5 N-(3-Acetyl-2-{3-[4-(3,4-dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-5-methyl-phenyl)-acetamide,

N-(2-{3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-4-methyl-phenyl)-acetamide,

10 N-(2-{3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-5-fluoro-phenyl)-acetamide,

1-(7-{3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-indol-1-yl)-ethanone,

N-(4-{3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-biphenyl-3-yl)-acetamide,

15 N-(2-{3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-4-fluoro-phenyl)-acetamide,

N-(2-{3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-5-methyl-phenyl)-acetamide,

20 N-(2-{3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

N-(5-Chloro-2-{3-[4-(4-chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

N-(3-Acetyl-2-{3-[4-(4-chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-5-methyl-phenyl)-acetamide,

25 N-(2-{3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-4-methyl-phenyl)-acetamide,

N-(2-{3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-5-fluoro-phenyl)-acetamide,

30 1-(7-{3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-indol-1-yl)-ethanone,

N-(4-{3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-biphenyl-3-yl)-acetamide,

N-(2-{3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-4-fluorophenyl)-acetamide,

5 N-(2-{3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-5-methylphenyl)-acetamide,

N-(2-{3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

10 N-{5-Chloro-2-[3-(8-chloro-1,3,4,5-tetrahydro-pyrido[4,3-b]indol-2-yl)-2-hydroxypropoxy]-phenyl}-acetamide,

N-{3-Acetyl-2-[3-(8-chloro-1,3,4,5-tetrahydro-pyrido[4,3-b]indol-2-yl)-2-hydroxypropoxy]-5-methyl-phenyl}-acetamide,

N-{2-[3-(8-Chloro-1,3,4,5-tetrahydro-pyrido[4,3-b]indol-2-yl)-2-hydroxy-propoxy]-4-methyl-phenyl}-acetamide,

15 N-{2-[3-(8-Chloro-1,3,4,5-tetrahydro-pyrido[4,3-b]indol-2-yl)-2-hydroxy-propoxy]-5-fluoro-phenyl}-acetamide,

1-(8-Chloro-1,3,4,5-tetrahydro-pyrido[4,3-b]indol-2-yl)-3-(1H-indol-7-yloxy)-propan-2-ol,

20 1-{7-[3-(8-Chloro-1,3,4,5-tetrahydro-pyrido[4,3-b]indol-2-yl)-2-hydroxy-propoxy]-indol-1-yl}-ethanone,

N-{4-[3-(8-Chloro-1,3,4,5-tetrahydro-pyrido[4,3-b]indol-2-yl)-2-hydroxy-propoxy]-biphenyl-3-yl}-acetamide,

N-{2-[3-(8-Chloro-1,3,4,5-tetrahydro-pyrido[4,3-b]indol-2-yl)-2-hydroxy-propoxy]-4-fluoro-phenyl}-acetamide,

25 N-{2-[3-(8-Chloro-1,3,4,5-tetrahydro-pyrido[4,3-b]indol-2-yl)-2-hydroxy-propoxy]-5-methyl-phenyl}-acetamide,

N-{2-[3-(8-Chloro-1,3,4,5-tetrahydro-pyrido[4,3-b]indol-2-yl)-2-hydroxy-propoxy]-phenyl}-acetamide,

30 N-{5-Chloro-2-[3-(8-fluoro-1,3,4,5-tetrahydro-pyrido[4,3-b]indol-2-yl)-2-hydroxypropoxy]-phenyl}-acetamide,

N-{3-Acetyl-2-[3-(8-fluoro-1,3,4,5-tetrahydro-pyrido[4,3-b]indol-2-yl)-2-hydroxy-propoxy]-5-methyl-phenyl}-acetamide,

N-{2-[3-(8-Fluoro-1,3,4,5-tetrahydro-pyrido[4,3-b]indol-2-yl)-2-hydroxy-propoxy]-4-methyl-phenyl}-acetamide,

5 N-{5-Fluoro-2-[3-(8-fluoro-1,3,4,5-tetrahydro-pyrido[4,3-b]indol-2-yl)-2-hydroxy-propoxy]-phenyl}-acetamide,

1-(8-Fluoro-1,3,4,5-tetrahydro-pyrido[4,3-b]indol-2-yl)-3-(1H-indol-7-yloxy)-propan-2-ol,

10 1-{7-[3-(8-Fluoro-1,3,4,5-tetrahydro-pyrido[4,3-b]indol-2-yl)-2-hydroxy-propoxy]-indol-1-yl}-ethanone,

N-{4-[3-(8-Fluoro-1,3,4,5-tetrahydro-pyrido[4,3-b]indol-2-yl)-2-hydroxy-propoxy]-biphenyl-3-yl}-acetamide,

N-{4-Fluoro-2-[3-(8-fluoro-1,3,4,5-tetrahydro-pyrido[4,3-b]indol-2-yl)-2-hydroxy-propoxy]-phenyl}-acetamide,

15 N-{2-[3-(8-Fluoro-1,3,4,5-tetrahydro-pyrido[4,3-b]indol-2-yl)-2-hydroxy-propoxy]-5-methyl-phenyl}-acetamide,

N-{2-[3-(8-Fluoro-1,3,4,5-tetrahydro-pyrido[4,3-b]indol-2-yl)-2-hydroxy-propoxy]-phenyl}-acetamide,

20 N-(2-{3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

3-(2-{3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propionic acid methyl ester,

1-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-3-(2,6-dimethoxy-phenoxy)-propan-2-ol,

1-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-3-(2-methoxy-phenoxy)-propan-2-ol,

25 2-{3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-N,N-dimethylbenzamide,

1-(2-{3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propan-1-one,

30 1-(2-{3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-ethanone,

3-(2-{3-[3-(4-Fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-
propionic acid methyl ester,

1-(2,6-Dimethoxy-phenoxy)-3-[3-(4-fluoro-phenoxy)-pyrrolidin-1-yl]-propan-2-ol,

1-[3-(4-Fluoro-phenoxy)-pyrrolidin-1-yl]-3-(2-methoxy-phenoxy)-propan-2-ol,

5 (2-{3-[3-(4-Fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-benzoylamino)-
acetic acid methyl ester,

(2-{3-[3-(4-Chloro-phenoxy)methyl]-piperidin-1-yl]-2-hydroxy-propoxy}-
benzoylamino)-acetic acid methyl ester,

2-(2-{3-[3-(4-Fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-benzoylamino)-
10 2-methyl-propionic acid methyl ester,

2-{3-[3-(4-Fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-N,N-dimethyl-
benzamide,

1-(2-{3-[3-(4-Fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-6-methoxy-
phenyl)-ethanone,

15 1-(2-{3-[3-(4-Fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propan-
1-one,

1-(2-{3-[3-(4-Fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-
ethanone,

N-[2-(3-{[1-(3,4-dichlorobenzyl)-4-piperidiny]amino}-2-hydroxypropoxy)-4-
20 methylphenyl]acetamide,

3-(2-{3-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-
propionic acid methyl ester,

1-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-3-(2-methoxy-phenoxy)-propan-2-ol,

(2-{3-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-
25 benzoylamino)-acetic acid methyl ester,

2-{3-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-N,N-dimethyl-
benzamide,

1-(2-{3-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-6-methoxy-
phenyl)-ethanone,

1-(2-{3-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propan-1-one,

1-(2-{3-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-ethanone,

5 *N*-(2-{3-[4-(3,4-dichloroanilino)-1-piperidiny]-2-hydroxypropoxy} phenyl)acetamide,
3-(2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propionic acid methyl ester,

1-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-3-(2,6-dimethoxy-phenoxy)-propan-2-ol,

1-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-3-(2-methoxy-phenoxy)-propan-2-ol,

10 (2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-benzoylamino)-acetic acid, methyl ester,

2-(2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-benzoylamino)-2-methyl-propionic acid methyl ester,

2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-*N,N*-dimethyl-
15 benzamide,

1-(2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-6-methoxy-phenyl)-ethanone,

1-(2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propan-1-one,

20 1-(2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-ethanone,

N-(2-{3-[3-(4-Cyano-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

3-(2-{3-[3-(4-Cyano-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-
25 propionic acid methyl ester,

(2-{3-[3-(4-Cyano-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-benzoylamino)-acetic acid methyl ester,

2-{3-[3-(4-Cyano-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-*N,N*-dimethyl-benzamide,

30 4-{1-[2-Hydroxy-3-(2-propionyl-phenoxy)-propyl]-pyrrolidin-3-yloxy}-benzonitrile,

N-(2-{2-Hydroxy-3-[3-(4-methoxy-phenoxy)-pyrrolidin-1-yl]-propoxy}-phenyl)-acetamide,

N-(4-chloro-2-{3-[4-(3,4-dichloroanilino)-1-piperidiny]-2-hydroxypropoxy} phenyl)acetamide,

5 3-(2-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propionic acid methyl ester,

1-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-3-(2-methoxy-phenoxy)-propan-2-ol,
(2-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-benzoylamino)-acetic acid methyl ester,

10 2-(2-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-benzoylamino)-2-methyl-propionic acid methyl ester,

2-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-N,N-dimethylbenzamide,

1-(2-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-6-methoxy-phenyl)-ethanone,

1-(2-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propan-1-one,

1-(2-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-ethanone,

20 N-(2-{3-[4-(3,4-Difluoro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

3-(2-{3-[4-(3,4-Difluoro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propionic acid methyl ester,

25 2-{3-[4-(3,4-Difluoro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-N,N-dimethylbenzamide,

1-(2-{3-[4-(3,4-Difluoro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propan-1-one,

(2-{3-[4-(3,4-Difluoro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-benzoylamino)-acetic acid methyl ester,

N-(2-{3-[3-(3,4-Difluoro-phenoxy)methyl]-piperidin-1-yl}-2-hydroxy-propoxy)-phenyl)-acetamide,

N-(2-{3-[4-(4-Fluoro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

5 3-(2-{3-[4-(4-Fluoro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propionic acid methyl ester,

1-(2,6-Dimethoxy-phenoxy)-3-[4-(4-fluoro-phenoxy)-piperidin-1-yl]-propan-2-ol,

1-[4-(4-Fluoro-phenoxy)-piperidin-1-yl]-3-(2-methoxy-phenoxy)-propan-2-ol,

1-(2-{3-[4-(4-Fluoro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-ethanone,

10 2-{3-[4-(4-Fluoro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-N,N-dimethyl-benzamide,

1-(2-{3-[4-(4-Fluoro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propan-1-one,

(2-{3-[4-(4-Fluoro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-benzoylamino)-acetic acid methyl ester,

15 N-(2-{3-[3-(4-Fluoro-phenoxy)methyl]-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

3-(2-{3-[3-(4-Fluoro-phenoxy)methyl]-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propionic acid methyl ester,

20 1-[3-(4-Fluoro-phenoxy)methyl]-piperidin-1-yl]-3-(2-methoxy-phenoxy)-propan-2-ol,

1-(2-{3-[3-(4-Fluoro-phenoxy)methyl]-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-ethanone,

2-(2-{3-[3-(4-Fluoro-phenoxy)methyl]-piperidin-1-yl]-2-hydroxy-propoxy}-benzoylamino)-2-methyl-propionic acid methyl ester,

25 2-{3-[3-(4-Fluoro-phenoxy)methyl]-piperidin-1-yl]-2-hydroxy-propoxy}-N,N-dimethyl-benzamide,

1-(2-{3-[3-(4-Fluoro-phenoxy)methyl]-piperidin-1-yl]-2-hydroxy-propoxy}-6-methoxy-phenyl)-ethanone,

30 N-(2-{3-[4-(4-Acetylamino-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

N-(4-{1-[3-(2-Acetyl-phenoxy)-2-hydroxy-propyl]-piperidin-4-yloxy}-phenyl)-acetamide,

N-(4-cyano-2-{3-[4-(3,4-dichloroanilino)-1-piperidiny]-2-hydroxypropoxy}phenyl)acetamide,

5 3-(2-{3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propionic acid methyl ester,

1-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-3-(2-methoxy-phenoxy)-propan-2-ol,

1-(2-{3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-ethanone,

10 2-(2-{3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-benzoylamino)-2-methyl-propionic acid methyl ester,

2-{3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-N,N-dimethyl-benzamide,

15 1-(2-{3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-6-methoxy-phenyl)-ethanone,

1-(2-{3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propan-1-one,

(2-{3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-benzoylamino)-acetic acid methyl ester,

20 N-(2-{3-[3-(4-Chloro-phenoxy)methyl]-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

3-(2-{3-[3-(4-Chloro-phenoxy)methyl]-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propionic acid methyl ester,

25 1-(2-{3-[3-(4-Chloro-phenoxy)methyl]-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-ethanone,

2-{3-[3-(4-Chloro-phenoxy)methyl]-piperidin-1-yl]-2-hydroxy-propoxy}-N,N-dimethyl-benzamide,

1-(2-{3-[3-(4-Chloro-phenoxy)methyl]-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propan-1-one,

N-[2-({(1*R*,2*R*)-2-[4-(3,4-dichlorophenoxy)-1-piperidinyl]-1-hydroxycyclopentyl}methoxy)phenyl]acetamide,

Methyl (2*S*,4*R*)-1-{3-[2-(acetylamino)phenoxy]-2-hydroxypropyl}-4-[(4-chlorobenzyl)oxy]-2-pyrrolidinecarboxylate hydrochloride,

5 N-(2-{3-[4-(3,4-Dichloroanilino)-1-piperidinyl]-2-hydroxypropoxy}-4-methylphenyl)acetamide,

N-(2-{3-[4-(4-Chloroanilino)-1-piperidinyl]-2-hydroxypropoxy}phenyl)acetamide,

N-(4-Chloro-2-{3-[4-(4-chloroanilino)-1-piperidinyl]-2-hydroxypropoxy}phenyl)-acetamide,

10 N-(2-{3-[4-(4-Chloroanilino)-1-piperidinyl]-2-hydroxypropoxy}-4-cyanophenyl)acetamide,

N-(2-{3-[4-(4-Chloroanilino)-1-piperidinyl]-2-hydroxypropoxy}-4-methylphenyl)acetamide,

15 N-(5-Chloro-2-{3-[4-(4-fluoroanilino)-1-piperidinyl]-2-hydroxypropoxy}phenyl)acetamide,

N-(5-Chloro-2-{3-[4-(3,4-difluoroanilino)-1-piperidinyl]-2-hydroxypropoxy}phenyl)acetamide,

N-(5-Cyano-2-{3-[4-(4-fluoroanilino)-1-piperidinyl]-2-hydroxypropoxy}-phenyl)acetamide,

20 N-(5-Cyano-2-{3-[4-(3,4-difluoroanilino)-1-piperidinyl]-2-hydroxypropoxy}phenyl)acetamide,

N-(2-{3-[4-(4-Fluoroanilino)-1-piperidinyl]-2-hydroxypropoxy}-4-methylphenyl)acetamide,

25 N-(2-{3-[4-(3,4-Difluoroanilino)-1-piperidinyl]-2-hydroxypropoxy}-4-methylphenyl)acetamide,

N-(2-{3-[3(*S*)-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-(*R*)-hydroxy-propoxy-phenyl)acetamide,

N-(2-{3-[3*S*-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2*S*-hydroxy-propoxy}-phenyl)-acetamide hydrochloride,

N-(2-{3-[3(R)-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-(S)-hydroxy-propoxy-phenyl}acetamide,

N-[5-Chloro-2-({(2S)-3-[(3S)-3-(4-chloro-phenoxy)pyrrolidinyl]-2-hydroxypropyl}oxy)phenyl]acetamide,

5 N-[5-Chloro-2-({(2R)-3-[(3R)-3-(4-chloro-phenoxy)pyrrolidinyl]-2-hydroxypropyl}oxy)phenyl]acetamide,

N-[5-Chloro-2-({(2S)-3-[(3R)-3-(4-chloro-phenoxy)pyrrolidinyl]-2-hydroxypropyl}oxy)phenyl]acetamide,

10 N-[5-Chloro-2-({(2R)-3-[(3S)-3-(4-chloro-phenoxy)pyrrolidinyl]-2-hydroxypropyl}oxy)phenyl]acetamide,

N-(2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-4,5-difluoro-phenyl)-acetamide,

N-{5-Chloro-2-[2-hydroxy-3-(3-phenoxy-pyrrolidin-1-yl)-propoxy]-phenyl}-acetamide,

15 N-(5-Chloro-2-{2-hydroxy-3-[3-(4-nitro-phenoxy)-pyrrolidin-1-yl]-propoxy}-phenyl)-acetamide,

N-(5-Acetyl-2-{3-[3-(3,4-dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

20 4-Acetylamino-3-{3-[3-(3,4-dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-benzoic acid methyl ester,

N-(3-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-naphthalen-2-yl)-acetamide,

N-(2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-5-cyano-phenyl)-acetamide,

25 4-Acetylamino-3-{3-[3-(4-chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-benzoic acid methyl ester,

N-(3-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-naphthalen-2-yl)-acetamide,

30 N-(5-Cyano-2-{3-[4-(3,4-dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

N-(2-{3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-5-trifluoromethyl-phenyl)-acetamide,

N-(5-Chloro-2-{3-[3-(4-fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide trifluoroacetate,

5 *N*-(5-Acetyl-2-{3-[3-(4-chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide trifluoroacetate,

N-(2-{3-[3-(4-Chlorophenoxy)-1-pyrrolidinyl]-2-hydroxypropoxy}phenyl)-methanesulfonamide,

10 *N*-(5-Chloro-2-[3-[3,4-dichlorophenoxy)-1-pyrrolidinyl]-2-hydroxypropoxy)-phenyl)urea,

1-(3-{2-[(Aminocarbonyl)amino]phenoxy}-2-hydroxypropyl)-3-(4-chlorophenoxy)pyrrolidinium 2,2,2-trifluoroacetate,

1-(3-{2-[(Aminocarbonyl)amino]phenoxy}-2-hydroxypropyl)-3-(3,4-dichlorophenoxy)pyrrolidinium 2,2,2-trifluoroacetate,

15 1-(3-{2-[(Aminocarbonyl)amino]-4-chlorophenoxy}-2-hydroxypropyl)-3-(4-chlorophenoxy)pyrrolidinium 2,2,2-trifluoroacetate,

N-(2-{3-[3-(4-Chlorophenoxy)-1-pyrrolidinyl]-2-hydroxypropoxy}phenyl)-*N'*-ethylurea hydrochloride,

20 *N*-(2-{3-[3-(4-Chlorophenoxy)-1-pyrrolidinyl]-2-hydroxypropoxy}phenyl)-*N'*-methylurea hydrochloride,

(2*S*,4*S*)-1-{3-[2-(Acetylamino)phenoxy]-2-hydroxypropyl}-4-(4-chlorophenoxy)-2-pyrrolidinecarboxylic acid; compound with trifluoroacetic acid,

Ethyl (2*S*,4*S*)-1-{3-[2-(acetylamino)phenoxy]-2-hydroxypropyl}-4-(3,4-dichlorophenoxy)-2-pyrrolidinecarboxylate; trifluoroacetic acid salt,

25 *N*-[2-({(2*S*)-3-[(2*S*,4*S*)-4-(4-Chlorophenoxy)-2-(hydroxymethyl)pyrrolidinyl]-2-hydroxypropyl}oxy)phenyl]acetamide; trifluoroacetic acid salt,

N-[2-({(2*R*)-3-[(2*S*,4*S*)-4-(4-Chlorophenoxy)-2-(hydroxymethyl)pyrrolidinyl]-2-hydroxypropyl}oxy)phenyl]acetamide; trifluoroacetic acid salt,

30 *N*-(2-{3-[3-(4-Chlorophenoxy)-1-pyrrolidinyl]-2-hydroxy-2-methylpropoxy}phenyl)acetamide hydrochloride,

N-(2-((1*S**,2*R**,3*S**)-3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-cyclopentyloxy)-phenyl)-acetamide,

N-(2-((1*R**,2*R**,3*S**)-3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-cyclopentyloxy)-phenyl)-acetamide,

5 *N*-(2-((2*R**,3*R**)-3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-butoxy)-phenyl)-acetamide,

N-(2-((1*S**,2*R**,3*S**)-3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-cyclopentyloxy)-phenyl)-acetamide,

10 *N*-(2-((2*R**,3*S**)-3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-butoxy)-phenyl)-acetamide,

N-(2-((2*R**,3*R**)-3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-butoxy)-phenyl)-acetamide,

N-(2-((2*R**,3*S**)-3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-butoxy)-phenyl)-acetamide,

15 *N*-(2-((1*S**,2*R**,3*S**)-3-[4-(3-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-cyclopentyloxy)-phenyl)-acetamide,

N-[5-Chloro-2-(((1*S*,2*R*,3*S*)*-3-[4-(3,4-dichlorophenoxy)-1-piperidinyl]-2-hydroxycyclopentyl)oxy)phenyl]acetamide,

20 *N*-[4-Fluoro-2-(((1*S*,2*R*,3*S*)*-3-[4-(3,4-dichlorophenoxy)-1-piperidinyl]-2-hydroxycyclopentyl)oxy)phenyl]acetamide,

N-(2-{3-[4-(3,4-Dichlorobenzyl)-1-piperazinyl]-2-hydroxypropoxy}-phenyl)acetamide dihydrochloride,

N-(2-{3-[4-(3,4-Dichlorobenzyl)-1-piperazinyl]-2-hydroxypropoxy}-4-fluorophenyl)acetamide,

25 *N*-(2-{3-[4-(3,4-Dichlorobenzyl)-2,5-dimethyl-1-piperazinyl]-2-hydroxypropoxy}phenyl)acetamide,

N-(5-Chloro-2-{3-[4-(3,4-dichlorobenzyl)-1-piperazinyl]-2-hydroxypropoxy}phenyl)acetamide,

30 *N*-(5-Chloro-2-{3-[4-(3,4-dichlorobenzyl)-2,5-dimethyl-1-piperazinyl]-2-hydroxypropoxy}phenyl)acetamide,

N-(2-{3-[4-(3,4-Dichlorobenzyl)-2,5-dimethyl-1-piperazinyl]-2-hydroxypropoxy}-4-methylphenyl)acetamide,

N-(2-{3-[4-(3,4-Dichlorobenzyl)-2,5-dimethyl-1-piperazinyl]-2-hydroxypropoxy}-4-fluorophenyl)acetamide,

5 *N*-(2-{3 [(*S***R**)-4-(3,4-Dichlorobenzyl)-2,5-dimethyl-1-piperazinyl]-2-hydroxypropoxy} phenyl)acetamide,

N-(2-{3 [(*S***R**)-4-(4-Chlorobenzyl)-2,5-dimethyl-1-piperazinyl]-2-hydroxypropoxy} phenyl)acetamide,

10 *N*-(5-Chloro-2-{3-[(*S***R**)-4-(3,4-dichlorobenzyl)-2,5-dimethylpiperazinyl]-2-hydroxypropoxy} phenyl)acetamide,

N-(5-Chloro-2-{3-[(*S***R**)-4-(4-chlorobenzyl)-2,5-dimethylpiperazinyl]-2-hydroxypropoxy} phenyl)acetamide,

1-(5-Chloro-2-{3-[4-(4-chlorobenzoyl)-1-piperazinyl]-2-hydroxypropoxy} phenyl)-1-ethanone,

15 *N*-(5-Cyano-2-{3-[(*S***R**)-4-(3,4-dichlorobenzyl)-2,5-dimethylpiperazinyl]-2-hydroxypropoxy} phenyl)acetamide,

N-(2-{3-[(*S***R**)-4-(4-Chlorobenzyl)-2,5-dimethylpiperazinyl]-2-hydroxypropoxy}-5-cyanophenyl)acetamide,

20 *N*-(5-Chloro-2-{3-[4-(4-chlorobenzyl)-1-piperazinyl]-2-hydroxypropoxy}-phenyl)acetamide,

N-(4-Chloro-2-{3-[4-(4-chlorobenzyl)-2,5-dimethyl-1-piperazinyl]-2-hydroxypropoxy} phenyl)acetamide,

N-(2-{3-[4-(4-Chlorobenzoyl)-1-piperazinyl]-2-hydroxypropoxy}-5-cyanophenyl)acetamide,

25 *N*-(2-{3-[4-(4-Chlorobenzoyl)-1-piperazinyl]-2-hydroxypropoxy}-4-methylphenyl)acetamide,

N-[5-Chloro-2-((1*R*,2*S*,3*R*)-3-[(3*S*)-3-(4-chlorophenoxy)pyrrolidinyl]-2-hydroxycyclopentyl}oxy)phenyl]acetamide,

30 *N*-(2-[(2*S*)-(3-[(3*S*)-3-[(4-Chlorophenyl)oxy]-1-pyrrolidinyl)-2-hydroxypropyl)oxy]-4-fluorophenyl)acetamide,

N-[2-({(2*S*)-3-[(3*S*)-3-(4-Chlorobenzyl)pyrrolidinyl]-2-hydroxypropyl} oxy)phenyl]acetamide hydrochloride,

N-(5-Chloro-2-{3-[3-(4-chloro-benzyl)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide trifluoroacetic acid salt,

5 *N*-(2-{3-[3-(4-Chlorophenoxy)-1-pyrrolidinyl]-2-hydroxypropoxy}-4-methylphenyl)-1-pyrrolidinecarboxamide trifluoroacetate,

N-(2-{3-[3-(4-Chlorophenoxy)-1-pyrrolidinyl]-2-hydroxypropoxy}-4-hydroxyphenyl)acetamide trifluoroacetate,

10 *N*-[2-({(2*S*)-3-[4-(3,4-Dichlorophenoxy)-1-piperidinyl]-2-hydroxypropyl} oxy)-4-fluorophenyl]acetamide trifluoroacetic acid salt,

N-(2-(3-(4-Chloro-phenoxy)-pyrrolidin-1-yl)-2-hydroxy-propoxy)-4,6-difluorophenyl)-acetamide hydrochloride,

N-[2-({(2*S*)-3-[(2*S*,4*S*)-4-(4-Chlorophenoxy)-2-methylpyrrolidinyl]-2-hydroxypropyl} oxy)-4-fluorophenyl]acetamide trifluoroacetic acid salt,

15 *N*-[2-({(2*S*)-3-[(3*R*)-3-(4-Chlorobenzyl)pyrrolidinyl]-2-hydroxypropyl} oxy)phenyl]acetamide hydrochloride,

N-{2-[(2*R*)-(3-{(3*S*)-3-[(4-Chlorophenyl)oxy]-1-pyrrolidinyl}-2-hydroxypropyl)oxy]-4-fluorophenyl} acetamide,

20 *N*-[2-({(2*S*)-3-[(2*R*,4*S*)-4-(4-Chlorophenoxy)-2-methylpyrrolidinyl]-2-hydroxypropyl} oxy) phenyl]acetamide trifluoroacetic acid salt,

N-{2-[(2*S*)-(3-{(3*R*)-3-[(4-Chlorophenyl)oxy]-1-pyrrolidinyl}-2-hydroxypropyl)oxy]-4-fluorophenyl} acetamide,

N-(2-{3-[3-(4-Chlorophenoxy)-1-pyrrolidinyl]-2-hydroxypropoxy}-4-methylphenyl)-*N,N*-dimethylurea trifluoroacetate,

25 *N*-(2-{3-[3-(4-Chloroanilino)-1-pyrrolidinyl]-2-hydroxypropoxy}phenyl)acetamide,

N-{2-[(3-{3-[(4-Chlorophenyl)oxy]-1-pyrrolidinyl}-2-hydroxy-1-methylpropyl)oxy]phenyl} acetamide hydrochloride,

N-(2-{3-[3-(4-Chlorophenoxy)-1-pyrrolidinyl]-2-hydroxypropoxy}-4-methoxyphenyl)acetamide hydrochloride,

N-(2-[3-(4-Chloro-benzyloxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy)-phenyl)-acetamide trifluoroacetic acid salt,

N-(2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-2-methyl-propoxy}-phenyl)-acetamide,

5 N-(2-((1*S*,2*R*,3*S*)*-3-[(3*S*)-3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-cyclopentyloxy}-5-chloro-phenyl)-acetamide (diastereomeric mixture),

N-[2-((2*R*,3*S*)*-3-[(3*S*)-3-(4-Chlorophenoxy)pyrrolidinyl]-2-hydroxybutyl}oxy)-4-methylphenyl]acetamide (diastereomeric mixture),

10 N-{2-[(3-{4-[(3,4-Dichlorophenyl)oxy]-1-piperidinyl}-2-hydroxy-2-methylpropyl)oxy]-4-fluorophenyl}acetamide hydrochloride,

N-(2-((1*S*,2*R*,3*S*)*-3-[(3*S*)-3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-cyclopentyloxy)-4-fluoro-phenyl)-acetamide (diastereomeric mixture),

N-(5-Chloro-2-{3-[3-(3,4-difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

15 N-(5-Chloro-2-{3-[3-(4-fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

N-(4-Cyano-2-{3-[4-(3,4-dichloroanilino)-1-piperidinyl]-2-hydroxypropoxy}phenyl)acetamide,

20 N-(4-Hydroxy-2-((1*S*,2*R*,3*S*)*-3-[(3*S*)-3-(4-chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-cyclopentyloxy)-phenyl)-acetamide (diastereomeric mixture),

N-(4-Hydroxy-2-((1*S*,2*R*,3*S*)-3-[(3*S*)-3-(4-chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-cyclopentyloxy)-phenyl)-acetamide,

N-(4-Hydroxy-2-((1*R*,2*S*,3*R*)-3-[(3*S*)-3-(4-chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-cyclopentyloxy)-phenyl)-acetamide,

25 N-[2-((1*S*,2*R*,3*S*)-3-[(3*S*)-3-(4-Chlorophenoxy)pyrrolidinyl]-2-hydroxycyclopentyl}oxy)phenyl]acetamide,

N-[2-((1*R*,2*S*,3*R*)-3-[(3*S*)-3-(4-Chlorophenoxy)pyrrolidinyl]-2-hydroxycyclopentyl}oxy)phenyl]acetamide,

30 N-[5-Chloro-2-((1*S*,2*R*,3*S*)-3-[(3*S*)-3-(4-chlorophenoxy)pyrrolidinyl]-2-hydroxycyclopentyl}oxy)phenyl]acetamide,

N-{5-Chloro-2-[[[(1*S*,2*R*,3*S*)*-3-[[1-(4-chlorobenzyl)-4-piperidinyl]amino]-2-hydroxycyclopentyl]oxy]phenyl}acetamide (racemic mixture), and

N-[2-({(2*S*)-3-[(3*S*)-3-(4-Chlorophenoxy)pyrrolidinyl]-2-hydroxypropyl}oxy)-4-hydroxyphenyl]acetamide.

5


The present invention further provides a process for the preparation of a compound of formula (I) as defined above which comprises,

(a) reacting a compound of general formula

10




wherein R is as defined in formula (I), with a compound of general formula




wherein Q, R², R⁴, R⁵, R⁶, R⁷ and R⁸ are as defined in formula (I); or

15 (b) reacting a compound of general formula



wherein R, R⁴, R⁵, R⁶, R⁷ and R⁸ are as defined in formula (I), with a compound of general formula

20



wherein L¹ represents a hydrogen atom or an activating group (e.g. Li when Q is CH₂) and Q and R² are as defined in formula (I);


and optionally thereafter converting the compound of formula (I) to a further compound of formula (I); and, if desired, forming a pharmaceutically acceptable salt or solvate of the compound of formula (I).

In one aspect, the invention provides a process for the preparation of a compound of formula (I') as hereinbefore defined which comprises,

(a) reacting a compound of general formula




wherein R is as defined in formula (I'), with a compound of general formula




wherein Q, R², R⁴, R⁵, R⁶, R⁷ and R⁸ are as defined in formula (I'); or

(b) reacting a compound of general formula



wherein R, R⁴, R⁵, R⁶, R⁷ and R⁸ are as defined in formula (I'), with a compound of general formula



wherein L¹ represents a hydrogen atom or an activating group (e.g. Li when Q is CH₂) and Q and R² are as defined in formula (I');


and optionally thereafter converting the compound of formula (I') to a further compound of formula (I'); and, if desired, forming a pharmaceutically acceptable salt or solvate of the compound of formula (I').

In another aspect, the invention provides a process for the preparation of a compound of formula (I'') as hereinbefore defined which comprises,

(a) reacting a compound of general formula




5 wherein R is as defined in formula (I''), with a compound of general formula




wherein Q, R², R⁴, R⁵, R⁶, R⁷ and R⁸ are as defined in formula (I''); or

(b) reacting a compound of general formula



wherein R, R⁴, R⁵, R⁶, R⁷ and R⁸ are as defined in formula (I''), with a compound of general formula



wherein L¹ represents a hydrogen atom or an activating group (e.g. Li when Q is CH₂) and Q and R² are as defined in formula (I'');


and optionally thereafter converting the compound of formula (I'') to a further compound of formula (I''); and, if desired, forming a pharmaceutically acceptable salt or solvate of the compound of formula (I'').

In yet another aspect, the invention provides a process for the preparation of a compound of formula (I''') as hereinbefore defined which comprises,

(a) reacting a compound of general formula




wherein R is as defined in formula (I'''), with a compound of general formula




wherein Q, R², R⁴, R⁵, R⁶, R⁷ and R⁸ are as defined in formula (I'''); or

(b) reacting a compound of general formula



wherein R, R⁴, R⁵, R⁶, R⁷ and R⁸ are as defined in formula (I'''), with a compound of general formula



wherein L¹ represents a hydrogen atom or an activating group (e.g. Li when Q is CH₂) and Q and R² are as defined in formula (I''');

and optionally thereafter converting the compound of formula (I''') to a further compound of formula (I'''); and, if desired, forming a pharmaceutically acceptable salt or solvate of the compound of formula (I''').

The processes of the invention may conveniently be carried out in a solvent, e.g. an organic solvent such as an alcohol (e.g. methanol or ethanol), a hydrocarbon (e.g. toluene) or acetonitrile at a temperature of, for example, 15°C or above such as a temperature in the range from 20 to 120°C.

Compounds of formulae (II), (II'), (II''), (II'''), (III), (III'), (III''), (III'''), (IV), (IV'), (IV''), (IV'''), (V), (V'), (V'') and (V''') are either commercially available, are well known in the literature or may be prepared easily using known techniques.

Compounds of formula (I), (I'), (I'') or (I''') can be converted into further compounds of formula (I), (I'), (I'') or (I''') using standard procedures. For example, a compound of formula (I) in which R¹⁵ represents -NHC(O)CH₃ can be converted to a further compound of formula (I) in which R¹⁵ represents -NH₂ by a hydrolysis reaction in the presence of hydrochloric acid.

It will be appreciated by those skilled in the art that in the processes of the present invention certain functional groups such as hydroxyl or amino groups in the starting reagents or intermediate compounds may need to be protected by protecting groups. Thus, the preparation of the compounds of formula (I), (I'), (I'') or (I''') may involve, at an appropriate stage, the removal of one or more protecting groups.

The protection and deprotection of functional groups is described in 'Protective Groups in Organic Chemistry', edited by J.W.F. McOmie, Plenum Press (1973) and 'Protective Groups in Organic Synthesis', 2nd edition, T.W. Greene and P.G.M. Wuts, Wiley-Interscience (1991).

The compounds of formula (I), (I'), (I'') or (I''') above may be converted to a pharmaceutically acceptable salt or solvate thereof, preferably an acid addition salt such as a hydrochloride, hydrobromide, phosphate, acetate, fumarate, maleate, tartrate, citrate, oxalate, methanesulphonate or *p*-toluenesulphonate.

Compounds of formula (I), (I'), (I'') or (I''') are capable of existing in stereoisomeric forms.

It will be understood that the invention encompasses the use of all geometric and optical isomers of the compounds of formula (I), (I'), (I'') or (I''') and mixtures thereof including racemates. The use of tautomers and mixtures thereof also form an aspect of the present invention. Enantiomerically pure forms are particularly desired.

The compounds of formula (I), (I'), (I'') or (I''') have activity as pharmaceuticals, in particular as modulators of chemokine receptor (especially MIP-1 α chemokine receptor) activity, and may be used in the treatment of autoimmune, inflammatory, proliferative and hyperproliferative diseases and immunologically-mediated diseases including rejection of transplanted organs or tissues and Acquired Immunodeficiency Syndrome (AIDS).

Examples of these conditions are:

- (1) **(the respiratory tract)** airways diseases including chronic obstructive pulmonary disease (COPD) such as irreversible COPD; asthma, such as bronchial, allergic, intrinsic, extrinsic and dust asthma, particularly chronic or inveterate asthma (e.g. late asthma and airways hyper-responsiveness); bronchitis; acute, allergic, atrophic rhinitis and chronic rhinitis including rhinitis caseosa, hypertrophic rhinitis, rhinitis purulenta, rhinitis sicca and rhinitis medicamentosa; membranous rhinitis including croupous, fibrinous and pseudomembranous rhinitis and scrofulous rhinitis; seasonal rhinitis including rhinitis nervosa (hay fever) and vasomotor rhinitis; sarcoidosis, farmer's lung and related diseases, fibroid lung and idiopathic interstitial pneumonia;
- (2) **(bone and joints)** rheumatoid arthritis, seronegative spondyloarthropathies (including ankylosing spondylitis, psoriatic arthritis and Reiter's disease), Behcet's disease, Sjogren's syndrome and systemic sclerosis;
- (3) **(skin)** psoriasis, atopic dermatitis, contact dermatitis and other eczematous dermatides, seborrhectic dermatitis, Lichen planus, Pemphigus, bullous Pemphigus, Epidermolysis bullosa, urticaria, angiodermas, vasculitides, erythemas, cutaneous eosinophilias, uveitis, Alopecia areata and vernal conjunctivitis;
- (4) **(gastrointestinal tract)** Coeliac disease, proctitis, eosinophilic gastro-enteritis, mastocytosis, Crohn's disease, ulcerative colitis, food-related allergies which have effects remote from the gut, e.g., migraine, rhinitis and eczema;

- (5) (other tissues and systemic disease) multiple sclerosis, atherosclerosis, Acquired Immunodeficiency Syndrome (AIDS), lupus erythematosus, systemic lupus, erythematosus, Hashimoto's thyroiditis, myasthenia gravis, type I diabetes, nephrotic syndrome, eosinophilia fascitis, hyper IgE syndrome, lepromatous leprosy, sezary syndrome and idiopathic thrombocytopenia pupura;
- (6) (allograft rejection) acute and chronic following, for example, transplantation of kidney, heart, liver, lung, bone marrow, skin and cornea; and chronic graft versus host disease;
- (7) cancers, especially non-small cell lung cancer (NSCLC) and squamous sarcoma;
- (8) diseases in which angiogenesis is associated with raised CXCR2 chemokine levels (e.g. NSCLC); and
- (9) cystic fibrosis, stroke, re-perfusion injury in the heart, brain, peripheral limbs and sepsis.

Thus, the present invention provides a compound of formula (I), (I'), (I'') or (I'''), or a pharmaceutically-acceptable salt or solvate thereof, as hereinbefore defined for use in therapy.

In a further aspect, the present invention provides the use of a compound of formula (I), (I'), (I'') or (I'''), or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore defined in the manufacture of a medicament for use in therapy.

In the context of the present specification, the term "therapy" also includes "prophylaxis" unless there are specific indications to the contrary. The terms "therapeutic" and "therapeutically" should be construed accordingly.

The invention also provides a method of treating an inflammatory disease in a patient suffering from, or at risk of, said disease, which comprises administering to the patient a therapeutically effective amount of a compound of formula (I), (I'), (I'') or (I'''), or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore defined.

5

The invention still further provides a method of treating an airways disease in a patient suffering from, or at risk of, said disease, which comprises administering to the patient a therapeutically effective amount of a compound of formula (I), (I'), (I'') or (I'''), or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore defined.

10

For the above-mentioned therapeutic uses the dosage administered will, of course, vary with the compound employed, the mode of administration, the treatment desired and the disorder indicated. The daily dosage of the compound of formula (I), (I'), (I'') or (I''') may be in the range from 0.001 mg/kg to 30 mg/kg.

15

The compounds of formula (I), (I'), (I'') or (I''') and pharmaceutically acceptable salts and solvates thereof may be used on their own but will generally be administered in the form of a pharmaceutical composition in which the formula (I), (I'), (I'') or (I''') compound/salt/solvate (active ingredient) is in association with a pharmaceutically acceptable adjuvant, diluent or carrier. Depending on the mode of administration, the pharmaceutical composition will preferably comprise from 0.05 to 99 %w (per cent by weight), more preferably from 0.05 to 80 %w, still more preferably from 0.10 to 70 %w, and even more preferably from 0.10 to 50 %w, of active ingredient, all percentages by weight being based on total composition.

25

The present invention also provides a pharmaceutical composition comprising a compound of formula (I), (I'), (I'') or (I'''), or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore defined, in association with a pharmaceutically acceptable adjuvant, diluent or carrier.

30

The invention further provides a process for the preparation of a pharmaceutical composition of the invention which comprises mixing a compound of formula (I), (I'), (I'') or (I'''), or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore defined, with a pharmaceutically acceptable adjuvant, diluent or carrier.

The pharmaceutical compositions may be administered topically (e.g. to the lung and/or airways or to the skin) in the form of solutions, suspensions, heptafluoroalkane aerosols and dry powder formulations; or systemically, e.g. by oral administration in the form of tablets, capsules, syrups, powders or granules, or by parenteral administration in the form of solutions or suspensions, or by subcutaneous administration or by rectal administration in the form of suppositories or transdermally.

The invention will now be further explained by reference to the following illustrative examples, in which ¹H NMR spectra were recorded on Varian Unity Inova 400. The central solvent peak of chloroform-*d* (δ_H 7.27 ppm) were used as internal standard. Low resolution mass spectra and accurate mass determination were recorded on a Hewlett-Packard 1100 LC-MS system equipped with APCI/ESI ionisation chambers.

All solvents and commercial reagents were laboratory grade and used as received.

The nomenclature used for the compounds was generated with ACD/IUPAC Name Pro.

Example 1

***N*-(2-{3-[3R,S-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2R,S-hydroxy-propoxy}-phenyl)-acetamide hydrochloride**

(i) 3-Hydroxy-pyrrolidine-1-carboxylic acid tert-butyl ester

A solution of pyrrolidin-3-ol (16.25g, 186.5 mmol) and di-tert-butyl-dicarbonate (40.7g, 186.5 mmol) in dry THF (50 ml) under nitrogen was stirred over night. Concentration at reduced pressure and purification by flash chromatography on silica (EtOAc : heptane, 7 : 3) gave 31.9 g (91 %) of the subtitle compound.

¹H-NMR (400MHz, DMSO-*d*₆): δ 4.87 (d, 1H, J = 3.4 Hz), 4.21 (bs, 1H), 3.31-3.22 (m, 3H), 3.10 (d, 1H, J = 11.5 Hz), 1.83 (m, 1H), 1.72 (m, 1H), 1.39 (s, 9H).

APCI-MS: m/z 132 [MH⁺-56]

5 (ii) **3-(4-Chloro-phenoxy)-pyrrolidine**

3-Hydroxy-pyrrolidine-1-carboxylic acid tert butyl ester (2.1g, 9.9 mmol) and triphenyl phosphine (2.59g, 9.9 mmol) were dissolved in dry THF (35 ml) under nitrogen. The solution was cooled to 0°C and 4-chlorophenol (1.28g, 9.9 mmol) dissolved in dry THF (10 ml) was added followed by diethyl azodicarboxylate (DEAD) (1.55 ml, 9.9 mmol).

10 After 15 minutes the ice bath was removed and the reaction was stirred overnight.

The reaction mixture was concentrated under reduced pressure and the residue was stirred with ether. The solid triphenyl phosphine oxide was filtered off. The solution was washed three times with sodium hydroxide (1M) and concentrated. The BOC-protected product was purified by flash chromatography on silica using EtOAc/ heptane as eluant. It was
15 dissolved in dichloromethane (35 ml) and trifluoroacetic acid (17 ml). The reaction mixture was stirred at room temperature overnight, concentrated and purified by flash chromatography on silica (MeOH:CHCl₃:NH₃, 100:100:1) to give the subtitle compound (1.72g, 88%).

20 ¹H-NMR (400MHz, DMSO-*d*₆): δ 7.30 (d, 2H, J = 8.9 Hz), 6.91 (d, 2H, J = 8.9 Hz), 4.82 (m, 1H), 3.03 (dd, 1H, J = 12.3, 5.4 Hz), 2.82 (m, 3H), 1.99 (m, 1H), 1.72 (m, 1H).

APCI-MS: m/z 198 [MH⁺]

25 (iii) **N-(2-{3-[3R,S-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2R,S-hydroxy-propoxy}-phenyl)-acetamide hydrochloride**

A solution of 3-(4-Chloro-phenoxy)-pyrrolidine (0.059 g, 0.298 mmol) and N-acetyl-2-(2,3-epoxypropoxy)aniline (0.062 g, 0.299 mmol) in EtOH (1.5 ml, 99.5%) was stirred for 3 hours at 75°C in a sealed vial. The solvent was evaporated after completion of the reaction and the residue was purified on silica (CH₂Cl₂:MeOH, 98:2 to 97:3) to give 88 mg
30 of the free amine of the title compound. The amine was dissolved in MeOH : water 1:1

(30 ml), and the solution was acidified with 2M hydrochloric acid. The methanol was evaporated and the residual water solution was lyophilized to give 92 mg (70%) of the title compound as a white solid.

5 APCI-MS: m/z 405.2, 407.2 [MH⁺, isotope pattern]

Example 2

***N*-(5-Chloro-2-{3-[3R,S-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2R,S-hydroxy-propoxy}-phenyl)-acetamide hydrochloride**

10

(i) *N*-(5-Chloro-2-hydroxy-phenyl)-acetamide

A solution of 4-amino-2-chlorophenol (2.0g, 13.9 mmol) and acetic anhydride (1.77g, 17.3 mmol) in water (40 ml) was stirred vigorously for 5 minutes. The reaction mixture was then heated with stirring to 60°C for 30 minutes, and was then allowed to cool. A pink solid was formed and the precipitate was collected by filtration, washed twice with water, and dried to give 1.8g (70%) of the subtitle compound.

15

¹H-NMR (400 MHz, DMSO-*d*₆): δ 10.09 (1H, s); 9.25 (1H, bs); 7.93 (1H, s); 6.93 (1H, dd, *J* 8.8, 2.7 Hz); 6.84 (1H, d, *J* 8.6 Hz); 2.09 (3H, s)

20

APCI-MS: m/z 186.0 [MH⁺]

(ii) *N*-(5-Chloro-2-oxiranylmethoxy-phenyl)-acetamide

A solution of *N*-(5-Chloro-2-hydroxy-phenyl)-acetamide (0.499 g, 2.68 mmol), K₂CO₃ (0.60 g, 4.35 mmol) and epibromohydrin (0.405 g, 2.95 mmol) in DMF (5 ml) was heated with stirring at 50°C for 2 hours. The mixture was then partitioned between EtOAc and water 40+40 ml. The organic phase was washed twice with water and once with brine and finally concentrated in vacuo to give a crude product. The crude product was purified on silica (heptane : EtOAc, 1 : 1), to give 0.43 g (66%) of a white solid.

25

¹H-NMR (400MHz, CDCl₃): δ 8.46 (1H, d, *J* 2.3 Hz); 7.90 (1H, bs); 6.98 (1H, dd, *J* 8.7, 2.4 Hz); 6.83 (1H, d, *J* 8.8 Hz); 4.36 (1H, dd, *J* 11.5, 2.4 Hz); 3.94 (1H, dd, *J* 11.6, 6.0 Hz); 3.41-3.36 (1H, m); 2.97 (1H, dd, *J* 4.7, 4.2 Hz); 2.80 (1H, dd, *J* 4.6, 2.6 Hz); 2.23 (3H, s)

5 (iii) ***N*-(5-Chloro-2-{3-[3R,S-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2R,S-hydroxypropoxy}-phenyl)-acetamide hydrochloride**

Prepared by a process analogous to that described in Example 1, step (iii).

Example 3

10 ***N*-(2-{3-[4-(3,4-dichlorophenoxy)-1-piperidinyl]-2-hydroxypropoxy}phenyl)-acetamide**

Prepared according to the methods described in Example 1. Purified and isolated as the free amine in 73% yield by C₁₈-column chromatography (H₂O:CH₃CN, 0.1M NH₄OAc
15 buffer, gradient 30% to 95% CH₃CN).

APCI-MS *m/z*: 453, 455 [MH⁺]

¹H NMR (400 MHz, CDCl₃): δ 7.32(d, 1H), 7.01(d, 1H), 6.85-8.80(m, 2H), 6.78-6.69 (m, 3H), 4.31 (m, 1H), 4.15-4.09(m, 1H), 4.18-3.18(bs, 3H), 2.91(m, 1H), 2.71(m, 1H),
20 2.62-2.52(m, 3H), 2.35(m, 1H), 2.05-1.93(m, 2H), 1.89-1.77(m, 2H)

Example 4

1-(2-aminophenoxy)-3-[4-(3,4-dichlorophenoxy)-1-piperidinyl]-2-propanol dihydrochloride

25 *N*-(2-{3-[4-(3,4-dichlorophenoxy)-1-piperidinyl]-2-hydroxypropoxy}phenyl)acetamide (1.418g, 3.13 mmol) was dissolved in 50ml HCl (35%/aq, puriss) and refluxed overnight. The product precipitated and was filtered and dried to give 0.835 g (65%) of the title compound.

APCI-MS m/z: 411, 413 [MH⁺]

¹H NMR (400 MHz, CDCl₃): δ 8.39-3.31 (m, 2H), 7.31(d, 1H), 7.01-6.98(m, 3H),
6.94-6.91(m, 1H), 6.75(dd, 1H), 4.31(m, 1H), 4.12-4.02 (m, 2H), 3.92(dd, 1H),
2.90(m, 1H), 2.69(m, 1H), 2.62-2.51(m, 2H), 2.46(dd, 1H), 2.34(m, 1H), 2.18(s, 3H),
2.04-1.93(m, 2H), 1.89-1.77(m, 2H).

Example 5

**N-(2-{3-[3-(3,4-dichlorophenoxy)-1-pyrrolidinyl]-2-hydroxypropoxy}phenyl)-
acetamide hydrochloride**

Prepared according to the methods described in Example 1 to give 68mg (68%) of the title
compound as a white solid.

APCI-MS m/z: 439, 441 [MH⁺]

Example 6

**2-{3-[4-(4-Fluoro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-benzoic acid methyl
ester**

(i) 2-Oxiranylmethoxy-benzoic acid methyl ester

Prepared according to the method described in Example 2, step (ii).

¹H-NMR: (400 MHz, CDCl₃): δ 7.81 (1H, dd, *J* 7.7 , 1.7Hz); 7.46 (1H, dt, *J* 7.7 , 1.7 Hz);
7.05-6.98 (2H, m); 4.33 (1H, dd, *J* 11.3 , 3.0 Hz); 4.11 (1H, dd, *J* 11.3 , 4.8 Hz); 3.90
(3H, s); 3.43-3.37 (1H, m); 2.93-2.90 (2H, m)

**(ii) 2-{3-[4-(4-Fluoro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-benzoic acid
methyl ester**

Prepared according to the method described in Example 1, step (ii). Isolated as the free
amine

¹H-NMR: (400 MHz, CDCl₃): δ 7.81 (1H, dd, *J* 8.1 , 1.8 Hz); 7.46 (1H, dt, *J* 7.8 , 1.7 Hz); 7.03-6.91 (4H, m); 6.86-6.82 (2H, m); 4.28-4.10 (3H, m); 4.08-4.00 (1H, m); 3.88 (3H, s); 2.92-2.84 (1H, m); 2.83-2.76 (1H, m); 2.66-2.53 (2H, m); 2.46 (1H, t, *J* 10.2 Hz); 2.36 (1H, t, *J* 10.2 Hz); 2.02-1.92 (2H, m); 1.86-1.74 (2H, m); 1.63 (1H, bs)
APCI-MS: m/z 404.2 [MH⁺]

Example 7

2-(2-{3-[4-(3,4-Difluoro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-benzoylamino)-2-methyl-propionic acid methyl ester

(i) Methyl 2-[(2-hydroxybenzoyl)amino]-2-methylpropanoate

To a solution of 2-(chlorocarbonyl)phenyl acetate (20 mmol, 3.96g) in toluene (50ml) were *N*-ethyl-*N,N*-diisopropylamine (22 mmol, 2.84g) and 2-methylalanine (22 mmol, 2.27g) added. After stirring the reaction mixture at room temperature overnight, the mixture was diluted with 250ml toluene and was washed with 1.8% HCl/aq (250 ml) and sat. NaCl/aq (250 ml). The organic phase was dried over Na₂SO₄ and concentrated under reduced pressure. The residue was dissolved in MeOH (50mL) and 3 drops of conc. H₂SO₄ were added. The mixture was refluxed for 2 hours and concentrated under reduced pressure. The residue was dissolved in 250mL EtOAc and washed with sat. NaHCO₃/aq (250 ml) and sat. NaCl/aq (250 ml). The organic phase was dried over Na₂SO₄ and concentrated at reduced pressure. The resulting crude material was used without further purification.
APCI-MS m/z: 238 [MH⁺]

(ii) Methyl 2-methyl-2-{[2-(2-oxiranylmethoxy)benzoyl]amino}propanoate

A solution of methyl 2-[(2-hydroxybenzoyl)amino]-2-methylpropanoate, K₂CO₃ (20 mmol, 2.68g) and 2-(chloromethyl)oxirane (22 mmol, 2.03g) in acetonitrile (60 ml) was stirred at reflux temperature overnight. The reaction mixture was diluted with EtOAc and washed with 1.8% HCl/aq (250 ml) and sat. NaCl/aq (250 ml). The organic phase was dried over Na₂SO₄ and concentrated at reduced pressure. The residue was purified on C₁₈-column

(H₂O:CH₃CN, 0.1M NH₄OAc buffer, gradient 10% to 95% CH₃CN) to give the subtitle compound (244mg, 5% yield, two steps).

APCI-MS m/z: 294 [MH⁺]

¹H NMR (400 MHz, CDCl₃) : δ 8.40(s, 1H), 8.14 (dd, 1H), 7.41 (dt, 1H), 7.07 (t, 1H), 6.90 (d, 1H), 4.44 (dd, 1H), 4.07 (dd, 1H), 3.74 (s, 3H), 3.45 (m, 1H), 2.94 (dd, 1H), 2.84 (dd, 1H), 1.64 (d, 6H)

(iii) **2-(2-{3-[4-(3,4-Difluoro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-benzoylamino)-2-methyl-propionic acid methyl ester**

A toluene solution of 4-(3,4-difluorophenoxy)piperidine (0.03 ml, 0.5 M) was mixed with a toluene solution of methyl 2-methyl-2-[[2-(2-oxiranylmethoxy)benzoyl]amino]-propanoate (0.03 ml, 0.5 M). The mixture was diluted with 0.20 ml toluene and 0.05 ml methanol. The reaction mixture was stirred overnight at 100° C in sealed vials. The product were concentrated in vacuo and used without any purification.

APCI-MS m/z: 507 [MH⁺]

¹H NMR (400 MHz, CDCl₃) : δ 8.13(s, 1H), 7.90 (dd, 1H), 7.33 (dt, 1H), 7.07-6.96 (m, 2H), 6.89 (d, 1H), 6.73-6.68 (m, 1H), 6.58-6.55 (m, 1H), 4.77-4.72 (m, 1H), 4.49 (bs, 1H), 4.20-4.13 (m, 2H), 3.69 (s, 3H), 3.58-3.44 (m, 2H), 3.39-3.26 (m, 4H), 2.54-2.40 (m, 2H), 2.13-2.04 (m, 2H), 1.60 (d, 6H)

Example 8

***N*-[2-(((1*R*,2*S*,3*R*)*-3-[4-(3,4-dichlorophenoxy)-1-piperidinyl]-2-hydroxycyclopentyl)oxy)phenyl]acetamide and *N*-[2-(((1*S*,2*S*,3*R*)*-3-[4-(3,4-dichlorophenoxy)-1-piperidinyl]-2-hydroxycyclopentyl)oxy)phenyl]acetamide**

(i) ***N*-[2-(2-cyclopenten-1-yloxy)phenyl]acetamide**

To a suspension of sodium hydride (60 proc. in paraffin; 297 mg, 7.43 mmol, 1.1 equiv.) in DMF (3 ml) a solution of 2-acetamido-phenol (1.02 g, 6.75 mmol, 1.0 equiv.) in DMF (12 ml) was added dropwise at 0°C. After 30 minutes chlorocyclopent-2-ene (R. Moffett, *Organic Synthesis*, Wiley: New York 1963, Collect. Vol. IV, p.238-241) (762 mg, 0.76 ml, 7.43 mmol, 1.1 equiv.) was added by a syringe and stirring was continued overnight. Aqueous work-up followed by flash chromatography on silica gel (heptane/ethyl acetate, 2:1 continued to 1:1) afforded 992 mg (68 %) of the subtitle compound as a dark, yellow oil.

¹H-NMR (400 MHz, CDCl₃): δ 8.35 (1H, d, J 8.0Hz), 7.73 (1H, bs), 7.00 (1H, td, J 7.9, 1.5Hz), 6.90-6.95 (2H, m), 6.17 (1H, m), 5.95 (1H, m), 5.36 (1H, d, J 5.9Hz), 2.59 (1H, m), 2.38 (2H, m), 2.17 (3H, s), 1.97 (1H, m).

MS-ESI+ : m/z 218.1 [MH⁺].

(ii) *N*-[2-(6-oxabicyclo[3.1.0]hex-2-yloxy)phenyl]acetamide

To an ice bath cooled solution of *N*-[2-(2-cyclopenten-1-yloxy)phenyl]acetamide (149 mg, 686 μmol, 1.0 equiv.) in dichloromethane (4 ml) *m*-chloroperbenzoic acid (85 proc.; 146 μmol, 1.1 equiv.) was added. After stirring overnight with slowly warming up to an ambient temperature the reaction mixture was diluted by *tert*butyl(methyl)ether, washed successively by a sat. sodium bisulfate solution, 5 proc. sodium hydroxide and brine and dried over sodium sulfate. Evaporation of the solvent and flash chromatography on silica gel (ethyl acetate/heptane, 2:3 continued to ethyl acetate) yielded 93 mg (58 %) of the subtitle compound as a mixture of the *trans*, (minor) and the *cis* (major) diastereoisomeric epoxides as a pale yellow oil. The *cis/trans* ratio was determined as 2:1 by ¹H-NMR.

¹H-NMR (400 MHz, CDCl₃): δ 8.39 (1H [A], m), 8.34 (1H [B], d, J 8.2Hz), 7.91 (1H [A], bs), 7.59 (1H, [B], bs), 6.92-7.25 (3H [A] + 3H [B], m), 4.89 (1H [B], d, J 5.2Hz), 4.77 (1H [A], td, J 8.0, 1.3Hz), 3.66 (1H [B], m), 3.64 (1H [B], m), 3.60 (1H, [A], m), 3.54 (1H [A], m), 2.23 (1H [B], d, J 8.4Hz), 2.21 (3H [A], s), 2.19 (3H [B], s), 2.10 (2H [A], m), 1.72-1.92 (m), 1.53-1.63 (m) (2H [A] + 3H [B]). (A=*trans*, B=*cis*)

MS-ESI+: m/z 234.1 [MH+].

(iii) *N*-[2-(((1*R*,2*S*,3*R*)*-3-[4-(3,4-dichlorophenoxy)-1-piperidiny]-2-hydroxycyclopentyl}oxy)phenyl]acetamide and

N-[2-(((1*S*,2*S*,3*R*)*-3-[4-(3,4-dichlorophenoxy)-1-piperidiny]-2-hydroxycyclopentyl}oxy)phenyl]acetamide

N-{2-(6-oxabicyclo[3.1.0]hex-2-yloxy)phenyl}acetamide (racemic mixture of the trans and cis diastereoisomers) (87 mg, 373 μ mol, 1.0 equiv.) and 4-(3,4-dichlorophenoxy)piperidine (92 mg, 394 μ mol, 1.06 equiv.) were dissolved in 2 M lithium perchlorate in acetonitrile (3 ml) and heated in a sealed tube over night at 85 °C. Aqueous work-up and flash chromatography of the crude on silica gel (heptane/ethyl acetate/methanol/ammonia = 1:3:0:0 continued to 0:90:10:1 to 0:80:20:3) led to the separation of two diastereoisomeric addition products to give 24 mg (14 %) of the (1*S*,2*S*,3*R*) diastereoisomer (first eluted) and 75 mg (42 %) of the second eluted (1*R*,2*S*,3*R*) diastereoisomer.

For (1*S*,2*S*,3*R*) diastereoisomer:

¹H-NMR (400 MHz, CDCl₃): δ 8.27 (1H, dd, J 7.6, 1.7Hz), 7.91 (1H, s), 7.29 (1H, d, J 8.9Hz), 6.88-7.00 (4H, m), 6.73 (1H, dd, J 8.9, 2.8Hz), 4.45 (1H, m), 4.28 (1H, hept, J 3.6Hz), 4.18 (1H, dd, J 7.1, 4.6Hz), 2.87 (3H, m), 2.71 (1H, q, J 7.5Hz), 2.15 (3H, s), 2.11 (1H, m), 1.78-2.02 (7H, m).

MS-APCI+: m/z 479.1 [MH+].

For (1*R*,2*S*,3*R*) diastereoisomer:

¹H-NMR (400 MHz, CDCl₃): δ 8.20-8.25 (2H, m), 7.29 (1H, d, J 8.9Hz), 6.91-7.00 (4H, m), 6.74 (1H, dd, J 8.9, 2.8Hz), 4.46 (1H, bq, J 4.8Hz), 4.29 (1H, m), 4.13 (1H, d, J 7.2Hz), 2.95 (2H, m), 2.84 (2H, m), 2.50 (2H, m), 2.15 (3H, s), 1.93-2.07 (5H, m), 1.82 (2H, m), 1.58 (1H, m).

MS-APCI+ : m/z 479.1 [MH+].

Example 9

***N*-[2-((2,3-*trans*)-3-[4-(3,4-dichlorophenoxy)-1-piperidinyl]-2-hydroxycyclohexyl)oxy)phenyl]acetamide**

(i) *N*-[2-(2-cyclohexen-1-yloxy)phenyl]acetamide

2-Cyclohexenol (491 mg, 0.49 ml, 5.00 mmol, 1.0 equiv.), 2-acetamidophenol (756 mg, 5.00 mmol, 1.0 equiv.) and triphenylphosphine (1.44 g, 5.50 mmol, 1.1 equiv.) were dissolved in THF (10 ml) and kept at ambient temperature by a water bath. After dropwise addition of diethyl azodicarbonic acid (871 mg, 0.78 ml, 5.00 mmol, 1.0 equiv.), dissolved in THF (3 ml), the reaction mixture was stirred over night. Extractive work-up and flash chromatography on silica gel (heptane/*tert*butyl(methyl)ether = 1:1) afforded 224 mg (19 %) the title compound as a yellow oil.

MS-ESI+: m/z 232.2 [MH⁺].

(ii) *N*-[2-(7-oxabicyclo[4.1.0]hept-2-yloxy)phenyl]acetamide

To a solution of *N*-[2-(2-cyclohexen-1-yloxy)phenyl]acetamide (76 mg, 329 μ mol, 1.0 equiv.) in dichloromethane (5 ml) *m*-chlorobenzoic acid (85 proz.; 121 mg, 559 μ mol, 1.7 equiv.) was added at 0°C. Stirring was continued overnight while the reaction mixture was allowed to warm up slowly to room temperature. The heterogenous mixture was diluted with ethyl acetate and washed with sat. sodium sulfite, 5 % sodium hydroxide and brine. Drying over sodium sulfate, evaporation of the solvent and flash chromatography on silica gel provided 59 mg (73 %) of the title compound as a mixture of the diastereoisomers (ratio A:B = *trans*:*cis* = 5:3 [¹H-NMR]).

¹H-NMR (400 MHz, CDCl₃): δ 8.35 (1H [A] + 1H [B], m), 8.02 (1H [A], bs), 7.70 (1H [B], bs), 6.95-7.04 (3H [A] + 3H [B], m), 4.62 (1H [A], dd, *J* 8.4, 5.5, 2.1 Hz), 4.55 (1H [B], dd, *J* 7.5, 6.7 Hz), 3.30-3.36 (2H [A] + 1H [B], m), 3.19 (1H [B], t, *J* 3.6 Hz), 1.26-2.23 (10H [A] + 10H [B], m).

LC/MS-ESI+: m/z 248.1 [MH⁺ (A)], 248.2 [MH⁺ (B)].

(iii) *N*-[2-((2,3-*trans*)-3-[4-(3,4-dichlorophenoxy)-1-piperidinyl]-2-hydroxycyclohexyl)oxy)phenyl]acetamide

The diastereoisomeric mixture of *N*-[2-(7-oxabicyclo[4.1.0]hept-2-yloxy)phenyl]acetamide (59 mg, 239 μ mol, 1.0 equiv.) and 4-(3,4-dichlorophenoxy)piperidine (56 mg, 239 μ mol, 1.0 equiv.) were dissolved in 2 M lithium perchlorate in acetonitrile (2 ml) and heated in a sealed tube over night at 85 °C. Aqueous work-up and flash chromatography on silica gel (heptane/ethyl acetate/methanol = 50:100:3) gave 86 mg (75 %) as a yellow oil in a diastereoisomeric ratio of 69:31 = A:B (¹H-NMR). No separation of the diastereoisomers on reversed phase columns could be observed. The relative stereochemistry of the major and minor diastereoisomers, respectively, could not be assigned due to the complex spectrum of the mixture.

¹H-NMR (400 MHz, CDCl₃): δ 9.48 (1H [A], bs), 9.25 (1H [B], bs), 8.46 (1H [A] + 1H [B], t, J 9.1 Hz), 7.22-7.32 (2H [A] + 1H [B], m), 6.93-7.08 (4H [A] + 5H [B], m), 6.72-6.76 (1H [A] + 1H [B], m), 4.08-4.30 (3H [A] + 3H [B], m), 3.55-3.64 (2H [A] + 1H [B], m), 2.96-3.07 (2H [A] + 2H [B], m), 2.71 (2H [A] + 3H [B], m), 2.19 (3H [A], s), 2.16 (3H [B], s), 1.47-2.37 (10 H [A] + 10H [B], m).

MS-ESI+: m/z 493.1 [MH⁺ (A, B)].

The following compounds were prepared by routes analogous to those described in the previous Examples.

Example 10

N-(5-Chloro-2-{3-[3-(3,4-dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide

APCI-MS: m/z 473.1 475.1 [MH⁺]

Example 11

N-(3-Acetyl-2-{3-[3-(3,4-dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-5-methyl-phenyl)-acetamide

APCI-MS: m/z 495.1 497.1 [MH⁺]

5

Example 12

N-(2-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-4-methyl-phenyl)-acetamide

10 APCI-MS: m/z 453.1 455.1 [MH⁺]

Example 13

N-(2-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-5-fluoro-phenyl)-acetamide

15

APCI-MS: m/z 457.1 459.1 [MH⁺]

Example 14

1-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-3-(1H-indol-7-yloxy)-propan-2-ol

20

APCI-MS: m/z 421.1 423.1 [MH⁺]

Example 15

1-(7-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-indol-1-yl)-ethanone

25

APCI-MS: m/z 463.1 465.1 [MH⁺]

Example 16

N-(4-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-biphenyl-3-yl)-acetamide

APCI-MS: m/z 515.1 517.1 [MH⁺]

5

Example 17

N-(2-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-4-fluorophenyl)-acetamide

10 APCI-MS: m/z 457.1 459.1 [MH⁺]

Example 18

N-(2-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-5-methylphenyl)-acetamide

15

APCI-MS: m/z 453.1 455.1 [MH⁺]

Example 19

N-(2-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide

20

APCI-MS: m/z 439.1 441.1 [MH⁺]

Example 20

N-(5-Chloro-2-{3-[3-(4-chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide

25

APCI-MS: m/z 439.1 441.1 [MH⁺]

30 **Example 21**

N-(3-Acetyl-2-{3-[3-(4-chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-5-methyl-phenyl)-acetamide

APCI-MS: m/z 461.1 [MH⁺]

5

Example 22

N-(2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-4-methyl-phenyl)-acetamide

10 APCI-MS: m/z 419.1 [MH⁺]

Example 23

N-(2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-5-fluoro-phenyl)-acetamide

15

APCI-MS: m/z 423.1 [MH⁺]

Example 24

1-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-3-(1H-indol-7-yloxy)-propan-2-ol

20

APCI-MS: m/z 387.1 [MH⁺]

Example 25

1-(7-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-indol-1-yl)-ethanone

25

APCI-MS: m/z 429.1 [MH⁺]

Example 26

N-(4-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-biphenyl-3-yl)-acetamide

APCI-MS: m/z 481.1 [MH⁺]

5

Example 27

N-(2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-4-fluorophenyl)-acetamide

10 APCI-MS: m/z 423.1 [MH⁺]

Example 28

N-(2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-5-methylphenyl)-acetamide

15

APCI-MS: m/z 419.1 [MH⁺]

Example 29

N-(2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide

20

APCI-MS: m/z 405.1 [MH⁺]

Example 30

N-(5-Chloro-2-{3-[3-(4-fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide

25

APCI-MS: m/z 423.1 [MH⁺]

30

Example 31

N-(3-Acetyl-2-{3-[3-(4-fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-5-methyl-phenyl)-acetamide

APCI-MS: m/z 445.3 [MH⁺]

Example 32

N-(2-{3-[3-(4-Fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-4-methyl-phenyl)-acetamide

APCI-MS: m/z 403.3 [MH⁺]

Example 33

N-(5-Fluoro-2-{3-[3-(4-fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide

APCI-MS: m/z 407.1 [MH⁺]

Example 34

1-[3-(4-Fluoro-phenoxy)-pyrrolidin-1-yl]-3-(1H-indol-7-yloxy)-propan-2-ol

APCI-MS: m/z 371.1 [MH⁺]

Example 35

1-(7-{3-[3-(4-Fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-indol-1-yl)-ethanone

APCI-MS: m/z 413.1 [MH⁺]

Example 36

N-(4-{3-[3-(4-Fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-biphenyl-3-yl)-acetamide

APCI-MS: m/z 465.3 [MH⁺]

5

Example 37

N-(4-Fluoro-2-{3-[3-(4-fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide

10 APCI-MS: m/z 407.1 [MH⁺]

Example 38

N-(2-{3-[3-(4-Fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-5-methyl-phenyl)-acetamide

15

APCI-MS: m/z 403.1 [MH⁺]

Example 39

N-(2-{3-[3-(4-Fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide

20

APCI-MS: m/z 389.1 [MH⁺]

Example 40

N-(5-Chloro-2-{3-[3-(3,4-difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide

25

APCI-MS: m/z 441.1 [MH⁺]

30 **Example 41**

N-(3-Acetyl-2-{3-[3-(3,4-difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-5-methyl-phenyl)-acetamide

APCI-MS: m/z 463.3 [MH⁺]

5

Example 42

N-(2-{3-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-4-methyl-phenyl)-acetamide

10 APCI-MS: m/z 421.1 [MH⁺]

Example 43

N-(2-{3-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-5-fluoro-phenyl)-acetamide

15

APCI-MS: m/z 425.1 [MH⁺]

Example 44

1-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-3-(1H-indol-7-yloxy)-propan-2-ol

20

APCI-MS: m/z 389.1 [MH⁺]

Example 45

1-(7-{3-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-indol-1-yl)-ethanone

25

APCI-MS: m/z 431.1 [MH⁺]

Example 46

N-(4-{3-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-biphenyl-3-yl)-acetamide

APCI-MS: m/z 483.3 [MH⁺]

5

Example 47

N-(2-{3-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-4-fluorophenyl)-acetamide

10 APCI-MS: m/z 425.1 [MH⁺]

Example 48

N-(2-{3-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-5-methylphenyl)-acetamide

15

APCI-MS: m/z 421.1 [MH⁺]

Example 49

N-(2-{3-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide

20

APCI-MS: m/z 407.1 [MH⁺]

Example 50

25 N-(5-Chloro-2-{3-[4-(3,4-dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide

APCI-MS: m/z 487.1 489.1 [MH⁺]

30 **Example 51**

N-(3-Acetyl-2-{3-[4-(3,4-dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-5-methyl-phenyl)-acetamide

APCI-MS: m/z 509.1 511.1 [MH⁺]

5

Example 52

N-(2-{3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-4-methyl-phenyl)-acetamide

10 APCI-MS: m/z 467.1 469.1 [MH⁺]

Example 53

N-(2-{3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-5-fluoro-phenyl)-acetamide

15

APCI-MS: m/z 471.1 473.1 [MH⁺]

Comparison Example 54

1-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-3-(1H-indol-7-yloxy)-propan-2-ol

20

APCI-MS: m/z 435.1 437.1 [MH⁺]

Example 55

1-(7-{3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-indol-1-yl)-ethanone

25

APCI-MS: m/z 477.1 479.1 [MH⁺]

Example 56

N-(4-{3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-biphenyl-3-yl)-acetamide

APCI-MS: m/z 429.1 431.1 [MH⁺]

5

Example 57

N-(2-{3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-4-fluorophenyl)-acetamide

10 APCI-MS: m/z 471.1 473.1 [MH⁺]

Example 58

N-(2-{3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-5-methylphenyl)-acetamide

15

APCI-MS: m/z 467.1 469.1 [MH⁺]

Example 59

N-(2-{3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide

20

APCI-MS: m/z 453.1 455.1 [MH⁺]

Example 60

N-(5-Chloro-2-{3-[4-(4-chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide

25

APCI-MS: m/z 453.1 455.1 [MH⁺]

30 Example 61

N-(3-Acetyl-2-{3-[4-(4-chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-5-methyl-phenyl)-acetamide

APCI-MS: m/z 475.3 [MH⁺]

5

Example 62

N-(2-{3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-4-methyl-phenyl)-acetamide

10 APCI-MS: m/z 433.1 [MH⁺]

Example 63

N-(2-{3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-5-fluoro-phenyl)-acetamide

15

APCI-MS: m/z 437.1 [MH⁺]

Comparison Example 64

1-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-3-(1H-indol-7-yloxy)-propan-2-ol

20

APCI-MS: m/z 401.1 [MH⁺]

Example 65

1-(7-{3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-indol-1-yl)-ethanone

25

APCI-MS: m/z 443.1 [MH⁺]

Example 66

N-(4-{3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-biphenyl-3-yl)-acetamide

APCI-MS: m/z 495.3 [MH⁺]

5

Example 67

N-(2-{3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-4-fluoro-phenyl)-acetamide

10 APCI-MS: m/z 437.1 [MH⁺]

Example 68

N-(2-{3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-5-methyl-phenyl)-acetamide

15

APCI-MS: m/z 433.1 [MH⁺]

Example 69

N-(2-{3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide

20

APCI-MS: m/z 419.1 [MH⁺]

Example 70

N-{5-Chloro-2-[3-(8-chloro-1,3,4,5-tetrahydro-pyrido[4,3-b]indol-2-yl)-2-hydroxy-propoxy]-phenyl}-acetamide

25

APCI-MS: m/z 448.1 450.1 [MH⁺]

30 **Example 71**

N-{3-Acetyl-2-[3-(8-chloro-1,3,4,5-tetrahydro-pyrido[4,3-b]indol-2-yl)-2-hydroxy-propoxy]-5-methyl-phenyl}-acetamide

APCI-MS: m/z 470.1 [MH⁺]

5

Example 72

N-{2-[3-(8-Chloro-1,3,4,5-tetrahydro-pyrido[4,3-b]indol-2-yl)-2-hydroxy-propoxy]-4-methyl-phenyl}-acetamide

10 APCI-MS: m/z 428.1 [MH⁺]

Example 73

N-{2-[3-(8-Chloro-1,3,4,5-tetrahydro-pyrido[4,3-b]indol-2-yl)-2-hydroxy-propoxy]-5-fluoro-phenyl}-acetamide

15

APCI-MS: m/z 432.1 [MH⁺]

Example 74

1-(8-Chloro-1,3,4,5-tetrahydro-pyrido[4,3-b]indol-2-yl)-3-(1H-indol-7-yloxy)-propan-2-ol

20

APCI-MS: m/z 396.1 [MH⁺]

Example 75

1-{7-[3-(8-Chloro-1,3,4,5-tetrahydro-pyrido[4,3-b]indol-2-yl)-2-hydroxy-propoxy]-indol-1-yl}-ethanone

25

APCI-MS: m/z 438.1 [MH⁺]

30 **Example 76**

N-{4-[3-(8-Chloro-1,3,4,5-tetrahydro-pyrido[4,3-b]indol-2-yl)-2-hydroxy-propoxy]-biphenyl-3-yl}-acetamide

APCI-MS: m/z 490.1 [MH⁺]

5

Example 77

N-{2-[3-(8-Chloro-1,3,4,5-tetrahydro-pyrido[4,3-b]indol-2-yl)-2-hydroxy-propoxy]-4-fluoro-phenyl}-acetamide

10 APCI-MS: m/z 432.1 [MH⁺]

Example 78

N-{2-[3-(8-Chloro-1,3,4,5-tetrahydro-pyrido[4,3-b]indol-2-yl)-2-hydroxy-propoxy]-5-methyl-phenyl}-acetamide

15

APCI-MS: m/z 428.1 [MH⁺]

Example 79

N-{2-[3-(8-Chloro-1,3,4,5-tetrahydro-pyrido[4,3-b]indol-2-yl)-2-hydroxy-propoxy]-phenyl}-acetamide

20

APCI-MS: m/z 414.1 [MH⁺]

Example 80

25 N-{5-Chloro-2-[3-(8-fluoro-1,3,4,5-tetrahydro-pyrido[4,3-b]indol-2-yl)-2-hydroxy-propoxy]-phenyl}-acetamide

APCI-MS: m/z 432.1 [MH⁺]

30 **Example 81**

N-{3-Acetyl-2-[3-(8-fluoro-1,3,4,5-tetrahydro-pyrido[4,3-b]indol-2-yl)-2-hydroxy-propoxy]-5-methyl-phenyl}-acetamide

APCI-MS: m/z 454.3 [MH⁺]

5

Example 82

N-{2-[3-(8-Fluoro-1,3,4,5-tetrahydro-pyrido[4,3-b]indol-2-yl)-2-hydroxy-propoxy]-4-methyl-phenyl}-acetamide

10 APCI-MS: m/z 412.1 [MH⁺]

Example 83

N-{5-Fluoro-2-[3-(8-fluoro-1,3,4,5-tetrahydro-pyrido[4,3-b]indol-2-yl)-2-hydroxy-propoxy]-phenyl}-acetamide

15

APCI-MS: m/z 416.1 [MH⁺]

Example 84

1-(8-Fluoro-1,3,4,5-tetrahydro-pyrido[4,3-b]indol-2-yl)-3-(1H-indol-7-yloxy)-propan-2-ol

20

APCI-MS: m/z 380.1 [MH⁺]

Example 85

1-{7-[3-(8-Fluoro-1,3,4,5-tetrahydro-pyrido[4,3-b]indol-2-yl)-2-hydroxy-propoxy]-indol-1-yl}-ethanone

25

APCI-MS: m/z 422.1 [MH⁺]

30 **Example 86**

N-{4-[3-(8-Fluoro-1,3,4,5-tetrahydro-pyrido[4,3-b]indol-2-yl)-2-hydroxy-propoxy]-biphenyl-3-yl}-acetamide

APCI-MS: m/z 474.3 [MH⁺]

5

Example 87

N-{4-Fluoro-2-[3-(8-fluoro-1,3,4,5-tetrahydro-pyrido[4,3-b]indol-2-yl)-2-hydroxy-propoxy]-phenyl}-acetamide

10 APCI-MS: m/z 416.1 [MH⁺]

Example 88

N-{2-[3-(8-Fluoro-1,3,4,5-tetrahydro-pyrido[4,3-b]indol-2-yl)-2-hydroxy-propoxy]-5-methyl-phenyl}-acetamide

15

APCI-MS: m/z 412.1 [MH⁺]

Example 89

N-{2-[3-(8-Fluoro-1,3,4,5-tetrahydro-pyrido[4,3-b]indol-2-yl)-2-hydroxy-propoxy]-phenyl}-acetamide

20

APCI-MS: m/z 398.1 [MH⁺]

Example 90

N-(2-{3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide

25

APCI-MS m/z: 453, 455 [MH⁺]

30 **Example 91**

3-(2-{3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-
propionic acid methyl ester

APCI-MS m/z: 482, 484 [MH+]

5

Example 92

1-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-3-(2,6-dimethoxy-phenoxy)-propan-2-ol

APCI-MS m/z: 456, 458 [MH+]

10

Example 93

1-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-3-(2-methoxy-phenoxy)-propan-2-ol

APCI-MS m/z: 426, 428 [MH+]

15

Example 94

2-{3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-N,N-dimethyl-
benzamide

20 APCI-MS m/z: 467, 469 [MH+]

Example 95

1-(2-{3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-
propan-1-one

25

APCI-MS m/z: 452, 454 [MH+]

Example 96

1-(2-{3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-
ethanone

30

APCI-MS m/z: 438, 440 [MH+]

Example 97

5 **3-(2-{3-[3-(4-Fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-
propionic acid methyl ester**

APCI-MS m/z: 418 [MH+]

10 **Example 98**

1-(2,6-Dimethoxy-phenoxy)-3-[3-(4-fluoro-phenoxy)-pyrrolidin-1-yl]-propan-2-ol

APCI-MS m/z: 392 [MH+]

15 **Example 99**

1-[3-(4-Fluoro-phenoxy)-pyrrolidin-1-yl]-3-(2-methoxy-phenoxy)-propan-2-ol

APCI-MS m/z: 362 [MH+]

20 **Example 100**

**(2-{3-[3-(4-Fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-benzoylamino)-
acetic acid methyl ester**

APCI-MS m/z: 447 [MH+]

25

Example 101

**(2-{3-[3-(4-Chloro-phenoxy)methyl]-piperidin-1-yl]-2-hydroxy-propoxy}-
benzoylamino)-acetic acid methyl ester**

30 APCI-MS m/z: 491 [MH+]

Example 102

2-(2-{3-[3-(4-Fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-benzoylamino)-2-methyl-propionic acid methyl ester

5

APCI-MS m/z: 475 [MH+]

Example 103

2-{3-[3-(4-Fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-N,N-dimethyl-benzamide

10

APCI-MS m/z: 403 [MH+]

Example 104

1-(2-{3-[3-(4-Fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-6-methoxy-phenyl)-ethanone

15

APCI-MS m/z: 404 [MH+]

Example 105

1-(2-{3-[3-(4-Fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propan-1-one

20

APCI-MS m/z: 388 [MH+]

25

Example 106

1-(2-{3-[3-(4-Fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-ethanone

APCI-MS m/z: 374 [MH+]

30

Example 107

***N*-[2-(3-{[1-(3,4-dichlorobenzyl)-4-piperidinyl]amino}-2-hydroxypropoxy)-4-methylphenyl]acetamide**

5 APCI-MS: m/z 480[MH⁺]

Example 108

3-(2-{3-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propionic acid methyl ester

10

APCI-MS m/z: 436 [MH⁺]

Example 109

1-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-3-(2-methoxy-phenoxy)-propan-2-ol

15

APCI-MS m/z: 380 [MH⁺]

Example 110

(2-{3-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-benzoylamino)-acetic acid methyl ester

20

APCI-MS m/z: 465 [MH⁺]

Example 111

2-{3-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-N,N-dimethylbenzamide

25

APCI-MS m/z: 421 [MH⁺]

30

Example 112

1-(2-{3-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-6-methoxy-phenyl)-ethanone

APCI-MS m/z: 422 [MH⁺]

5

Example 113

1-(2-{3-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propan-1-one

10 APCI-MS m/z: 406 [MH⁺]

Example 114

1-(2-{3-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-ethanone

15

APCI-MS m/z: 392 [MH⁺]

Example 115

N-(2-{3-[4-(3,4-dichloroanilino)-1-piperidinyl]-2-hydroxypropoxy}phenyl)acetamide

20

APCI-MS: m/z 452.1 [MH⁺]

Example 116

3-(2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propionic acid methyl ester

25

APCI-MS m/z: 434 [MH⁺]

Example 117

1-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-3-(2,6-dimethoxy-phenoxy)-propan-2-ol

30

APCI-MS m/z: 408 [MH+]

Example 118

5 **1-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-3-(2-methoxy-phenoxy)-propan-2-ol**

APCI-MS m/z: 378 [MH+]

Example 119

10 **(2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-benzoylamino)-
acetic acid, methyl ester**

APCI-MS m/z: 463 [MH+]

15 **Example 120**

**2-(2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-benzoylamino)-2-
methyl-propionic acid methyl ester**

APCI-MS m/z: 491 [MH+]

20

Example 121

**2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-N,N-dimethyl-
benzamide**

25 APCI-MS m/z: 419 [MH+]

Example 122

**1-(2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-6-methoxy-
phenyl)-ethanone**

30

APCI-MS m/z: 420 [MH+]

Example 123

1-(2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propan-
5 1-one

APCI-MS m/z: 404 [MH+]

Example 124

10 1-(2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-ethanone

APCI-MS m/z: 390 [MH+]

Example 125

15 N-(2-{3-[3-(4-Cyano-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-
acetamide

APCI-MS m/z: 396 [MH+]

20 **Example 126**

3-(2-{3-[3-(4-Cyano-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propionic
acid methyl ester

APCI-MS m/z: 425 [MH+]

25

Example 127

(2-{3-[3-(4-Cyano-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-benzoylamino)-
acetic acid methyl ester

30 APCI-MS m/z: 454 [MH+]

Example 128

2-{3-[3-(4-Cyano-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-N,N-dimethyl-benzamide

APCI-MS m/z: 410 [MH⁺]

Example 129

4-{1-[2-Hydroxy-3-(2-propionyl-phenoxy)-propyl]-pyrrolidin-3-yloxy}-benzonitrile

APCI-MS m/z: 395 [MH⁺]

Example 130

N-(2-{2-Hydroxy-3-[3-(4-methoxy-phenoxy)-pyrrolidin-1-yl]-propoxy}-phenyl)-acetamide

APCI-MS m/z: 401 [MH⁺]

Example 131

N-(4-chloro-2-{3-[4-(3,4-dichloroanilino)-1-piperidinyl]-2-hydroxypropoxy}phenyl)acetamide

APCI-MS: m/z 486 [MH⁺]

Example 132

3-(2-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propionic acid methyl ester

APCI-MS m/z: 468, 470 [MH⁺]

Example 133

1-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-3-(2-methoxy-phenoxy)-propan-2-ol

APCI-MS m/z: 412, 414 [MH+]

5

Example 134

(2-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-benzoylamino)-
acetic acid methyl ester

10 APCI-MS m/z: 497, 499 [MH+]

Example 135

2-(2-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-
benzoylamino)-2-methyl-propionic acid methyl ester

15

APCI-MS m/z: 525, 527 [MH+]

Example 136

2-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-N,N-dimethyl-
20 benzamide

APCI-MS m/z: 453, 455 [MH+]

Example 137

25 1-(2-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-6-methoxy-
phenyl)-ethanone

APCI-MS m/z: 454, 456 [MH+]

30 **Example 138**

1-(2-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propan-1-one

APCI-MS m/z: 438, 440 [MH+]

5

Example 139

1-(2-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-ethanone

10 APCI-MS m/z: 424, 426 [MH+]

Example 140

N-(2-{3-[4-(3,4-Difluoro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide

15

APCI-MS m/z: 421 [MH+]

Example 141

3-(2-{3-[4-(3,4-Difluoro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propionic acid methyl ester

20

APCI-MS m/z: 450 [MH+]

Example 142

2-{3-[4-(3,4-Difluoro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-N,N-dimethylbenzamide

25

APCI-MS m/z: 435 [MH+]

30 **Example 143**

1-(2-{3-[4-(3,4-Difluoro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propan-1-one

APCI-MS m/z: 420 [MH+]

5

Example 144

(2-{3-[4-(3,4-Difluoro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-benzoylamino)-acetic acid methyl ester

10 APCI-MS m/z: 479 [MH+]

Example 145

N-(2-{3-[3-(3,4-Difluoro-phenoxy)methyl]-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide

15

APCI-MS m/z: 435 [MH+]

Example 146

N-(2-{3-[4-(4-Fluoro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide

20

APCI-MS m/z: 403 [MH+]

Example 147

3-(2-{3-[4-(4-Fluoro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propionic acid methyl ester

25

APCI-MS m/z: 432 [MH+]

30 **Example 148**

1-(2,6-Dimethoxy-phenoxy)-3-[4-(4-fluoro-phenoxy)-piperidin-1-yl]-propan-2-ol

APCI-MS m/z: 406 [MH+]

5 **Example 149**

1-[4-(4-Fluoro-phenoxy)-piperidin-1-yl]-3-(2-methoxy-phenoxy)-propan-2-ol

APCI-MS m/z: 376 [MH+]

10 **Example 150**

1-(2-{3-[4-(4-Fluoro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-ethanone

APCI-MS m/z: 388 [MH+]

15 **Example 151**

2-{3-[4-(4-Fluoro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-N,N-dimethyl-benzamide

APCI-MS m/z: 417 [MH+]

20

Example 152

1-(2-{3-[4-(4-Fluoro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propan-1-one

25 APCI-MS m/z: 402 [MH+]

Example 153

(2-{3-[4-(4-Fluoro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-benzoylamino)-acetic acid methyl ester

30

APCI-MS m/z: 461 [MH+]

Example 154

N-(2-{3-[3-(4-Fluoro-phenoxy)methyl]-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-
5 acetamide

APCI-MS m/z: 417 [MH+]

Example 155

10 3-(2-{3-[3-(4-Fluoro-phenoxy)methyl]-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-
propionic acid methyl ester

APCI-MS m/z: 446 [MH+]

15 **Example 156**

1-[3-(4-Fluoro-phenoxy)methyl]-piperidin-1-yl]-3-(2-methoxy-phenoxy)-propan-2-ol

APCI-MS m/z: 390 [MH+]

20 **Example 157**

1-(2-{3-[3-(4-Fluoro-phenoxy)methyl]-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-
ethanone

APCI-MS m/z: 402 [MH+]

25

Example 158

2-(2-{3-[3-(4-Fluoro-phenoxy)methyl]-piperidin-1-yl]-2-hydroxy-propoxy}-
benzoylamino)-2-methyl-propionic acid methyl ester

30 APCI-MS m/z: 503 [MH+]

Example 159

2-{3-[3-(4-Fluoro-phenoxy-methyl)-piperidin-1-yl]-2-hydroxy-propoxy}-N,N-dimethyl-benzamide

5

APCI-MS m/z: 431 [MH⁺]

Example 160

1-(2-{3-[3-(4-Fluoro-phenoxy-methyl)-piperidin-1-yl]-2-hydroxy-propoxy}-6-methoxy-phenyl)-ethanone

10

APCI-MS m/z: 432 [MH⁺]

Example 161

N-(2-{3-[4-(4-Acetyl-amino-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide

15

APCI-MS m/z: 442 [MH⁺]

Example 162

N-(4-{1-[3-(2-Acetyl-phenoxy)-2-hydroxy-propyl]-piperidin-4-yloxy}-phenyl)-acetamide

20

APCI-MS m/z: 427 [MH⁺]

25

Example 163

N-(4-cyano-2-{3-[4-(3,4-dichloroanilino)-1-piperidinyl]-2-hydroxypropoxy}phenyl)acetamide

30

APCI-MS: m/z 477 [MH⁺]

Example 164

3-(2-{3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propionic acid methyl ester

5

APCI-MS m/z: 448 [MH+]

Example 165

1-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-3-(2-methoxy-phenoxy)-propan-2-ol

10

APCI-MS m/z: 392 [MH+]

Example 166

1-(2-{3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-ethanone

15

APCI-MS m/z: 404 [MH+]

Example 167

2-(2-{3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-benzoylamino)-2-methyl-propionic acid methyl ester

20

APCI-MS m/z: 505 [MH+]

Example 168

2-{3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-N,N-dimethyl-benzamide

25

APCI-MS m/z: 433 [MH+]

30 **Example 169**

1-(2-{3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-6-methoxy-phenyl)-ethanone

APCI-MS m/z: 434 [MH+]

5

Example 170

1-(2-{3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propan-1-one

10 APCI-MS m/z: 418 [MH+]

Example 171

(2-{3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-benzoylamino)-acetic acid methyl ester

15

APCI-MS m/z: 477 [MH+]

Example 172

N-(2-{3-[3-(4-Chloro-phenoxy)methyl]-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide

20

APCI-MS m/z: 433 [MH+]

Example 173

3-(2-{3-[3-(4-Chloro-phenoxy)methyl]-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propionic acid methyl ester

25

APCI-MS m/z: 462 [MH+]

30 Example 174

1-(2-{3-[3-(4-Chloro-phenoxy)methyl]-piperidin-1-yl}-2-hydroxy-propoxy)-phenyl-ethanone

APCI-MS m/z: 418 [MH+]

5

Example 175

2-{3-[3-(4-Chloro-phenoxy)methyl]-piperidin-1-yl}-2-hydroxy-propoxy-N,N-dimethyl-benzamide

10 APCI-MS m/z: 447 [MH+]

Example 176

1-(2-{3-[3-(4-Chloro-phenoxy)methyl]-piperidin-1-yl}-2-hydroxy-propoxy)-phenyl-propan-1-one

15

APCI-MS m/z: 432 [MH+]

Example 177

20 N-[2-((1R,2R)-2-[4-(3,4-dichlorophenoxy)-1-piperidinyl]-1-hydroxycyclopentyl)methoxy)phenyl]acetamide

Example 178

Methyl (2S,4R)-1-{3-[2-(acetylamino)phenoxy]-2-hydroxypropyl}-4-[(4-chlorobenzyl)oxy]-2-pyrrolidinecarboxylate hydrochloride

25

Examples 179-189

Starting materials:

A) (3,4-Dichloro-phenyl)-piperidin-4-yl-amine

In a nitrogen filled reaction vessel 4-oxo-piperidine-1-carboxylic acid tert-butyl ester (2.46g, 12.3 mmol) and 3,4-dichloro-phenylamine (1.0 g, 6.17 mmol) were dissolved in dichloromethane (28 ml) and acetic acid (2.12 ml). Sodium triacetoxymethylborohydride (3.67 g, 17.3 mmol) was added at room temperature. The reaction was stirred over night and then poured into a sodium hydrogencarbonate solution (5%). The water phase was shaken three times with ethyl acetate (EtOAc). The combined organic phase was dried over sodium sulfate, evaporated and purified by flash chromatography (EtOAc : Heptane 3:7) giving 1.7 g, 81 % of pure compound. The BOC-protected title compound was dissolved in dichloromethane (26 ml) and trifluoro acetic acid (13 ml) and stirred at room temperature for 3h, evaporated and dissolved in diethyl ether and sodium hydroxide (1 M). The organic layer was separated and the water phase washed twice with ether. The combined organic layer was washed with a small portion of brine, dried over sodium sulfate and evaporated to give 1.15g (76%) of the title compound.

¹H-NMR (400MHz, DMSO-d₆): δ 7.20 (d, 1H, J = 8.9 Hz), 6.73 (d, 1H, J = 2.7 Hz), 6.54 (dd, 1H, J = 8.8, 2.7 Hz), 5.95 (d, 1H, J = 8.1 Hz), 3.22 (m, 1H), 2.91 (bd, 2H, J = 12.6 Hz), 2.51 (m, 2H), 2.02 (bs, 1H), 1.81 (bd, 2H, J = 12.4 Hz), 1.18 (m).

APCI-MS: m/z 245 [M⁺]

B) (4-Chloro-phenyl)-piperidin-4-yl-amine

Was synthesised in the same way as (A) from 4-oxo-piperidine-1-carboxylic acid tert-butyl ester (3.59 g, 18.0 mmol), 4-chloro-phenylamine (1.15 g, 9.0 mmol) and sodium triacetoxymethylborohydride (5.34 g, 25.2 mmol) in dichloromethane (40 ml) and acetic acid (3.1 ml). The deprotection was run in dichloromethane (37 ml) and trifluoro acetic acid (18 ml). Yield 1.5 g, 79 %

¹H-NMR (400MHz, DMSO-d₆): δ 7.04 (d, 2H, J = 8.9 Hz), 6.55 (d, 2H, J = 8.9 Hz), 5.62 (d, 1H, J = 8.1 Hz), 3.18 (m, 1H), 2.92 (bd, 2H, J = 12.6 Hz), 2.50 (m, 2H), 1.99 (bs, 1H), 1.82 (d, 2H, J = 12.7 Hz), 1.18 (m, 2H).

APCI-MS: m/z 211 [MH⁺]

C) (4-Fluoro-phenyl)-piperidin-4-yl-amine

Was synthesised in the same way as (A) from 4-oxo-piperidine-1-carboxylic acid tert-butyl ester (3.59 g, 18.0 mmol), 4-fluoro-phenylamine (1.0 g, 9.0 mmol) and sodium triacetoxymethylborohydride (5.34 g, 25.2 mmol) in dichloromethane (40 ml) and acetic acid (3.1 ml). The deprotection was run in dichloromethane (37 ml) and trifluoro acetic acid (18 ml). Yield 1.1 g, 63 %

¹H-NMR (400MHz, DMSO-d₆): δ 6.85 (t, 2H, J = 9.0 Hz), 6.51 (dd, 2H, J = 9.1, 4.6 Hz), 5.27 (d, 1H, J = 8.2 Hz), 3.13 (m, 1H), 2.89 (bd, 2H, J = 12.5 Hz), 2.48 (m, 2H), 1.80 (bd, 2H, J = 12.3 Hz), 1.14 (m, 2H).

APCI-MS: m/z 195 [MH⁺]

D) (3,4-Difluoro-phenyl)-piperidin-4-yl-amine

Was synthesised in the same way as (A) from 4-oxo-piperidine-1-carboxylic acid tert-butyl ester (3.59 g, 18.0 mmol), 3,4-difluoro-phenylamine (1.16 g, 9.0 mmol) and sodium triacetoxymethylborohydride (5.34 g, 25.2 mmol) in dichloromethane (40 ml) and acetic acid (3.1 ml). The deprotection was run in dichloromethane (37 ml) and trifluoro acetic acid (18 ml). Yield 1.26 g, 66 %

¹H-NMR (400MHz, DMSO-d₆): δ 7.05 (dt, 1H, J = 10.8, 9.2 Hz), 6.50 (ddd, 1H, J = 14.1, 7.0, 2.8 Hz), 6.32 (bd, 1H, J = 9.20 Hz), 5.64 (d, 1H, J = 8.14 Hz), 3.17 (m, 1H), 2.90 (bd, 2H, J = 12.6 Hz), 2.50 (m, 2H), 2.00 (bs, 1H), 1.81 (bd, 2H, J = 12.6 Hz), 1.16 (m, 2H)

APCI-MS: m/z 213 [MH⁺]

Example 179

N-(2-{3-[4-(3,4-Dichloroanilino)-1-piperidinyl]-2-hydroxypropoxy}-4-methylphenyl)acetamide

APCI-MS: m/z 466 [MH⁺]

Example 180**N-(2-{3-[4-(4-Chloroanilino)-1-piperidinyl]-2-hydroxypropoxy}phenyl)acetamide**5 APCI-MS: m/z 418[MH⁺]**Example 181****N-(4-Chloro-2-{3-[4-(4-chloroanilino)-1-piperidinyl]-2-hydroxypropoxy}phenyl)-
acetamide**

10

APCI-MS: m/z 452[MH⁺]**Example 182****N-(2-{3-[4-(4-Chloroanilino)-1-piperidinyl]-2-hydroxypropoxy}-4-
15 cyanophenyl)acetamide**APCI-MS: m/z 443[MH⁺]**Example 183**20 **N-(2-{3-[4-(4-Chloroanilino)-1-piperidinyl]-2-hydroxypropoxy}-4-
methylphenyl)acetamide**APCI-MS: m/z 432[MH⁺]25 **Example 184****N-(5-Chloro-2-{3-[4-(4-fluoroanilino)-1-piperidinyl]-2-
hydroxypropoxy}phenyl)acetamide**APCI-MS: m/z 436[MH⁺]

30

Example 185

N-(5-Chloro-2-{3-[4-(3,4-difluoroanilino)-1-piperidinyl]-2-hydroxypropoxy}phenyl)acetamide

5 APCI-MS: m/z 454[MH+]

Example 186

N-(5-Cyano-2-{3-[4-(4-fluoroanilino)-1-piperidinyl]-2-hydroxypropoxy}-phenyl)acetamide

10

APCI-MS: m/z 427[MH+]

Example 187

N-(5-Cyano-2-{3-[4-(3,4-difluoroanilino)-1-piperidinyl]-2-hydroxypropoxy}phenyl)acetamide

15

APCI-MS: m/z 445[MH+]

Example 188

N-(2-{3-[4-(4-Fluoroanilino)-1-piperidinyl]-2-hydroxypropoxy}-4-methylphenyl)acetamide

20

APCI-MS: m/z 416[MH+]

Example 189

N-(2-{3-[4-(3,4-Difluoroanilino)-1-piperidinyl]-2-hydroxypropoxy}-4-methylphenyl)acetamide

25

APCI-MS: m/z 434[MH+]

30

Example 190**N-(2-{3-[3(S)-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-(R)-hydroxy-propoxy-phenyl}acetamide****i) 3-(S)-(4-Chloro-phenoxy)-pyrrolidine CF₃COOH**

To a solution of triphenyl phosphine (4.2 g, 16.02 mmol) in THF (75 mL) was added diethyl azodicarboxylate (2.52 mL) at 0 °C, after 15 min 4-chlorophenol (2.05 g, 16.02 mmol) was added and after another 10 min 3-hydroxy-pyrrolidine-1-carboxylic acid tert-butyl ester (3.0 g, 16.02 mmol) in THF (20 mL) was added slowly. After addition was complete the ice bath was removed and the reaction mixture was kept at room temperature overnight. The solvent was removed in vacuo and the residue was stirred with diethyl ether, the solid triphenyl phosphine was filtered off. The residue was purified by flash chromatography (0-0.5% MeOH in CHCl₃) to give the subtitled compound (3.65 g, 76%) which was dissolved in dichloromethane (60 mL) and trifluoroacetic acid (15 mL) was added. The reaction mixture was kept at room temperature for 30 min. The solvent was removed in vacuo. The residue was dissolved in dichloromethane, diethylether and hexane were added. The solid was filtered off to give the subtitled compound 3.70 g, 97%.

¹H-NMR (CDCl₃, 400 MHz): δ 10.20 (s, 1H), 9.99 (s, 1H), 7.25 (d, J 8.8 Hz, 2H), 6.78 (d, J 8.8 Hz, 2H), 4.99 (m, 1H), 3.41 (m, 4H), 2.30 (m 1H), 2.20 (m, 1H).

APCI-MS: m/z 198 (MH⁺).

ii) N-(2-{3-[3(S)-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-(R)-hydroxy-propoxy-phenyl}acetamide

A mixture of 3-(S)-(4-chloro-phenoxy)-pyrrolidine.CF₃COOH (312 mg, 1.0 mmol), N-acetyl-2-(2,3-epoxypropoxy)aniline (207 mg, 1.0 mmol), K₂CO₃ (560 mg) in EtOH (10 mL) was stirred at 65 °C for 4 h. The solvent was removed in vacuo. The residue was partitioned between ethyl acetate and water. The organic layer was washed with aqueous NHCl solution, then with water. The organic layer was dried over Na₂SO₄, filtered, concentrated. The residue was purified by flash chromatography (0-3% MeOH in CHCl₃)

to give a mixture of diastereomers (310 mg, 77%). The diastereomers were separated by HPLC to give N-(2-{3-[3(S)-(4-chloro-phenoxy)-pyrrolidin-1-yl]-2-(R)-hydroxy-propoxy-phenol}acetamide (57 mg)

5 ¹H-NMR (CDCl₃, 400 MHz): δ 8.36 (m, 1H), 8.25 (s, 1H), 7.25 (m, 2H), 6.99 (m, 2H), 6.93 (m, 1H), 6.75 (m, 2H), 4.80 (m, 1H), 4.08 (m, 2H), 3.96 (m, 1H), 3.11 (dd, J 5.9 10.5 Hz, 1H), 3.01 (m, 1H), 2.82 (m, 2H), 2.59 (m, 1H), 2.51 (dd, J 3.2, 12.0 Hz, 1H), 2.29 (m, 1H), 2.19 (s, 3H), 2.01 (m, 1H). APCI-MS: m/z 405 (MH⁺).

10 **Example 191**

N-(2-{3-[3S-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2S-hydroxy-propoxy}-phenyl)-acetamide hydrochloride

The reaction was performed analogously to Example 190.

15

¹H-NMR (CDCl₃, 400 MHz): δ 8.35 (m, 1H), 8.26 (s, 1H), 7.24 (m, 2H), 6.99 (m, 2H), 6.92 (m, 1H), 6.75 (m, 2H), 4.80 (m, 1H), 4.12 (m, 2H), 3.95 (m, 1H), 2.95 (m, 2H), 2.80 (m, 3H), 2.52 (dd, 3.4, 12.2 Hz, 1H), 2.30 (m, 1H), 2.19 (s, 3H), 2.01 (m, 1H). APCI-MS: m/z 405 (MH⁺).

20

Example 192

N-(2-{3-[3(R)-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-(S)-hydroxy-propoxy-phenyl}acetamide

25 The reaction was performed analogously to Example 190.

¹H-NMR (CDCl₃, 400 MHz): δ 8.36 (m, 1H), 8.25 (s, 1H), 7.25 (m, 2H), 6.99 (m, 2H), 6.93 (m, 1H), 6.75 (m, 2H), 4.80 (m, 1H), 4.08 (m, 2H), 3.96 (m, 1H), 3.11 (dd, J 5.9 10.5 Hz, 1H), 3.01 (m, 1H), 2.82 (m, 2H), 2.59 (m, 1H), 2.51 (dd, J 3.2, 12.0 Hz, 1H), 2.29 (m, 30 1H), 2.19 (s, 3H), 2.01 (m, 1H).

APCI-MS: m/z 405 (MH⁺).

Example 193

N-[5-Chloro-2-(((2S)-3-[(3S)-3-(4-chloro-phenoxy)pyrrolidinyl]-2-hydroxypropyl}oxy)phenyl]acetamide

The reaction was performed analogously to Example 190.

¹H-NMR (CDCl₃, 400 MHz): δ 8.45 (m, 1H), 8.36 (br. S, 1H), 7.23 (m, 2H), 6.95 (m, 1H), 6.85 (m, 1H), 6.75 (m, 2H), 4.80 (m, 1H), 4.07 (m, 2H), 3.91 (m, 1H), 2.95 (m, 2H), 2.80 (m, 3H), 2.49 (dd, J 3.2, 12.0 Hz, 1H), 2.30 (m, 1H), 2.19 (s, 3H), 2.03 (m, 1H).

APCI-MS: m/z 439 (MH⁺).

Example 194

N-[5-Chloro-2-(((2R)-3-[(3R)-3-(4-chloro-phenoxy)pyrrolidinyl]-2-hydroxypropyl}oxy)phenyl]acetamide

The reaction was performed analogous to Example 190.

¹H-NMR (CDCl₃, 400 MHz): δ 8.45 (m, 1H), 8.36 (br. S, 1H), 7.23 (m, 2H), 6.95 (m, 1H), 6.85 (m, 1H), 6.75 (m, 2H), 4.80 (m, 1H), 4.07 (m, 2H), 3.91 (m, 1H), 2.95 (m, 2H), 2.80 (m, 3H), 2.49 (dd, J 3.2, 12.0 Hz, 1H), 2.30 (m, 1H), 2.19 (s, 3H), 2.03 (m, 1H).

APCI-MS: m/z 439 (MH⁺).

Example 195

N-[5-Chloro-2-(((2S)-3-[(3R)-3-(4-chloro-phenoxy)pyrrolidinyl]-2-hydroxypropyl}oxy)phenyl]acetamide

The reaction was performed analogously to Example 190.

¹H-NMR (CDCl₃, 400 MHz): δ 8.45 (m, 1H), 8.34 (br. S, 1H), 7.22 (m, 2H), 6.94 (m, 1H), 6.85 (m, 1H), 6.75 (m, 2H), 4.80 (m, 1H), 4.08 (m, 2H), 3.90 (m, 1H), 3.11 (dd, J 5.9, 10.5 Hz, 1H), 3.02 (m, 1H), 2.81 (m, 2H), 2.58 (m 1H), 2.49 (dd, J 3.5, 12.1 Hz, 1H), 2.30 (m, 1H), 2.18 (s, 3H), 2.01 (m, 1H).

5 APCI-MS: m/z 439 (MH⁺).

Example 196

N-[5-Chloro-2-((2R)-3-[(3S)-3-(4-chloro-phenoxy)pyrrolidinyl]-2-hydroxypropyl)oxy)phenyl]acetamide

10

The reaction was performed analogous to Example 190.

¹H-NMR (CDCl₃, 400 MHz): δ 8.45 (m, 1H), 8.34 (br. S, 1H), 7.22 (m, 2H), 6.94 (m, 1H), 6.85 (m, 1H), 6.75 (m, 2H), 4.80 (m, 1H), 4.08 (m, 2H), 3.90 (m, 1H), 3.11 (dd, J 5.9, 10.5
15 Hz, 1H), 3.02 (m, 1H), 2.81 (m, 2H), 2.58 (m 1H), 2.49 (dd, J 3.5, 12.1 Hz, 1H), 2.30 (m, 1H), 2.18 (s, 3H), 2.01 (m, 1H).

APCI-MS: m/z 439 (MH⁺).

Example 197

20 **N-(2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-4,5-difluoro-phenyl)-acetamide**

i) 4,5-Difluoro-2-nitro-phenol

In a flask was dissolved 3,4-Difluorophenol (3.10 g, 23.7 mmole) in acetic acid (15 ml). To
25 the stirred solution was added dropwise a solution of fuming HNO₃ (1.25 g, 29.7 mmole)
in acetic acid (6 ml). The temperature was kept under 50°C during the entire addition. After
completed addition, the mixture was stirred for another hour. The reaction mixture was
then poured onto ice-water, giving precipitation of a yellowish solid. The solid was
collected by filtration, and dried. The solid was purified on silica (Heptane : EtOAc 5:1),

giving the sub-title compound (2.05 g, 50%) as a yellow oil, which crystallizes on standing.

¹H-NMR (400 MHz, CDCl₃) δ: 10.61 (1H, s); 8.00 (1H, dd, *J* 9.6 8.2 Hz); 7.00 (1H, dd, *J* 10.4 6.8 Hz)

ii) N-(4,5-Difluoro-2-hydroxy-phenyl)-acetamide

In a flask was added the product obtained in i) (0.59 g, 3.37 mmole), and acetic acid (10 ml). The solution was heated with stirring to 90°C, and Tin (powder, 1.60 g, 13.5 mmole) was added. The flask was sealed and heated with stirring for another hour, and the hot solution was filtered through celite. The filter was then washed with another 10 ml of hot acetic acid. To the filtrate was added water (25 ml) and acetic anhydride (0.5 ml, 5.29 mmole), and the resulting mixture was heated with stirring at 60°C for 20 minutes. The mixture was allowed to cool, and was partitioned between EtOAc and water. The organic phase was collected and washed with water and brine. The organic phase was evaporated to give the 0.63 g (100%) of the sub-title compound as a solid.

¹H-NMR (400 MHz, DMSO-*d*₆) δ: 10.25 (1H, s); 9.31 (1H, bs); 7.88 (1H, dd, *J* 12.8 7.9 Hz); 6.83 (1H, dd, *J* 12.1 7.7 Hz); 2.08 (3H, s)

iii) N-(4,5-Difluoro-2-oxiranylmethoxy-phenyl)-acetamide

In a vial was added the compound obtained in ii) (0.4 g, 2.137 mmole), epibromohydrine (0.35 g, 2.55 mmole), K₂CO₃ (0.6 g, 4.4 mmole) and DMF (2 ml). The vial was sealed and heated with stirring (2 hours, 60°C). The mixture was then partitioned between EtOAc and water, and the organic phase was washed twice with water and once with brine, and was finally evaporated to give a brown solid. The crude epoxide was purified on silica, to give 0.27 g (52%) of the sub-title compound as a slightly pink solid.

¹H-NMR (400 MHz, CDCl₃) δ: 8.37 (1H, dd, *J* 12.2 8.8 Hz); 7.85 (1H, bs); 6.78 (1H, dd, *J* 11.2 7.1 Hz); 4.34 (1H, dd, 11.5 2.2 Hz); 3.90 (1H, dd, 11.6 6.3 Hz); 3.40-3.36 (1H, m); 2.98 (1H t, *J* 4.5 Hz); 2.81 (1H, dd, *J* 4.7 6.3 Hz); 2.22 (3H, s)

5 iv) *N*-(2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-4,5-difluoro-phenyl)-acetamide

In a vial was added the compound obtained in iii) (0.059 g, 0.24 mmole), 3-(4-chlorophenoxy)-pyrrolidine (0.048 g, 0.24 mmole) and ethanol (2 ml, 99.5%). The vial was sealed and the content was heated with stirring to 75°C for 2 hours. The crude solution was
10 evaporated and the obtained oil purified on silica to the title compound which was lyophilized as the hydrochloride. The title compound was obtained as a white solid (0.075 g, 65%). The compound was a mixture of four stereoisomers, which had an effect on the NMR-spectra.

15 ¹H-NMR (400 MHz, DMSO-*d*₆) δ: 10.78-10.30 (1H, m); 9.30 (1H, s); 8.07 (1H, dd, *J* 12.8 9.3 Hz); 7.40-7.34 (2H, m); 7.23 (1H, dd, *J* 12.7 7.5 Hz); 7.05-6.99 (2H, m); 6.19 (1H, bs); 5.23-5.11 (1H, m); 4.35 (1H, bs); 4.08-3.97 (1.5H, m), 3.96-3.90 (1H, m); 3.84-3.70 (1.5H, m); 3.63-3.23 (4H, m); 2.66-2.00 (5H, m)

20 APCI-MS: *m/z* 411.1 [MH⁺]

Example 198

N-(5-Chloro-2-[2-hydroxy-3-(3-phenoxy-pyrrolidin-1-yl)-propoxy]-phenyl)-acetamide

25 The compound was prepared analogously to Example 197.

¹H-NMR (400 MHz, DMSO-*d*₆) δ: 10.80-10.36 (1H, m); 9.26 (1H, s); 8.14 (1H, s); 7.32 (2H, t, *J* 8.35 Hz); 7.11-6.95 (5H, m); 6.31-6.02 (1H, m); 5.24-5.12 (1H, m); 4.37 (1H, bs); 4.10-3.97 (1.5H, m); 3.95-3.88 (1H, m) 3.84-3.68 (1.5H, m); 3.64-3.26 (4H, m); 2.65-2.52
30 (0.5H, m); 2.35-2.02 (4.5H, m)

APCI-MS: m/z 405.2 [MH⁺]

Example 199

***N*-(5-Chloro-2-{2-hydroxy-3-[3-(4-nitro-phenoxy)-pyrrolidin-1-yl]-propoxy}-phenyl)-acetamide**

The compound was prepared analogously to Example 197.

¹H-NMR (400 MHz, DMSO-*d*₆) δ: 10.95-10.48 (1H, m); 9.26 (1H, s); 8.24 (2H, d, *J* 9.6 Hz); 8.13 (1H, bs); 7.23-7.17 (2H, m); 7.12-7.02 (2H, m); 6.20 (1H, bs); 5.43-5.30 (1H, m); 4.38 (1H, m); 4.18-4.06 (0.5H, m); 4.05-3.97 (1H, m); 3.95-3.87 (1H, m); 3.86-3.72 (1.5H, m); 3.69-3.27 (4H, m); 2.73-2.60 (0.5H, m); 2.46-2.08 (4.5H, m)

APCI-MS: m/z 450.1 [MH⁺]

Example 200

***N*-(5-Acetyl-2-{3-[3-(3,4-dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide**

The compound was prepared analogously to Example 197.

APCI-MS: m/z 481.2, 483.2 [MH⁺]

Example 201

4-Acetylamino-3-{3-[3-(3,4-dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-benzoic acid methyl ester

The compound was prepared analogously to Example 197.

APCI-MS: m/z 497.1, 499.2 [MH+]

Example 202

N-(3-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-naphthalen-
5 2-yl)-acetamide

The compound was prepared analogously to Example 197.

APCI-MS: m/z 489.2, 491.2 [MH+]

10

Example 203

N-(2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-5-cyano-phenyl)-
acetamide

15 The title compound was prepared according to the method in Example 197.

APCI-MS: m/z 430.2 [MH+]

Example 204

4-Acetylamino-3-{3-[3-(4-chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-
20 benzoic acid methyl ester

The compound was prepared analogously to Example 197.

APCI-MS: m/z 463.2 [MH+]

25

Example 205

N-(3-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-naphthalen-2-
yl)-acetamide

30 The title compound was prepared according to the method in Example 197.

APCI-MS: m/z 455.2 [MH⁺]

Example 206

5 ***N*-(5-Cyano-2-{3-[4-(3,4-dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide**

The title compound was prepared according to the method in Example 197.

10 APCI-MS: m/z 478.2 480.1 [MH⁺]

Example 207

***N*-(2-{3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-5-trifluoromethyl-phenyl)-acetamide**

15 The title compound was prepared according to the method in Example 197.

APCI-MS: m/z 521.1 523.2 [MH⁺]

20 **Example 208**

***N*-(5-Chloro-2-{3-[3-(4-fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide trifluoroacetate**

The title compound was prepared according to the method in Example 197.

25 APCI-MS: m/z 423.1, 424.9 [MH⁺]

Example 209

30 ***N*-(5-Acetyl-2-{3-[3-(4-chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide trifluoroacetate**

i) N-(5-Acetyl-2-oxiranylmethoxy-phenyl)-acetamide

To a solution of 4-acetyl-2-nitrophenol (0.50g, 2.76mmol) in THF (20 ml) was added 10% Pd/C (0.15g). The resultant mixture was hydrogenated with H₂ at 1 atm for 5 hours and
5 was then filtered through celite and evaporated to give 0.63g of a red oil. Water (20 ml) and acetic anhydride (0.35g, 3.44mmol) was added and the mixture was stirred vigorously for 5 minutes. The reaction mixture was then heated with stirring to 60°C for 30 minutes, and was then allowed to cool. A red solid was formed and the precipitate was collected by filtration, washed with water and dried to give 0.27g (1.40mmol) of N-(5-Acetyl-2-
10 hydroxy-phenyl)-acetamide. This was dissolved in DMF (5ml). K₂CO₃ (0.34g, 2.45mmol) and epibromohydrin (0.21g, 1.54mmol) was added and the resulting mixture was heated with stirring at 50°C for 3 hours. The mixture was partitioned between EtOAc and water 40+40ml. The organic phase was washed twice with water and once with brine and finally concentrated in vacuo to give a red oil. The crude product was purified on silica
15 (Heptane/EtOAc, 1:2-1:4) to give 110 mg (16%) of the subtitle compound.

¹H-NMR (400MHz, CDCl₃): δ 9.03 (1H, d, J1.9Hz), 7.81 (1H, bs), 7.74 (1H, dd, J8.6, 2.3Hz), 6.96 (1H, d, J8.6Hz), 4.48 (1H, dd, J11.3, 2.4Hz) 4.00 (1H, dd, J11.4, 6.4Hz), 3.45-3.40 (1H, m), 2.99 (1H, t, J4.4Hz), 2.79 (1H, dd, J4.7, 2.6 Hz), 2.59 (3H, s), 2.26 (3H, s).

ii) N-(5-Acetyl-2-{3-[3-(4-chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide trifluoroacetate

The title compound was prepared according to the method described in Example 197.

25 APCI-MS: m/z 447, 449 [MH⁺]

Example 210

N-(2-{3-[3-(4-Chlorophenoxy)-1-pyrrolidinyl]-2-hydroxypropoxy}phenyl)-methanesulfonamide

i) 1-(2-Aminophenoxy)-3-[3-(4-chlorophenoxy)-1-pyrrolidinyl]-2-propanol dihydrochloride

A mixture of *N*-(2-{3-[3-(4-chlorophenoxy)-1-pyrrolidinyl]-2-hydroxypropoxy}phenyl)acetamide (0.95 g, 2.34 mmol) and concentrated hydrochloric acid (25 mL) was heated (100–105 °C) for 3 hours then allowed to stand at room temperature overnight. The mixture was concentrated at reduced pressure to a third of its volume basified with saturated sodium hydrogen carbonate. The resulting suspension was extracted twice with ethyl acetate. The organic extracts were dried, the solvent was evaporated at reduced pressure to give a pale brown oil. This oil was dissolved in a minimum amount of methanol, diluted with ethyl ether and the product precipitated by addition of HCl-saturated ethyl ether. The product was filtered to afford the subtitle product (0.93 g, 91.2%).

APCI-MS: *m/z* 363 [*MH*⁺] for the free base.

ii) *N*-(2-{3-[3-(4-Chlorophenoxy)-1-pyrrolidinyl]-2-hydroxypropoxy}phenyl)-methanesulfonamide

Methanesulfonyl chloride (35 mg, 0.3 mmol) was added to cold (0 °C), stirred mixture of the above amine (110 mg, 0.25 mmol) and pyridine (0.4 mL) in dry dichloromethane (10 mL). The mixture was then stirred at room temperature for 1.5 hour then concentrated. The residue was partitioned between ethyl acetate and water. The organic phase was concentrated and the residue was purified by flash chromatography (silica gel, dichloromethane-methanol, 25:1) to afford the title compound (68 mg, 61.8%) as a foam.

¹H-NMR (400MHz, CDCl₃): δ 7.51 (dd, 1H, *J* = 1.4 and 8.0 Hz), 7.22 (m, 2H), 7.10 (m, 1H), 7.68 (m, 1H), 6.92 (d, 1H, *J* = 9.0 Hz), 6.76 (m, 2H), 5.78 (very bs, 1H), 4.80 (m, 1H), 4.20 (m, 1H), 4.08 (m, 1H), 3.98 (m, 1H), 3.16 (m, 1H), 3.01 (m, 1H), 2.96 (s, 3H), 2.89 (m, 2H), 2.74 (m, M), 2.68 (dd, 1H, *J* = 4.0 and 12.2 Hz), 2.3 (m, 1H), and 2.02 (m, 1H).

¹³C-NMR, 400 MHz, CDCl₃): δ 155.9, 149.4, 129.4, 126.9, 125.8, 125.77, 125.75, 122.29, 122.26, 122.17, 115.5, 113.52, 113.50, 76.52, 76.49, 72.15, 72.09, 67.18, 67.08, 60.24, 60.07, 57.96, 57.94, 53.18, 52.98, 39.1, 31.92, 31.90.

5 APCI-MS: m/z 441[MH⁺].

Example 211

N-(5-Chloro-2-[3-[3,4-dichlorophenoxy]-1-pyrrodinyl]-2-hydroxypropoxy)-phenyl)urea

10

i) N-(5-Chloro-2-hydroxyphenyl)urea

A solution of potassium cyanate (6.14 g, 75.6 mmol) in water (50 mL) was added dropwise to a stirred suspension of 2-amino-4-chlorophenol (5.00 g, 34.8 mmol) in a mixture of acetic acid (350 mL) and water (250 mL) and the resulting solution was stirred at room temperature for 3 hours. The reaction mixture was extracted three times with ethyl ether. The ether extracts were combined and concentrated to a thick oil. A 10% solution of sodium hydrogen carbonate (250 mL) was added to the above oil. The solid product was filtered and washed several times with water and recrystallized (toluene containing a little methanol) to afford the subtitle compound (3.27 g, 50.4%)

20

¹H-NMR (400MHz, DMSO-*d*₆): δ 10.1 (s, 1H), 8.07 (d, 1H, J=2.2 Hz), 8.04 (s, 1H), 6.75-6.78 (m, 2H), 6.29 (bs, 2H).

¹³C-NMR: δ 156.0, 144.1, 130.0, 122.5, 120.2, 117.5, 115.2.

25

ii) N-[5-Chloro-2-(2-oxiranylmethoxy)phenyl]urea

A suspension of N-(5-chloro-2-hydroxyphenyl)urea (53 mg, 0.28 mmol), cesium carbonate (92 mg, 0.28 mmol) and epibromohydrine (49 mg, 0.36 mmol) in dry DMF (0.6 mL) was stirred at room temperature for 24 hours. The mixture was then partitioned between ethyl acetate and water. The organic phase was washed with water three times, dried and

30

concentrated to a solid residue. This crude product was recrystallized (ethyl ether and heptane to afford the subtitle compound (18 mg, 26.5%).

¹H-NMR (400MHz, DMSO-*d*₆): δ 8.20 (d, 1H, J=2.2 Hz), 8.00 (s, 1H), 7.00 (d, 1H, J=8.8 Hz), 6.88 (dd, 1H, J= 2.4 and 8.6 Hz), 6.40 (bs, 2H), 4.40 (dd, 1H, J= 2.2 and 12.0 Hz), 3.90 (dd, 1H, J= 6.6 and 12.0 Hz), 3.37 (m, 1H), 2.88 (t, 1H, H= 4.8 Hz), 2.74 (m, 1H).

iii) N-(5-Chloro-2-[3-[3,4-dichlorophenoxy]-1-pyrrodinyl]-2-hydroxypropoxy)-phenyl)urea

A solution of the subtitle compound (ii) (16 mg, 0.07 mmol) and 3-(3,4-dichlorophenoxy)pyrrolidine (17 mg, 0.07 mmol) in absolute ethanol (1 mL) for 2 hours. The solvent was removed under reduced pressure and the residue was purified by flash chromatography (dichloromethane-methanol, 15:1) to afford the title compound (11 mg, 33%).

¹H-NMR (400MHz, DMSO-*d*₆): δ 8.18 (d, 1H, J= 2.6 Hz), 7.94 (s, 1H), 7.5 (d, 1H, J= 0.0Hz), 7.16 (d, 1H, J= 2.1 Hz), 6.82-6.98 (m, 3H), 6.33 (bs, 2H), 4.98 (m, 1H), 4.90 (m, 1H), 3.85-4.07 (m, 3H), 2.63-2.93 (m, 5H), 2.21-2.30 (m, 1H), 1.74 (m, 1H).

APCI-MS: m/z 198 [MH⁺]

Example 212

1-(3-{2-[(Aminocarbonyl)amino]phenoxy}-2-hydroxypropyl)-3-(4-chlorophenoxy)pyrrolidinium 2,2,2-trifluoroacetate

i) N-(2-Hydroxyphenyl)urea

A solution of potassium cyanate (3.94 g, 48.6 mmol) in water (30 mL) was added during 15 min. to suspension of 2-aminophenol (2.41 g, 22.1 mmol) in 50% aqueous acetic acid (160 mL). The resulting solution was allowed to stand at room temperature overnight and then extracted with ethyl ether (3 times). The combined organic extracts was concentrated

to small volume and poured into cold saturated aqueous sodium hydrogen carbonate. The solid was filtered and washed with water to afford the subtitle compound (1.61 g, 47.9%).

5 ¹H-NMR (400MHz, DMSO-d₆): δ 9.88 (s, 1H), 7.97 (s, 1H), 7.80 (bd, 1H), 6.77 (m, 1 H), 6.68 (m, 1H), 6.17 (s, 2H),

ii) N-[2-(2-Oxiranylmethoxy)phenyl]urea

A solution of epibromohydrin (0.94 g, 6.84 mmol) in dry DMF (2mL) was added dropwise
10 to a stirred suspension of *N*-(2-hydroxyphenyl)urea (0.65 g, 4.27 mmol) and cesium carbonate (2.22 g, 6.84 mmol) in DMF (8 mL). After 2 hours the mixture was partitioned between water and ethyl acetate. The aqueous phase was extracted with ethyl acetate and the combined organic phase was washed with water (3 times), dried and concentrated. The
15 semi-solid residue was dissolved in dichloromethane/ethyl ether, filtered and heptane was added to turbidity. After standing at room temperature overnight, the solid was filtered to afford the subtitle compound (0.28 g, 32%).

¹H-NMR (400MHz, DMSO-d₆): δ 8.07(m, 1H), 7.82 (s, 1H), 6.97 (m, 1H), 6.85 (m, 2H), 6.24 (bs, 2H), 4.37 (dd, 1H, J=2,5 and 11.6 Hz), 3.89 (dd, 1H, J=6.4 and 11.6 Hz), 3.38 (m,
20 1H), 2.87 (t, 1H, J=4.6 Hz), 2.75 (dd, 1H, J=2.6 and 5.2 Hz).

APCI-MS: m/z 209 [MH⁺].

iii) 1-(3-{2-[(Aminocarbonyl)amino]phenoxy}-2-hydroxypropyl)-3-(4-chlorophenoxy)pyrrolidinium 2,2,2-trifluoroacetate
25

A solution of *N*-[2-(2-oxiranylmethoxy)phenyl]urea (78 mg, 0.37 mmol) and 3-(4-chlorophenoxy)pyrrolidine (70 mg, 0.36 mmol) in absolute ethanol (4 mL) was heated at reflux for 2.5 hours. The mixture was then concentrated and the residue was purified by HPLC to afford the title compound (102 mg, 54.5%).

¹H-NMR (400MHz, DMSO-d₆ + D₂O): δ 7.98 (bd, 1H, J=7.2 Hz), 7.36 (d, 2H, J=8.8 Hz), 6.95-7.02 (m, 3H), 6.88 (m, 2H), 5.15 (bd, 1H), 4.26 (m, 1H). The remaining 10 aliphatic protons appear as complicated overlapping multiplets between δ 2.04 and 4.04.

APCI-MS: m/z 406 [MH⁺] and 408 [MH⁺+2] for the free base.

Example 213

1-(3-{2-[(Aminocarbonyl)amino]phenoxy}-2-hydroxypropyl)-3-(3,4-dichlorophenoxy)pyrrolidinium 2,2,2-trifluoroacetate

A solution of *N*-[2-(2-oxiranylmethoxy)phenyl]urea (Described under example 22, step ii; 69 mg, 0.33 mmol) and 3-(3,4-dichlorophenoxy)pyrrolidine (77 mg, 0.33 mmol) in absolute ethanol (4.5 mL) was heated at 70°C for 2 hours. The residue after evaporation of the solvent was purified by HPLC to afford the title compound (133 mg, 72.3%).

¹H-NMR (400MHz, DMSO-d₆ + D₂O): δ 7.96 (bd, 1H, J=7.4 Hz), 7.54 (bd, 1H, J=9.0 Hz), 7.27 (bs, 1H), 6.84-7.00 (m, 4H), 5.20 (bd, 1H), 4.26 (m, 1H). The remaining 10 aliphatic protons appear as complicated overlapping multiplets between δ 2.05 and 4.03.

APCI-MS: m/z 439.9 [M] and 442 [M+2] for the free base.

Example 214

1-(3-{2-[(Aminocarbonyl)amino]-4-chlorophenoxy}-2-hydroxypropyl)-3-(4-chlorophenoxy)pyrrolidinium 2,2,2-trifluoroacetate

A solution of *N*-[2-(2-oxiranylmethoxy)phenyl]urea (described under Example 212, step ii; 47 mg, 0.22 mmol) and 3-(4-chlorophenoxy)pyrrolidine (41 mg, 0.2 mmol) in absolute ethanol (3 mL) was heated at 70°C for 1.5 hours. The solvent was then evaporated and the residue was purified by HPLC to afford the title compound (67 mg, 60.9%).

¹H-NMR (400MHz, CD₃OD): δ 8.04(s, 1H), 7.31 (d, 2H, J=8.6 Hz), 6.94-6.98 (m, 4H), 5.20 (bs, 1H), 4.40 (m, 1H). The remaining 10 aliphatic protons appear as complicated overlapping multiplets between δ 2.25 and 4.13.

5 APCI-MS: m/z 440.1 [M] and 442.1 [M+2] for the free base.

Example 215

N-(2-{3-[3-(4-Chlorophenoxy)-1-pyrrolidinyl]-2-hydroxypropoxy}phenyl)-*N'*-ethylurea hydrochloride

10

An ether solution of *N*-(2-{3-[3-(4-chlorophenoxy)-1-pyrrolidinyl]-2-hydroxypropoxy}phenyl)urea [obtained from the hydrochloride (110mg, 0.25mmol) by treatment with 1M NaOH and extraction with ether] was treated with ethyl isocyanate (16μl, 0.2mmol) in a sealed vial for 15h at ambient temperature. After evaporation and
15 purification by flash chromatography on silica (EtOAc/MeOH 100/5) the appropriate fractions were acidified with 1M HCl and lyophilized from acetic acid to give the title compound as a white amorphous solid (35mg, 37%). The substance is a mixture of two diastereomeric pairs.

20 APCI-MS: m/z 434 [MH⁺]

Example 216

N-(2-{3-[3-(4-Chlorophenoxy)-1-pyrrolidinyl]-2-hydroxypropoxy}phenyl)-*N'*-methylurea hydrochloride

25

To a solution of *N*-(2-{3-[3-(4-chlorophenoxy)-1-pyrrolidinyl]-2-hydroxypropoxy}phenyl)urea [obtained from the hydrochloride (44mg, 0.1mmol) by treatment with 1M NaOH and extraction with ether] in DCM (3ml) di(*tert*-butyl)tricarboxylate (26mg, 0.1mmol) was added, and the solution was set aside for 15 min.

Methylamine (2M in DCM, 100 μ l, 0.2mmol) was added and the solution was left to stand over night. After evaporation the crude product was purified by preparative RP-HPLC using acetonitrile and water containing 0.1% TFA as mobil phase. The appropriate fraction was concentrated *in vacuo* and extracted with 1M NaOH and ether.

The residue from the organic phase was acidified with 1M HCl and lyophilized from acetic acid to afford the title compound as a white amorphous solid (15mg, 30%). The substance is a mixture of two diastereomeric pairs.

APCI-MS: m/z 420 [MH⁺]

Example 217

(2S,4S)-1-{3-[2-(Acetylamino)phenoxy]-2-hydroxypropyl}-4-(4-chlorophenoxy)-2-pyrrolidinecarboxylic acid; compound with trifluoroacetic acid

(i) Methyl (2S,4S)-1-{3-[2-(acetylamino)phenoxy]-2-hydroxypropyl}-4-(4-chlorophenoxy)-2-pyrrolidinecarboxylate hydrochloride

Methyl (2S,4S)-4-(4-chlorophenoxy)-2-pyrrolidinecarboxylate (370mg, 1.0mmol) and *N*-[2-(2-oxiranylmethoxy)phenyl]acetamide (207mg, 1.0mmol) dissolved in *tert*-butanol (7ml) was stirred in a sealed vial at 100°C over night. Workup of the crude material by flash chromatography on silica (DCM/MeOH 100/2), acidification of the appropriate fractions with 1M HCl and lyophilization from acetic acid afforded the subtitled compound as a white amorphous solid (360mg, 72%). The substance is a diastereomeric mixture.

APCI-MS: m/z 463 [MH⁺]

ii) (2S,4S)-1-{3-[2-(Acetylamino)phenoxy]-2-hydroxypropyl}-4-(4-chlorophenoxy)-2-pyrrolidinecarboxylic acid; compound with trifluoroacetic acid

Compound (i) (50mg, 0.1mmol) was dissolved in acetonitrile (2ml) and water (3ml). Lithium hydroxide hydrate (8mg, 0.2mmol) dissolved in water (0.5ml) was added. The reaction was complete after 0.5h as determined by analytical HPLC. The mixture was

acidified with TFA and purified by preparative RP-HPLC using acetonitrile and water containing 0.1% TFA as mobile phase. The appropriate fraction was concentrated *in vacuo* and the residue lyophilized from acetic acid to give the title compound as a white amorphous solid (27mg, 48%). The substance is a diastereomeric mixture.

5

APCI-MS: m/z 449 [MH⁺], 431 [MH⁺, lactone, minor amount]

Example 218

Ethyl (2*S*,4*S*)-1-{3-[2-(acetylamino)phenoxy]-2-hydroxypropyl}-4-(3,4-
10 **dichlorophenoxy)-2-pyrrolidinecarboxylate; trifluoroacetic acid salt**

i) Methyl (2*S*,4*S*)-4-(3,4-dichlorophenoxy)-2-pyrrolidinecarboxylate

The compound was prepared by analogy with Example 217(ii) from N-Boc-*cis*-4-hydroxy-L-proline methylester and 3,4-dichlorophenol.

15

¹H-NMR (400MHz, DMSO-*d*₆): δ 9.64 (bs, 2H); 7.58 (d, 1H); 7.25 (d, 1H);
6.94 (dd, 1H); 5.24 (m, 1H); 4.66 (dd, 1H); 3.76 (s, 3H); 3.55 (dd, 1H); 3.47 (d, 1H);
2.67-2.58 (m, 1H); 2.38 (d, 1H)

20 APCI-MS: m/z 290, 292 [MH⁺, isotope pattern]

ii) Ethyl (2*S*,4*S*)-1-{3-[2-(acetylamino)phenoxy]-2-hydroxypropyl}-4-(3,4-
dichlorophenoxy)-2-pyrrolidinecarboxylate; trifluoroacetic acid salt

The compound was prepared by analogy with Example 217 from compound i) (404mg,
25 1.0mmol) and *N*-[2-(2-oxiranylmethoxy)phenyl]acetamide (207mg, 1.0mmol), with the exception that ethanol was used as solvent. Reesterification occurred, and after work-up and purification the title compound was isolated as a white solid (209mg, 33%).

The substance is a diastereomeric mixture.

30 APCI-MS: m/z 511, 513 [MH⁺, isotope pattern]

Example 219

N-[2-(((2*S*)-3-[(2*S*,4*S*)-4-(4-Chlorophenoxy)-2-(hydroxymethyl)pyrrolidinyl]-2-hydroxypropyl}oxy)phenyl]acetamide; trifluoroacetic acid salt *and*

N-[2-(((2*R*)-3-[(2*S*,4*S*)-4-(4-Chlorophenoxy)-2-(hydroxymethyl)pyrrolidinyl]-2-hydroxypropyl}oxy)phenyl]acetamide; trifluoroacetic acid salt

i) [(2*S*,4*S*)-4-(4-Chlorophenoxy)pyrrolidinyl]methanol

Methyl (2*S*,4*S*)-4-(4-chlorophenoxy)-2-pyrrolidinecarboxylate

(prepared from *cis*-4-hydroxy-L-proline methylester according to Example 217ii) (850 mg, 3.32 mmol) in dry THF (20 ml) was added during 40 min to a mixture of lithium aluminiumhydride (505 mg, 13.3 mmol) in dry THF (10 ml) at 0°C under an argon atmosphere. After stirring overnight at room temperature sodium sulfate decahydrate (1g) was added, followed by dropwise addition of water (0.5 ml), sodium hydroxide (15% w/v, 0.5 ml) and water (1.5 ml). Filtration and evaporation gave a syrup which was purified by flash chromatography on silica gel (dichloromethane/methanol/concentrated ammonia 100/20/1) to give the subtitle compounds (0.60 g, 79%).

¹H-NMR (400 MHz, CDCl₃): δ 7.22 (m, 2H), 6.78 (m, 2H), 4.79 (m, 1H), 3.62 (m, 2H), 3.39 (m, 1H), 3.23 (bd, 1H), 3.14 (dd, 1H, *J* 5.0 Hz, 12.2 Hz), 2.28 (m, 1H), 1.72 (m, 1H).

MS-APCI+: *m/z* 228 [MH⁺]

ii) *N*-[2-(((2*S*)-3-[(2*S*,4*S*)-4-(4-Chlorophenoxy)-2-(hydroxymethyl)pyrrolidinyl]-2-hydroxypropyl}oxy)phenyl]acetamide; trifluoroacetic acid salt *and*

N-[2-(((2*R*)-3-[(2*S*,4*S*)-4-(4-Chlorophenoxy)-2-(hydroxymethyl)pyrrolidinyl]-2-hydroxypropyl}oxy)phenyl]acetamide; trifluoroacetic acid salt

[(2*S*,4*S*)-4-(4-Chlorophenoxy)pyrrolidinyl]methanol (380 mg, 1.67 mmol) and *N*-[2-(2-oxiranylmethoxy)phenyl]acetamide (414 mg, 2.00 mmol) were dissolved in tert-butanol (5

ml) and stirred in a sealed vial at 100°C for 3h. The solution was concentrated and the residue was purified by flash chromatography on silica gel (dichloromethane/methanol 13/1) followed by preparative RP-HPLC using acetonitrile/water containing 0.1% trifluoroacetic acid as mobile phase. The appropriate fractions were lyophilized to give the title compounds (epimer A: 248 mg, 27 %, epimer B: 115 mg, 13 %; stereochemistry at epimeric centre not determined).

Epimer A:

¹H-NMR (400 MHz, MeOD): δ 7.82 (dd, 1H, *J* 1.3 Hz, 8.0 Hz), 7.31 (m, 2H), 7.14 (m, 1H), 7.04 (m, 1H), 6.96 (m, 3H), 5.20 (m, 1H), 4.40 (m, 1H), 4.11 (bd, 2H), 3.79–4.05 (m, 5H), 3.73 (dd, 1H, *J* 5.2 Hz, 12.5 Hz), 3.43 (dd, 1H, *J* 2.6 Hz, 13.0 Hz), 2.80 (m, 1H), 2.18 (s, 3H), 2.12 (m, 1H).

MS-APCI+: *m/z* 435 [MH⁺]

Epimer B:

¹H-NMR (400 MHz, MeOD): δ 7.79 (dd, 1H, *J* 1.3 Hz, 7.9 Hz), 7.32 (m, 2H), 7.14 (m, 1H), 7.04 (m, 1H), 6.97 (m, 3H), 5.18 (m, 1H), 4.49 (m, 1H), 3.83–4.19 (m, 7H), 3.69 (dd, 1H, *J* 4.8 Hz, 13.2 Hz), 3.34 (m, 1H), 2.72 (m, 1H), 2.18 (s, 3H), 2.07 (m, 1H).

MS-APCI+: *m/z* 435 [MH⁺]

Example 220

***N*-(2-{3-[3-(4-Chlorophenoxy)-1-pyrrolidinyl]-2-hydroxy-2-methylpropoxy}phenyl)acetamide hydrochloride**

i) *N*-{2-[(2-Methyl-2-propenyl)oxy]phenyl}acetamide

The compound (1.74 g, 85 %) was prepared from 3-chloro-2-methylpropene (1.09 g, 11.9 mmol) and 2-acetamidophenol (1.50 g, 9.92 mmol) analogously to that described in Example 8(i).

¹H-NMR (400 MHz, CDCl₃): δ 8.36 (dd, 1H, *J* 1.7 Hz, 7.8 Hz), 7.80 (bs, 1H), 6.98 (m, 2H), 6.87 (dd, 1H, *J* 1.6 Hz, 7.9 Hz), 5.08 (s, 1H), 5.04 (s, 1H), 4.51 (s, 2H), 2.21 (s, 3H), 1.85 (s, 3H).

5 **ii) *N*-{2-[(2-Methyl-2-oxiranyl)methoxy]phenyl}acetamide**

The compound (0.56 g, 65 %) was prepared from *N*-{2-[(2-methyl-2-propenyl)oxy]phenyl}acetamide (800 mg, 3.90 mmol) and *m*-chloroperbenzoic acid (80 %, 1.10 g, 5.22 mmol) analogously to that described in Example 8 (ii).

10 ¹H-NMR (400 MHz, CDCl₃): δ 8.36 (m, 1H), 8.01 (bs, 1H), 7.01 (m, 2H), 6.91 (m, 1H), 4.07 (m, 2H), 2.96 (d, 1H, *J* 4.8 Hz), 2.79 (d, 1H, *J* 4.8 Hz), 2.22 (s, 3H), 1.49 (s, 3H).

iii) *N*-(2-{3-[3-(4-Chlorophenoxy)-1-pyrrolidinyl]-2-hydroxy-2-methylpropoxy}phenyl)acetamide hydrochloride

15 The title compound (255 mg, 110 %) was prepared from *N*-{2-[(2-methyl-2-oxiranyl)methoxy]phenyl}acetamide (123 mg, 0.557 mmol) and 3-(4-chlorophenoxy)pyrrolidine (100 mg, 0.506 mmol) analogously to that described in Example 1 (iii).

20 ¹H-NMR (400 MHz, MeOD): δ 7.61 (m, 1H), 7.29 (m, 2H), 7.18 (m, 1H), 7.10 (m, 1H), 6.96 (m, 3H), 5.18 (m, 1H), 3.91-4.22 (m, 4H), 3.37-3.76 (m, 4H), 2.66 (m, ½ H), 2.37 (m, 1H), 2.25 (m, ½ H), 2.15 (m, 3H), 1.45 (m, 3H).

MS-APCI+: *m/z* 419 [MH⁺]

25 **General procedures and preparation of starting materials for Examples 221-230**

Preparation of the epoxides:

30 **A) *N*-(2-[(2*S*,3*S*)-3-Methyloxiranyl]methoxy}phenyl)acetamide**

i) N-{2-[(E)-1-Propenyloxy]phenyl}acetamide

A heterogenic mixture of commercially available 1-chloro-2-butene (Aldrich, predominantly *trans*) (453 mg, 0.49 mL, 5.00 mmol), 2-acetamidophenol (756 mg, 5.00 mmol) and potassium carbonate (691 mg, 5.00 mmol) in acetone (10 ml) was heated to reflux over night. After evaporation of the solvent the residue was taken up by ethylacetate and water. Washing of the organic phase with water, 5-proz. sodium hydroxide and brine and evaporation of the solvent afforded the crude which was purified by flash chromatography on silica gel (heptane/EtOAc = 3:2). Yield: 732 mg (71 %) of a brownish yellow solid. *Trans: cis* = 81:19 (ratio determined by ¹H-NMR; 400MHz, CDCl₃):

MS-APCI+ : m/z 206.1 [MH+].

ii) *M*-chloroperbenzoic acid (70-proz.; 270 mg, 1.92 mmol, 2.0 equiv.) was added to a ice bath cooled solution of compound i) (112 mg, 546 μmol) dissolved in dichloromethane (3 ml) and stirred without further cooling over night. After addition of ethylacetate the mixture was washed with sat. sodium sulfite, 5-proz. sodium hydroxide and brine. Drying over sodium sulfate, evaporation of the solvent and flash chromatography on silica gel afforded 86 mg (71 %) of the product as a beige solid in a *trans: cis*-ratio of 83:17 as determined by ¹H-NMR.

¹H-NMR (400MHz, CDCl₃; only the signals of the major isomer are given): δ 8.35 (1H, m), 7.90 (1H, br.s), 7.00 (2H, m), 6.88 (1H, m), 4.30 (1H, dd, J 11.4, 2.5Hz), 3.96 (1H, dd, J 11.4, 5.7Hz), 3.08 (2H, m), 2.21 (3H, s), 1.40 (3H, d, J 5.2Hz).

MS-APCI+ : m/z 222.1 [MH+].

B) N-(2-[(2*S*,3*R*)-3-Methyloxiranyl]methoxy)phenyl}acetamide

i) N-[2-(2-Butynyloxy)phenyl]acetamide

A mixture of 1-bromo-2-butene (1.39 g (10.4 mmol), 2-acetamidophenol (1.58 mg, 10.4 mmol), anhydrous potassium carbonate (1.44 g, 10.4 mmol) and sodium iodide (30 mg) in butanone (50 ml) was heated over night to reflux. After that the reaction mixture was filtrated, the filtrate was evaporated and the resulting residue was taken up by ethyl acetate. The obtained solution was washed with 5-proz. sodium hydroxide, brine and water, dried over sodium sulfate and evaporated. The crude was recrystallized out of heptane/MTB (1:1) yielding in 1.57 g (74 %) of a light brown needles.

MS-APCI+ : m/z 204.1 [MH+].

ii) N-{2-[(Z)-2-Butenyloxy]phenyl}acetamide

A mixture of the alkyne i) (357 mg, 1.76 mmol) and 5-% Pd/BaSO₄ (22 mg) in pyridine (2.0 mL) was hydrogenated for 2h 30min under atmospheric pressure at room temperature. At this point the starting material was completely consumed, but some overreduction to the corresponding alkane was observed by LC/MS. The reaction mixture was filtered on celite which was thoroughly washed with ethylacetate. Thereafter the filtrate was washed with 1 N HCl to acidic reaction and finally washed neutral with brine. Drying over sodium sulfate, evaporation of the solvent yielded in 318 mg (88 %) crude which contained beside the desired Z-olefin also some E-olefine and corresponding alkane as biproducts. The ratio as determined by ¹H-NMR (400MHz, CDCl₃) was E : Z : alkane = 83 : 10 : 7. The crude was used in the next step without further purification.

MS-APCI+ : m/z 206.1 [MH+].

iii) The olefine ii) (310 mg, 1.51 mmol) was dissolved in dichloromethane (10 ml) and m-chloroperbenzoic acid (80-proz.; 587 mg, 2.72 mmol, 1.8 equiv.) was added at 0°C. Stirring over night at ambient temperature was followed by evaporation of the solvent and taking up the resulting residue with ethylacetate, washing with sat. sodium sulfite, 5-% sodium hydroxide and brine and drying over sodium sulfate. Evaporation of the solvent and flash chromatography on silica gel (ethylacetate/heptane = 2:1 continued to

ethylacetate) gave 269 mg (81 %) of the epoxide in a *E:Z*-ratio of 82:18 (determined by ¹H-NMR) as a white powder.

¹H-NMR (400MHz, CDCl₃; only the signals of the major isomer are given): δ 8.37 (1H, dd, J 7.5, 2.1Hz), 7.81 (1H, br.s), 7.02 (2H, m), 6.91 (1H, dd, J 7.4, 1.7Hz), 4.32 (1H, dd, J 11.1, 3.6Hz), 4.03 (1H, dd, J 11.1, 6.9Hz), 3.33 (1H, dt, J 7.0, 4.0Hz), 3.24 (1H, dt, J 9.9, 5.5Hz), 2.21 (3H, s), 1.38 (3H, d, J 5.7Hz).

MS-APCI+ : m/z 222.1 [MH⁺].

C) *N*-{2-[(1*R*,2*S*,5*R*)*-6-Oxabicyclo[3.1.0]hex-2-yloxy]phenyl}acetamide

i) (2-Cyclopenten-1-yloxy)(triisopropyl)silane

A solution of 2-cyclopentenol (K. Alder, F. H. Flock, *Chem. Ber.* **1956**, *89*, 1732.) (2.31 g, 27.5 mmol), (triisopropyl)chlorsilane (5.30 g, 5.9 mL, 27.5 mmol), imidazole (2.06 g, 30.2 mmol) in DMF (50 mL) was stirred at room temperature for 3h and for an additional hour at 50°C. Then the solution was diluted with ethylacetate, washed 4 times with water and dried over sodium sulfate. Evaporation of the solvent yielded in 6.32 g (96 %) of the silylether 513/13 as a colourless liquid. No major impurities were visible in the ¹H-NMR-spectrum.

¹H-NMR (400MHz, CDCl₃): δ 5.88 (1H, m), 5.76 (1H, m), 4.98 (1H, m), 2.48 (1H, m), 2.27-2.17 (2H, m), 1.70 (1H, m), 1.12-1.05 (21H, m).

ii) Triisopropyl[(1*R*,2*R*,5*R*)*-6-oxabicyclo[3.1.0]hex-2-yloxy]silane

M-chloroperbenzoic acid (70-%; 5.41 g, 21.9 mmol, 1.7 equiv.) was added to a ice bath cooled solution of compound i) (3.10 g, 12.9 mmol) dissolved in dichloromethane (25 ml) and stirred without further cooling additional 90 min. After filtration of the reaction mixture, evaporation of the filtrate the residue was dissolved in ethylacetate, washed with sat. sodium sulfite, 5-proz. sodium hydroxide and brine and dried over sodium sulfate.

Evaporation of the solvent afforded the crude as a mixture of the diastereomeric epoxides in *trans/cis*-ratio of 78:22 (¹H-NMR). Separation by flash chromatography on silica gel (heptane/ethylacetate = 95:5 continued to 90:10) afforded 1.65 g (50 %) of the desired *trans*-epoxide (1*R*,2*R*,5*R*)* as the first eluated diastereomer. The total yield of both
5 diastereomeric epoxides was 2.86 g (87 %).

¹H-NMR (400MHz, CDCl₃): δ 4.39 (1H, d, J 3.4Hz), 3.54 (1H, d, J 2.5Hz), 3.37 (1H, d, J 2.5Hz), 1.94 (1H, m), 1.84 (dtd, J 13.7, 9.3, 1.1Hz), 1.60-1.55 (2H, m), 1.13-1.04 (21H, m).

10 **iii) (1*S*,2*R*,5*R*)*-6-Oxabicyclo[3.1.0]hexan-2-ol**

To a solution of the silylether ii) (280 mg, 1.09 mmol) in THF (2.0 mL) tetrabutylammonium fluorid (1.0 M in THF; 1.2 mL, 1.20 mmol) was added. After stirring for 3 h at room temperature the mixture was diluted with ethylacetate, washed with brine and dried over sodium sulfate. Chromatographic filtration on silica gel
15 (heptane/*tert*butylmethylether 2:1 continued to ethylacetate) afforded 79 mg (72 %) as a pale yellow oil.

¹H-NMR (400MHz, CDCl₃): δ 4.36 (1H, d, J 5.1Hz), 3.55 (1H, s), 3.42 (1H, d, J 1.5Hz), 1.99 (1H, m), 1.84 (1H, dddd, J 13.9, 10.1, 9.0, 1.1Hz), 1.69-1.53 (3H, m).

20 iv) The title compound was prepared according to the general protocol (I) below.

¹H-NMR (400MHz, CDCl₃): δ 8.38 (1H, m), 7.91 (1H, br.s), 7.02-6.98 (2H, m), 6.94 (1H, m), 4.78 (1H, td, J 8.0, 1.2Hz), 3.61 (1H, m), 3.54 (1H, d, J 2.7Hz), 2.21 (3H, s), 2.21 (1H, m), 2.10 (1H, dt, J 12.8, 12.0Hz), 1.76 (1H, dtd, J 14.3, 10.4, 1.4Hz), 1.58 (1H, m).

MS-APCI+ : m/z 234.1 [MH⁺].

General protocol (I) for the preparation of *N*-{2-[(1*R*,2*S*,5*R*)*-6-oxabicyclo[3.1.0]hex-2-yloxy]phenyl}acetamides
30

Step 1 – Mitsunobu coupling:

To a ice bath cooled solution of the epoxyalcohol 546/16 (1.0 equiv.), triphenylphosphine (1.2 equiv.) and a 2-nitrophenol (1.0 equiv.) in dry THF (2 mL/mmol) diethyl
5 diazodicarboxylic acid (1.2 equiv.) was added dropwise. Stirring was continued over night without further cooling. Aqueous, basic workup, followed by flash chromatography on silica gel (typical eluant: heptane/ethyl acetate = 1:1) afforded the 2-nitrophenolic esters which contained often an equimolar amount of the biproduct diethyl 1,2-hydrazinedicarboxylate.

Step 2 – hydrogenation:

A mixture of 2-nitrophenolic esters as obtained from step 1, diisopropyl(ethyl)amine (2.0 equiv.), acetic acid anhydrate (2.0 equiv.) and 5-proz. Pt/C (10 mg/mmol) in ethyl acetate (10 mL/mmol) was hydrogenated for 1h under atmospheric pressure at room temperature.
15 In the case of non-halogenated aromates Pd/C and shorter reaction times, typically about 5 min, could be applied. Thereafter the catalysator was filtered off by a celite filled filterfunnel and washed with ethanol. The filtrate was evaporated and the remaining residue was subjected an aqueous, basic work-up. Subsequent flash chromatography on silica gel (typical eluant: ethyl acetate/heptane = 2:1) afforded the respective acetamides in typical
20 yields of 50-70 % (2 steps).

D) *N*-{5-Chloro-2-[(1*R*,2*S*,5*R*)*-6-oxabicyclo[3.1.0]hex-2-yloxy]phenyl}acetamide

Preparation according to protocol (I).

25 ¹H-NMR (400MHz, CDCl₃): δ 8.47 (1H, d, J 2.5Hz), 7.89 (1H, br.s), 6.97 (1H, dd, J 8.8, 2.5Hz), 6.85 (1H, d, J 8.8Hz), 4.74 (1H, td, J 8.0, 1.3Hz), 3.58 (1H, m), 3.56 (1H, m), 2.21 (1H, m), 2.21 (3H, s), 2.09 (dt, J 13.0, 7.4Hz), 1.76 (1H, dtd, J 14.3, 10.1, 1.3Hz), 1.56 (1H, m).

MS-APCI+ : m/z 268.0 [MH+].

E) *N*-{4-Fluoro-2-[(1*R*,2*S*,5*R*)*-6-oxabicyclo[3.1.0]hex-2-yloxy]phenyl}acetamide

Preparation according to protocol (I).

¹H-NMR (400MHz, CDCl₃): δ 8.23 (1H, dd, J 10.7, 2.8Hz), 7.99 (1H, br.s), 6.89 (1H, dd, J 9.0, 5.5Hz), 6.69 (1H, ddd, J 11.1, 9.0, 3.1Hz), 4.70 (1H, t, J 7.8Hz), 3.56 (2H, s), 2.21 (1H, dd, J 14.7, 8.4Hz), 2.21 (3H, s), 2.08 (1H, dt, J 13.0, 8.2Hz), 1.75 (1H, dtm, J 14.3, 9.5Hz).

MS-APCI+ : m/z 252.1 [MH+].

General protocol (II) for the addition of aminocycles to substituted 2-(aryloxymethyl)oxiranes

Equimolar amounts of aminocycle and epoxide, dissolved in a saturated solution of LiClO₄ in acetonitrile (1ml/100μmol), were heated to 100°C in a sealed tube. Typical reaction times ranged from 3h for open chained epoxides to 18h for oxabicyclo[3.1.0]hexanes. After cooling down to ambient temperature the reaction mixture was diluted with ethyl acetate and subjected to an aqueous work-up. The crude products were usually obtained in quantitative yields and were purified by flash chromatography on silica gel (typical eluants: ethyl acetate/methanol = 80:20).

The Examples 221-230 below were prepared according to the general protocols (I) and (II).

Example 221

***N*-(2-{(1*S**,2*R**,3*S**)-3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-cyclopentyloxy}-phenyl)-acetamide**

¹H-NMR (400MHz, CDCl₃): δ 8.20 (1H, d, J 7.4Hz), 8.07 (1H, br.s), 7.21 (2H, m), 7.01-6.96 (2H, m), 6.92 (1H, dm, J 7.4Hz), 6.77 (2H, dm, J 8.8Hz), 4.77 (1H, m), 4.54 (1H, br.q, J 4.8Hz), 4.15 (1H, m), 3.04-2.91 (3H, m), 2.81 (1H, q, J 6.8Hz), 2.62 (1H, quint, J 7.3Hz),
5 2.29 (1H, m), 2.16 (3H, s), 2.13-1.90 (5H, m), 1.63 (1H, m).

MS-APCI+ : m/z 431.2 [MH+].

Example 222

10 **N-(2-((1*R**,2*R**,3*S**)-3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-cyclopentyloxy)-phenyl)-acetamide**

¹H-NMR (400MHz, CDCl₃): δ 8.26 (1H, m), 7.90 (1H, br.d, J 9.5Hz), 7.20 (2H, m), 6.97 (2H, m), 6.88 (1H, br.d, J 7.3Hz), 6.76 (2H, m), 4.76 (1H, m), 4.50 (1H, m), 4.21 (1H, dt, J
15 14.1, 5.5Hz), 3.00-2.89 (3H, m), 2.67-2.54 (2H, m), 2.28 (1H, m), 2.15 (3H, s), 2.11 (1H, m), 1.97 (2H, m), 1.87 (2H, m).

MS-APCI+ : m/z 431.2 [MH+].

20 Example 223

N-(2-((2*R,3*R**)-3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-butoxy)-phenyl)-acetamide**

¹H-NMR (400MHz, CDCl₃): δ 8.27 ((1H, dd, J 7.6, 1.6Hz), 8.07 (1H, br.s), 7.30 (1H, d, J
25 8.8Hz), 7.04-6.92 (4H, m), 6.74 (1H, dd, J 8.8, 2.9Hz), 4.26 (1H, m), 4.23 (1H, dd J 9.9, 2.7Hz), 4.06 (1H, m), 3.96 (1H, dd, J 9.9, 8.0Hz), 2.86-2.72 (3H, m), 2.58 (1H, m), 2.47 (1H, m), 2.18 (3H, s), 1.99 (2H, m), 1.80 (2H, m), 1.12 (3H, d, J 6.9Hz).

MS-APCI+ : m/z 469.1 [MH+].

Example 224

***N*-(2-((1*S**,2*R**,3*S**)-3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-cyclopentyloxy)-phenyl)-acetamide**

5 ¹H-NMR (400MHz, CDCl₃): δ 8.26 (1H, m), 8.20 (1H, br.s), 7.19 (2H, m), 7.02 (2H, m), 6.95 (1H, m), 6.84 (2H, m), 4.49 (1H, q, J 5.2Hz), 4.31 (1H, m), 4.15 (1H, m), 2.95 (2H, q, J 7.8Hz), 2.87 (1H, m), 2.51 (2H, br.q, J 10.2Hz), 2.19 (3H, s), 2.10-1.95 (5H, m), 1.86 (2H, m), 1.60 (1H, m).

10 MS-APCI+ : m/z 445.0 [MH+].

Example 225

***N*-(2-((2*R**,3*S**)-3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-butoxy)-phenyl)-acetamide**

15 ¹H-NMR (400MHz, CDCl₃): δ 8.47 (1H, br.s), 8.36 (1H, dd, J 7.2, 1.3Hz), 7.32 (1H, d, J 9.0), 7.04-6.93 (4H, m), 6.75 (1H, dd, J 9.0, 2.9Hz), 4.34 (1H, m), 4.11 (1H, m), 3.96 (1H, dd, J 10.5, 5.2 Hz), 3.66 (1H, m), 3.00-2.80 (2H, m), 2.71 (2H, m), 2.42 (1H, m), 2.19 (3H, s), 2.05 (1H, m), 1.94-1.81 (2H, m), 1.08 (3H, d, J 6.7Hz).

20 MS-APCI+ : m/z 466.9 [MH+].

Example 226

25 ***N*-(2-((2*R**,3*R**)-3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-butoxy)-phenyl)-acetamide**

¹H-NMR (400MHz, CDCl₃): δ 8.34 (1H, t, J 4.5Hz), 8.18 (1H, br.s), 7.22 (2H, m), 6.99 (2H, m), 6.93 (1H, m), 6.76 (2H, m), 4.78 (1H, m), 4.20 (1H, m), 4.07 (1H, dt, J 10.1, 2.9 Hz), 3.95 (1H, dd, J 10.1, 8.2Hz), 3.15 (0.5H, dd, J 10.7, 6.1Hz), 2.97-2.94 (1.5H, m), 2.89

(0.5H, q, J 7.6Hz), 2.82 (0.5H, dd, J 10.5, 2.5Hz), 2.73 (0.5H, qm, J 7.5Hz), 2.65 (0.5H, m), 2.58 (1H, m), 2.26 (1H, m), 2.01 (1H, m), 1.09 (3H, appears as dd, J 6.7, 1.7Hz).

MS-APCI+ : m/z 419.1 [MH+].

5

Example 227

N-(2-((2*R*^{*},3*S*^{*})-3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-butoxy)-phenyl)-acetamide

10 ¹H-NMR (400MHz, CDCl₃): δ 8.48 (1H, m), 8.37 (1H, m), 7.22 (2H, m), 7.04-6.93 (3H, m), 6.77 (2H, m), 4.80 (1H, m), 4.12 (1H, m), 3.97 (1H, dd, J 10.5, 5.3Hz), 3.65 (1H, m), 3.09 (0.5H, dd, J 10.3, 6.0Hz), 3.06-2.99 (1.5H, m), 2.94 (0.5H, q, J 8.0Hz), 2.87-2.67 (2.5H, m), 2.25 (1H, m), 2.02 (1H, m), 1.05 (3H, appears as dd, J 6.7, 5.2Hz).

15 MS-APCI+ : m/z 418.9 [MH+].

Example 228

N-(2-((1*S*^{*},2*R*^{*},3*S*^{*})-3-[4-(3-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-cyclopentyloxy)-phenyl)-acetamide

20

¹H-NMR (400MHz, CDCl₃): δ 8.25 (1H, m), 8.17 (1H, br.s), 7.18 (1H, t, J 8.1Hz), 7.01 (2H, m), 6.97-6.90 (3H, m), 6.79 (1H, dm, J 9.0Hz), 4.48 (1H, q, J 5.5Hz), 4.34 (1H, hept, J 3.5Hz), 4.15 (1H, dd, J 7.2, 5.5Hz), 2.95 (2H, q, J 7.4Hz), 2.87 (1H, m), 2.52 (2H, br.q, J 9.6Hz), 2.19 (3H, s), 2.09-1.94 (6H, m), 1.86 (2H, m), 1.59 (1H, m).

25

MS-APCI+ : m/z 445.1 [MH+].

Example 229

N-[5-Chloro-2-((1*S*,2*R*,3*S*)^{*}-3-[4-(3,4-dichlorophenoxy)-1-piperidinyl]-2-hydroxycyclopentyl)oxy]phenyl]acetamide

30

¹H-NMR (400MHz, CDCl₃): δ 8.38 (2H, m), 7.31 (1H, d, J 8.7Hz), 6.99 (1H, d, J 8.8Hz), 6.95 (1H, dd, J 8.8, 2.6Hz), 6.86 (1H, d, J 8.8Hz), 6.75 (1H, dd, J 8.8, 2.9Hz), 4.38 (1H, q, J 4.0Hz), 4.31 (1H, hept, J 3.7Hz), 4.10 (1H, dd, J ≈7, 6Hz), 3.00 (1H, q, J 7.1Hz), 2.91-2.82 (1H, m), 2.53 (2H, m), 2.17 (3H, s), 2.08-1.93 (5H, m), 1.85 (2H, m), 1.60 (1H, m).

MS-APCI+ : m/z 513.1 [MH+].

Example 230

***N*-[4-Fluoro-2-((1*S*,2*R*,3*S*)*-3-[4-(3,4-dichlorophenoxy)-1-piperidinyl]-2-hydroxycyclopentyl)oxy]phenyl]acetamide**

¹H-NMR (400MHz, CDCl₃): δ 8.58 (1H, br.s), 8.19 (1H, dd, J 10.9, 3.2Hz), 7.31 (1H, d, J 8.8Hz), 7.00 (1H, d, J 2.9Hz), 6.89 (1H, dd, J 9.0, 5.2Hz), 6.75 (1H, dd, J 9.0, 2.9Hz), 6.67 (1H, td, J 8.8, 3.1Hz), 4.35-4.29 (2H, m), 4.09 (1H, dd, J 7.8, 5.0Hz), 3.04 (1H, q, J 7.8Hz), 2.88 (2H, m), 2.54 (2H, m), 2.18 (3H, s), 2.08 (5H, m), 1.85 (2H, m), 1.61 (1H, m).

MS-APCI+ : m/z 497.2 [MH+].

Preparation of starting materials for Examples 231-248

A) (S*R*)-1-(3,4-Dichloro-benzyl)-2,5-dimethyl-piperazine

A solution of 1,2-dichloro-4-chloromethyl-benzene (1.1ml 7.89 mmol) in DMF (5 ml) was added to 2,5-dimethyl-piperazine (1.0g, 8.77 mmol) dissolved in DMF (25 ml). The reaction was stirred over night, poured into a mixture of EtOAc and sodium carbonate (5%). The water phase was washed twice with EtOAc and the combined organic phase once with brine, and dried over sodium sulfate. After evaporation the crude was dissolved in methanol. The dibensylated piperazine does not dissolve. The filtrate was filtered through a short silica column, using methanol as eluant and evaporated to give the pure product. Yield 812mg, 38%

¹H-NMR (400MHz, DMSO-*d*₆): δ 7.56 (d, 1H, J = 8.1 Hz), 7.52 (d, 1H, J = 1.8 Hz), 7.20 (dd, 1H, J = 8.2, 1.8 Hz), 3.97 (d, 1H, J = 14.1 Hz), 3.04 (d, 1H, J = 14.2 Hz), 2.76 (dd, 1H, J = 11.9, 3.0 Hz), 2.59 (m), 2.48 (dd, 1H, J = 11.9, 2.6 Hz), 2.37 (t, 1H, J = 10.8 Hz), 2.12 (m), 1.89 (s), 1.57 (t, 1H, J = 10.4 Hz), 1.00 (d, 1H, J = 6.1 Hz), 0.82 (d, 1H, J = 6.3 Hz).

APCI-MS: m/z 273 [M⁺]

B) (S^{*}R^{*})-1-(4-Chloro-benzyl)-2,5-dimethyl-piperazine

Was synthesized in the same way as A) from 1-chloro-4-chloromethyl-benzene (1.27 g, 7.89 mmol) and 2,5-dimethyl-piperazine (1.0g, 8.77 mmol) in DMF. Yield 701 mg, 37%

¹H-NMR (400MHz, DMSO-*d*₆): δ 7.36 (d, 2H, J = 8.4 Hz), 7.30 (d, 2H, J = 8.4 Hz), 3.97 (d, 1H, J = 13.9 Hz), 3.01 (d, 1H, J = 13.8 Hz), 2.75 (dd, 1H, J = 11.9, 3.0 Hz), 2.57 (m, 1H, J = 10.8, 2.6 Hz), 2.47 (dd, 1H, J = 10.9, 2.6 Hz), 2.36 (dd, 1H, J = 11.6, 10.1 Hz), 2.10 (m, 1H), 1.88 (bs, 1H), 1.53 (t, 1H, J = 10.5 Hz), 1.01 (d, 3H, J = 6.1 Hz), 0.80 (d, 3H, J = 6.4 Hz).

APCI-MS: m/z 239 [MH⁺]

C) 1-(3,4-Chlorobenzyl)piperazine

3,4-chlorobenzyl chloride (170mg, 0.872mmol) was added to a solution of piperazine (150mg, 1.74mmol) and triethyl amine(1ml) in DMF (10ml) at room temperature. After 2hrs the solution was concentrated in vacuo. The resulting residue was triturated under ether and the obtained solid was washed with water and then dissolved in methanol and co-evaporated with toluene to give the product, 89mg, as a solid.

APCI-MS: m/z 245, 247[MH⁺]

¹HNMR (400MHz, CD₃OD) δ 7.41 (d, 1H, J=2.0 Hz), 7.37 (d, 1H, J=8.2 Hz), 7.13 (dd, 1H, J=8.2, J=2.0 Hz), 3.5 (s, 2H), 3.05 (m, 4H), 2.57 (m, 4H)

D) 1-(4-Chlorobenzyl)piperazine

Was prepared by analogy to C) above.

Example 231

N-(2-{3-[4-(3,4-Dichlorobenzyl)-1-piperazinyl]-2-hydroxypropoxy}-phenyl)acetamide dihydrochloride

A solution of N-acetyl-2-(2,3-epoxypropoxy)aniline (87.53mg, 0.422mmol) and 1-(3,4-chlorobenzyl)piperazine in ethanol (10ml 99.5%) was refluxed for 3hrs. The solvent was distilled off under reduced pressure and the resulting residue was purified by silica gel column chromatography (dichloromethane/methanol 20:1) to give the title compound as a gum. Addition of 1.0M ethereal HCl solution gave a white solid product 78mg (40%).

APCI-MS: m/z 452, 454[MH⁺]

¹HNMR (400MHz, CD₃OD) δ 8.0 (1H, dd, J=1.53Hz, J=8.01Hz), 7.5 (1H, d, J=1.91Hz), 7.45 (1H d, J=8.2Hz), 7.23 (1H, dd, J=6.1Hz, J=2.1Hz), 6.89-7.08(4H, m), 4.15(1H, m), 3.9-4.1(2H, m), 3.48(2H, S) 2.45-2.60(10H, m), 2.17(3H, S).

Examples 232-248 were synthesized according to Example 231 with the starting materials A) to D) above.

Example 232

N-(2-{3-[4-(3,4-Dichlorobenzyl)-1-piperazinyl]-2-hydroxypropoxy}-4-fluorophenyl)acetamide

APCI-MS: m/z 470[MH⁺]

Example 233

N-(2-{3-[4-(3,4-Dichlorobenzyl)-2,5-dimethyl-1-piperazinyl]-2-hydroxypropoxy}phenyl)acetamide

APCI-MS: m/z 480[MH⁺]

Example 234

N-(5-Chloro-2-{3-[4-(3,4-dichlorobenzyl)-1-piperazinyl]-2-hydroxypropoxy}phenyl)acetamide

APCI-MS: m/z 486[MH⁺]

Example 235

N-(5-Chloro-2-{3-[4-(3,4-dichlorobenzyl)-2,5-dimethyl-1-piperazinyl]-2-hydroxypropoxy}phenyl)acetamide

APCI-MS: m/z 480[MH⁺]

Example 236

N-(2-{3-[4-(3,4-Dichlorobenzyl)-2,5-dimethyl-1-piperazinyl]-2-hydroxypropoxy}-4-methylphenyl)acetamide

APCI-MS: m/z 494[MH⁺]

Example 237

N-(2-{3-[4-(3,4-Dichlorobenzyl)-2,5-dimethyl-1-piperazinyl]-2-hydroxypropoxy}-4-fluorophenyl)acetamide

APCI-MS: m/z 498[MH⁺]

Example 238

N-(2-{3 [(S^{*}R^{*})-4-(3,4-Dichlorobenzyl)-2,5-dimethyl-1-piperazinyl]-2-hydroxypropoxy}phenyl)acetamide

APCI-MS: m/z 480[MH⁺]

Example 239

N-(2-{3 [(S^{*}R^{*})-4-(4-Chlorobenzyl)-2,5-dimethyl-1-piperazinyl]-2-hydroxypropoxy}phenyl)acetamide

APCI-MS: m/z 446[MH⁺]

Example 240

N-(5-Chloro-2-{3-[(S^{*}R^{*})-4-(3,4-dichlorobenzyl)-2,5-dimethylpiperazinyl]-2-hydroxypropoxy}phenyl)acetamide

APCI-MS: m/z 514[MH⁺]

Example 241

N-(5-Chloro-2-{3-[(S^{*}R^{*})-4-(4-chlorobenzyl)-2,5-dimethylpiperazinyl]-2-hydroxypropoxy}phenyl)acetamide

APCI-MS: m/z 480[MH⁺]

Example 242

1-(5-Chloro-2-{3-[4-(4-chlorobenzoyl)-1-piperazinyl]-2-hydroxypropoxy}phenyl)-1-ethanone

APCI-MS: m/z 451[MH⁺]

Example 243

N-(5-Cyano-2-{3-[(*S***R**)-4-(3,4-dichlorobenzyl)-2,5-dimethylpiperazinyl]-2-hydroxypropoxy}phenyl)acetamide

APCI-MS: m/z 505[MH⁺]

Example 244

N-(2-{3-[(*S***R**)-4-(4-Chlorobenzyl)-2,5-dimethylpiperazinyl]-2-hydroxypropoxy}-5-cyanophenyl)acetamide

APCI-MS: m/z 471[MH⁺]

Example 245

N-(5-Chloro-2-{3-[4-(4-chlorobenzyl)-1-piperazinyl]-2-hydroxypropoxy}-phenyl)acetamide

APCI-MS: m/z 452[MH⁺]

Example 246

N-(4-Chloro-2-{3-[4-(4-chlorobenzyl)-2,5-dimethyl-1-piperazinyl]-2-hydroxypropoxy}phenyl)acetamide

APCI-MS: m/z 460[MH⁺]

Example 247

N-(2-{3-[4-(4-Chlorobenzoyl)-1-piperazinyl]-2-hydroxypropoxy}-5-cyanophenyl)acetamide

APCI-MS: m/z 457[MH⁺]

Example 248

***N*-(2-{3-[4-(4-Chlorobenzoyl)-1-piperazinyl]-2-hydroxypropoxy}-4-methylphenyl)acetamide**

APCI-MS: m/z 446[MH⁺]

Example 249

***N*-[5-Chloro-2-(((1*R*,2*S*,3*R*)-3-[(3*S*)-3-(4-chlorophenoxy)pyrrolidinyl]-2-hydroxycyclopentyl}oxy)phenyl]acetamide**

MS-APCI⁺ : m/z 464.9 [MH⁺].

[α]_D²² = - 47.6 (CH₂C₂).

Example 250

***N*-(2-[(2*S*)-3-[(3*S*)-3-[(4-Chlorophenyl)oxy]-1-pyrrolidinyl]-2-hydroxypropyl)oxy]-4-fluorophenyl)acetamide**

APCI-MS: m/z 423.1 [M⁺]

Example 251

***N*-(2-[(2*S*)-3-[(3*S*)-3-(4-Chlorobenzyl)pyrrolidinyl]-2-hydroxypropyl}oxy)phenyl)acetamide hydrochloride**

i) 3-(4-Chlorobenzyl)pyrrolidine

To a solution of 3-(4-chlorobenzyl)-2-pyrrolidinone (420mg, 2mmol) in dry THF (25mL) and under N₂ LAH (190mg, 5mmol) was added in portions with stirring over a period of a couple of minutes. The temperature was increased to 60°C and the stirring continued for 2.5h. The mixture was quenched with 200μL water, 200μL 5M NaOH and 600μL water.

The solid Li- and Al-salts were filtered off and the filtrate was evaporated to give a colourless oil (387mg, 99%).

APCI-MS: m/z 196, 198 [MH⁺]

5

ii) (2S)-1-[3-(4-Chlorobenzyl)-1-pyrrolidinyl]-3-(2-nitrophenoxy)-2-propanol

A solution of compound (i) (387mg, 2mmol) and (2S)-2-[(2-nitrophenoxy)-methyl]oxirane (390mg, 2mmol) in ethanol (6mL) was refluxed until the reaction was complete (2h), as determined by LCMS. The solvent was evaporated to give an orange oil (650mg, 83%) which was used without further purification.

10

APCI-MS: m/z 391, 393 [MH⁺]

iii) (2S)-1-(2-Aminophenoxy)-3-[3-(4-chlorobenzyl)-1-pyrrolidinyl]-2-propanol

To a solution of compound (ii) (650mg, 1.67mmol) in ethanol (10mL) at 60°C a mixture of tin(II)chloride dihydrate (2.25g, 10mmol) and 35% hydrochloric acid (2.5mL) was added. The temperature increased rapidly to 75°C. The mixture was stirred at 60°C for further 30 min. After evaporation of the solvent the residue was extracted with 5M NaOH and ether. The organic phase was washed with water, dried and evaporated. The residue was purified by RP-HPLC with acetonitrile and water containing 0.1% TFA as mobile phase. The appropriate fraction was evaporated and the residue extracted with 1M NaOH and ether. The subtitle compound was obtained from the organic phase as a colourless oil (400mg, 66%).

20

APCI-MS: m/z 361, 363 [MH⁺]

25

iv) To a solution of compound (iii) (400mg, 1.1mmol) in DCM (10mL) acetic anhydride (200μL, 2.1mmol) was added and the mixture was left overnight. After evaporation the residue was dissolved in methanol and 1.5M sodiummethoxid in methanol (2mL) was added. The mixture was left for 2h, evaporated and taken up in ether and water. A mixture

30

of the two diastereomers was obtained from the organic phase. The diastereomers were separated by HPLC on a chiral column using a mixture of isohexane, 2-propanol and methanol as mobile phase. The isolated enantiomers were dissolved in methanol (1mL), acidified with 1M hydrochloric acid (1mL), diluted with water and lyophilized to give the
5 title compounds as white amorphous solids (156mg and 173mg).

The absolute stereoisomerism was not assigned.

APCI-MS: m/z 403, 405 [MH⁺]

10 Example 252

N-(5-Chloro-2-{3-[3-(4-chloro-benzyl)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide trifluoroacetic acid salt

i) 3-(4-Chloro-benzyl)-pyrrolidin-2-one

15 In a flask was added diisopropylamine (3.22 g, 31.8 mmole) and dry THF (60 ml). The content of the flask was kept under nitrogen, and was then cooled to -76°C. To the cool solution was dropwise added n-butyllithium (n-BuLi, 32 mmole, 20 ml, 1.6 M in hexane). After completed addition, the solution was stirred for 10 minutes, and a solution of 1-Trimethylsilanyl-pyrrolidin-2-one (5.00 g, 31.8 mmole, prepared according to literature
20 methods) in dry THF (5 ml) was added dropwise. The solution was then stirred for an additional 20 minutes and a solution of 4-Chlorobenzyl chloride (5.13 g, 32 mmole) in THF (5 ml) was added via a syringe during 5 minutes. The resulting mixture was stirred at -76°C for 1 hour, and was then allowed to reach the ambient temperature and was stirred over night. Water (40 ml) was added and the mixture was stirred vigorously for 60
25 minutes. The phases were separated and the organic phase was washed with brine, and was finally evaporated, giving an oil which crystallized on standing. The solid was triturated with heptane:EtOAc 2:1 and was filtered, giving a partly purified solid. The solid was purified on silica (DCM to DCM : MeOH 99:1 to 98:2 to 97:3 gradient) giving 1.3g (20%) of the sub-title compound.

¹H-NMR (400 MHz, CDCl₃) δ: 7.27 (2H, d, *J* 8.4Hz); 7.16 (2H, d, *J* 8.4Hz); 5.43 (1H, bs); 3.31-3.13 (3H, m); 2.74-2.61 (2H, m); 2.20-2.12 (1H, m); 1.88-1.77 (1H, m)

ii) 3-(4-Chloro-benzyl)-pyrrolidine

5 In a flask was dissolved the compound obtained in a) (0.20 g, 0.95 mmole), in dry THF (10 ml). LiAlH₄ (0.17 g, 4.53 mmole) was added in small portions over 10 minutes. After completed addition, the mixture was heated to 60°C for approximately 3h under nitrogen, and the reaction was monitored by LC-MS, and was quenched after completed reaction. Before quenching, the reaction was allowed to reach the ambient temperature, and water
10 (0.160 ml) was added cautiously drop by drop. NaOH (10% solution in water, 0.16 ml) was added dropwise, and finally another portion of water (0.48 ml). The mixture was stirred for 1 hour and was then filtered. The filtrate was concentrated in vacuo giving the sub-title compound (0.18 g, 97%) as a colorless oil.

15 APCI-MS: *m/z* 196.1 [M+H]

iii) *N*-(5-Chloro-2-{3-[3-(4-chloro-benzyl)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide trifluoroacetic acid salt

20 The title compound was prepared according to the method described in Example 1 iii) from the compound obtained in ii).

¹H-NMR (400 MHz, DMSO) δ: 9.93-9.62 (1H, m); 9.12 (1H, s); 8.11 (1H, s); 7.38 (2H, d, *J* 8.9 Hz); 7.29-7.23 (2H, m); 7.13-7.02 (2H, m); 6.11-6.02 (1H, m); 4.29-4.16 (1H, bs); 4.05-3.95 (1H, m); 3.95-3.87 (1H, m); 3.75-3.50 (2H, m); 3.40-3.22 (3H, m); 2.91-2.65
25 (3H, m); 2.62-2.52 (1H, m); 2.13 (3H, s); 2.11-1.94 (1H, m); 1.81-1.55 (1H, m)

Example 253

***N*-(2-{3-[3-(4-Chlorophenoxy)-1-pyrrolidinyl]-2-hydroxypropoxy}-4-methylphenyl)-1-pyrrolidinecarboxamide trifluoroacetate**

i) 2-[(5-Methyl-2-nitrophenoxy)methyl]oxirane

A mixture of 5-Methyl-2-nitrophenol (7.7g, 50mmol), potassium carbonate (13.8g, 0.1mmol) and epibromohydrine (8.25mL, 0.1mmol) was dissolved in DMF (100mL) and stirred 2-3 h at 100°C under an atmosphere of nitrogen.

5 The mixture was diluted with ether (0.5L) and extracted with water until pH=7.

The organic phase was evaporated and the residue was purified by flash-chromatography on silica (DCM) to give the sub-title compound as a yellow solid (8.65g, 83%).

¹H-NMR (400MHz, CDCl₃): δ 7.80 (d,1H); 6.91 (s,1H); 6.86 (d,1H); 4.39 (dd,1H); 4.15 (dd,1H); 3.43-3.37 (m, 1H); 2.93 (dd,1H); 2.89 (dd,1H); 2.42 (s,3H)

ii) 1-[3-(4-Chlorophenoxy)-1-pyrrolidinyl]-3-(5-methyl-2-nitrophenoxy)-2-propanol

A mixture of compound i) (1.05g, 5.0mmol) and 3-(4-chlorophenoxy)pyrrolidine (988mg, 5.0mmol) in ethanol (12mL) was refluxed for 2h. The solvent was evaporated to give the crude product as an orange oil, which was used without further purification.

APCI-MS: m/z 407, 409 [MH⁺]

iii) 1-(2-Amino-5-methylphenoxy)-3-[3-(4-chlorophenoxy)-1-pyrrolidinyl]-2-propanol

20 To a stirred solution of compound ii) (2.1g, 5.0mmol) in ethanol (10mL) tin(II) chloride dihydrate (5.6g, 25mmol) in 35% hydrochloric acid (6mL) was added at 50°C. An exothermic reaction started and the temperature increased rapidly to 75°C.

The mixture was maintained at 60°C for 0.5h. The cooled mixture was alkalized with 1M sodium hydroxide (180mL) and extracted with ether, the organic phase washed with water, dried and evaporated to give the subtitle compound as a pale yellow oil (1.34g, 71%).

NMR: Due to a mixture of two diastereomeric pairs integration will result in parts of protons.

¹H-NMR (400MHz, CDCl₃): δ 7.23 (d, 2H); 6.77 (d, 2H); 6.67-6.65 (m, 1H);

6.64-6.63 (m, 1H); 4.83-4.76 (m, 1H); 4.14-4.06 (m, 1H); 4.01 (d, 2H); 3.71 (bs, 2H); 3.41 (bs, 1H); 3.10 (dd, 0.5H); 3.01-2.90 (m, 1.75H); 2.87-2.70 (m, 2.7H); 2.66-2.57 (m, 1.5H); 2.28 (h, 1H); 2.25 (s, 3H); 2.06-1.95 (m, 1H)

5 APCI-MS: 377, 379 [MH⁺]

iv) N-(2-{3-[3-(4-Chlorophenoxy)-1-pyrrolidinyl]-2-hydroxypropoxy}-4-methylphenyl)-1-pyrrolidinecarboxamide trifluoroacetate

A solution of compound (iii) (75mg, 0.2mmol) and di(*tert*-butyl) tricarboxylate (53mg, 0.2mmol) in DCM (3mL) was stirred for 1h at ambient temperature. Pyrrolidine (33μL, 0.4mmol) was added and the stirring continued for 1h. The reaction was complete as determined by LCMS. TFA (1mL) was added and the solution was left for 1h. The volatile parts were evaporated and the crude product purified by preparative RP-HPLC using acetonitrile and water containing 0.1% TFA as mobile phase. The appropriate fraction was concentrated *in vacuo* and the residue lyophilized to give the title compound as a white amorphous solid (85mg, 72%).

NMR: Due to a mixture of two diastereomeric pairs integration will result in parts of protons. Data are from the free base.

20 ¹H-NMR (400MHz, CDCl₃): δ 8.03 (dd, 1H); 7.24 (d, 2H); 6.94 (bs, 1H); 6.82-6.74 (m, 2H); 6.78 (d, 2H); 6.73-6.68 (m, 1H); 4.85-4.76 (m, 1H); 4.11-3.94 (m, 3H); 3.52-3.42 (m, 5.6H); 3.11 (dd, 0.5H); 3.05-2.92 (m, 0.45H); 2.95 (d, 1H); 2.87-2.72 (m, 2.5H); 2.62-2.52 (m, 1.4H); 2.37-2.21 (m, 0.7H); 2.29 (s, 3H); 2.08-1.90(m, 4.6H)

25 APCI-MS: m/z 474, 476 [MH⁺]

Example 254

N-(2-{3-[3-(4-Chlorophenoxy)-1-pyrrolidinyl]-2-hydroxypropoxy}-4-hydroxyphenyl)acetamide trifluoroacetate

To a stirred solution of the free base of compound Example 265 iii) following (128mg, 0.29mmol) in DCM (4mL) under N₂ 1M boron tribromide in DCM (0.58mL, 0.58mmol) was added at ambient temperature. The heterogeneous mixture was stirred overnight and poured into methanol. After evaporation the crude product was purified by RP-HPLC using acetonitrile and water containing 0.1% TFA as mobile phase. The appropriate fraction was lyophilized to give the title compound as a white amorphous solid (113mg, 73%).

APCI-MS: m/z 421, 423 [MH⁺]

Example 255

***N*-[2-((2*S*)-3-[4-(3,4-Dichlorophenoxy)-1-piperidinyl]-2-hydroxypropyl)oxy]-4-fluorophenyl]acetamide trifluoroacetic acid salt**

i) (2*S*)-2-[(5-Fluoro-2-nitrophenoxy)methyl]oxirane

In a flask was added (R)-glycidol (0.994 g, 13.4 mmole) and triphenylphosphine (3.52 g, 13.4 mmole) and THF (20 ml, dried over molecular sieves), and 5-fluoro-2-nitrophenol (2.10 g, 13.4 mmole). The mixture was stirred until a homogeneous solution was obtained. The solution was cooled in an ice bath and diethylazodicarboxylate (DEAD, 2.11 ml, 13.4 mmole) was added dropwise over a few minutes. After completed addition, the flask was allowed to reach room temperature and stirred for an additional 2 hours. The solvent was removed in vacuo and to the residue was added chloroform (5-10 ml). The precipitate (PPh₃O) was removed by filtration and the solid was washed with an additional amount of chloroform (5-10 ml). The filtrate was added to a flash column (SiO₂, Heptane:Ethyl acetate 4:1), and purified to give 2.02 g (71%) of the sub-title compound as a crystalline material after concentration of pure fractions.

¹H-NMR (400 MHz, CDCl₃) δ: 7.97 (1H, dd, *J* 9.3, 6.0 Hz); 6.86 (1H, dd, *J* 10.0, 2.5 Hz); 6.80-6.74 (1H, m); 4.44 (1H, dd, *J* 11.4, 2.6 Hz); 4.12 (1H, dd, *J* 11.2, 5.1 Hz); 3.44-3.38 (1H, m); 2.95 (1H, t, *J* 4.5 Hz); 2.90 (1H, dd, *J* 4.8, 2.6)

ii) (2S)-1-[4-(3,4-Dichlorophenoxy)-1-piperidinyl]-3-(5-fluoro-2-nitrophenoxy)-2-propanol

In a vial was added 4-(3,4-dichlorophenoxy)-piperidine (0.123 g, 0.5 mmole) and the compound obtained in i) (0.106 g, 0.5 mmole) and ethanol (99.5%, 3 ml). The vial was sealed and the content was heated with stirring at 65°C for 3 hours, and the reaction was monitored on LC-MS. The vial was allowed to cool and the solvent was evaporated, giving an oil, which was purified on silica (DCM to DCM:MeOH 99:1 to 98:2 to 97:3 as a stepwise gradient). Evaporation of pure fractions gave 0.22 g (96%) of the sub-title compound as an oil.

APCI-MS (m/z): 459.1 [M+H]

iii) N-[2-({(2S)-3-[4-(3,4-Dichlorophenoxy)-1-piperidinyl]-2-hydroxypropyl}oxy)-4-fluorophenyl]acetamide trifluoroacetic acid salt

The compound obtained in ii) (0.22 g, 0.48 mmole) was dissolved in ethanol (99.5%, 7 ml) and heated with stirring to 60°C. A solution of SnCl₂ x 2H₂O (0.56 g, 5 equivalents) in concentrated hydrochloric acid (0.63 ml) was added and was stirred at 60°C for 1 hour. The mixture was then allowed to cool. The solution was alkalized by the addition of excess 2M NaOH, and the solution was extracted with diethyl ether (3 x 50 ml). The combined ethereal solutions were washed with brine and evaporated. The obtained oil was dissolved in THF (8 ml), and water (8 ml) was added, followed by the addition of acetic anhydride (50 µl, 0.52 mmole). The mixture was stirred at 40°C for 15 minutes. The organic solvent was removed in vacuo, and the residue was extracted with EtOAc (3 x 30 ml). The combined organic phases were washed with brine and were then concentrated in vacuo. The residual oil was purified on preparative HPLC giving 55g (20%, 98% purity) of the title compound as the trifluoro acetate, and as a white solid after lyophilization of pure fractions.

APCI-MS (m/z): 471.0, 472.0, 473.0 and 474.0 [M+H]

Example 256

***N*-(2-(3-(4-Chloro-phenoxy)-pyrrolidin-1-yl)-2-hydroxy-propoxy)-4,6-difluorophenyl)-acetamide hydrochloride**

i) 3,5-Difluoro-6-nitrophenol

To a stirred solution of 2,3,4-trifluoronitrobenzene (5g, 28.23mmol) in dry methanol (70ml) was added a solution of sodium (0.68, 29.46) in dry methanol (30ml). The solution was stirred until all starting material was consumed (~2h). After concentration water was added and the solution was extracted with ether, dried over Na₂SO₄, filtered and concentrated to a yellow residue (4.65g). To the solution of the yellow residue in dichloromethane (140ml) was added boron tribromide (1M in dichloromethane, 40ml) and stirred at room temperature overnight. Water was then added and the solution stirred for further 30 min. The organic phase was separated and the water phase was extracted with ether. The combined organic phase were dried over Na₂SO₄, filtered and concentrated in vacuo to give a brownish residue. The residue was taken up into ether and washed with 2M sodium hydroxide and water. The water and sodium hydroxide washings were combined and neutralized with 6M HCl and extracted with ether, dried over Na₂SO₄ and evaporated to give a yellow residue which was purified by flash chromatography on silica gel with EtOAc:Heptan; 1:2 as eluent to give the desired product 2g, 11.42mmol.

GC-MS: m/z 175(M⁺)

¹HNMR (400MHz, CD₃OD) δppm, 6.63–6.68(1H, m), 6.60-6.67(1H, dt)

ii) 2-[(3,5-Difluoro-2-nitrophenoxy)methyl]oxirane

To a mixture of 3,5-difluoro-6-nitrophenol (100mg, 0.571mmol) and potassium carbonate (394mg) in DMF (5ml) was added epibromohydrin (80mg, 0.582mmol) and was stirred at 70°C for 3hr. Water and ethyl acetate were added, the organic phase separated, dried and concentrated. The resulting residue was purified by chromatography (ethyl acetate : heptan 1:3) to give the desired product as a solid 161mg (0.696mmol).

GC-MS: m/z 231 (M+)

iii) 1-[3-(4-Chlorophenoxy)-1-pyrrolidinyl]-3-(3,5-difluoro-2-nitrophenoxy)-2-propanol

A solution of 3-(4-chlorophenoxy)pyrrolidine and 2-[(3,5-difluoro-2-nitrophenoxy)methyl]oxirane (50mg, 0.216mmol) in ethanol was refluxed for 3hrs. The solvent was distilled off under reduced pressure and the resulting residue purified by silica gel column chromatography (dichloromethane/methanol 20:1) to give 45mg (0.105mmol) of the title compound as solid.

iv) N-(2-(3-(4-Chloro-phenoxy)-pyrrolidin-1-yl)-2-hydroxy-propoxy)-4,6-difluorophenyl)-acetamide hydrochloride

Platinum oxide on carbon was added to a solution of 1-[3-(4-chlorophenoxy)-1-pyrrolidinyl]-3-(3,5-difluoro-2-nitrophenoxy)-2-propanol (40mg, 0.0932mmol) in ethanol and the mixture was hydrogenated for 4hrs at 1atm. The mixture was filtered through Celite and washed several times with warm ethanol and the combined filtrate were concentrated in vacuo. To the resulting yellow residue was taken up in dichloromethane and acetic anhydride was added to the solution. The solution was stirred at room temperature for 2hrs then concentrated. Addition of 1.0M ethereal hydrogen chloride solution gave the title product as solid 20mg

APCI-MS: m/z 441[MH⁺]

Example 257

N-[2-((2S)-3-[(2S,4S)-4-(4-Chlorophenoxy)-2-methylpyrrolidinyl]-2-hydroxypropyl)oxy)-4-fluorophenyl]acetamide trifluoroacetic acid salt

The title compound was prepared according to the procedure described in Example 260 following.

¹H-NMR (400 MHz, DMSO-*d*₆) δ: 9.89 (1H, bs); 9.05 (1H, s); 7.79 (1H, dd, *J* 8.8, 6.6 Hz); 7.37 (2H, d, *J* 9.6 Hz); 7.00-6.94 (3H, m); 6.75 (1H, dt, *J* 8.6, 2.6 Hz); 6.00 (1H, bs); 5.17-5.10 (1H, m); 4.32-4.20 (1H, m); 4.05 (1H, dd, *J* 10.1, 4.6 Hz); 3.97 (1H, dd, *J* 9.9, 5.7 Hz); 3.78-3.50 (3H, m); 3.47 (1H, t, *J* 11.6 Hz); 3.17 (1H, t, *J* 13.3 Hz); 2.83 (1H, p, *J* 6.9 Hz); 2.07 (3H, s); 1.90-1.80 (1H, m); 1.42 (3H, d, *J* 6.4 Hz)

Example 258

N-[2-({(2*S*)-3-[(3*R*)-3-(4-Chlorobenzyl)pyrrolidinyl]-2-hydroxypropyl}oxy)phenyl]acetamide hydrochloride

Prepared by the method described in Example 251.

Example 259

N-{2-[(2*R*)-3-[(3*S*)-3-[(4-Chlorophenyl)oxy]-1-pyrrolidinyl]-2-hydroxypropyl]oxy}-4-fluorophenyl}acetamide

APCI-MS: *m/z* 423.1 [M⁺].

Example 260

N-[2-({(2*S*)-3-[(2*R*,4*S*)-4-(4-chlorophenoxy)-2-methylpyrrolidinyl]-2-hydroxypropyl}oxy) phenyl]acetamide trifluoroacetic acid salt

i) 1-(*tert*-Butyl) 2-methyl (2*S*, 4*R*)-4-hydroxy-1,2-pyrrolidinedicarboxylate

In a flask was dissolved (2*S*,4*R*)-4-Hydroxy-proline hydrochloride (5.4 g, 30 mmole) in a mixture of THF (200 ml), water (170 ml) and NaOH (30 ml, 2 M in water, 60 mmole). To this emulsion was added di-*tert*-butyldicarbonate (Boc₂O, 6.54 g, 30 mmole), and the mixture was stirred vigorously for 1 hour. Ether (100 ml) was added and the phases were allowed to separate. The aqueous phase was extracted with an additional 100 ml of ether.

The aqueous phase was discarded and the combined organic phases were washed with 1M

HCl (aq.) and potassium carbonate (saturated, aq.) and brine. The extract was dried with Na₂SO₄ and was concentrated in vacuo to give a residue, which was purified on silica (Heptane:EtOAc 5:1 to 3:1 to 1:1 stepwise gradient, spots visualized with I₂/MeOH). Evaporation of pure fractions were concentrated in vacuo to give 4.2 g (57%) of the sub-
5 title compound as a colorless oil.

¹H-NMR (400 MHz, CDCl₃) δ: 4.50 (1H, bs); 4.45-4.35 (1H, m); 3.74 (3H, s); 3.64 (1H, dd, *J* 11.7, 4.3 Hz); 3.59-3.42 (1H, m); 2.35-2.20 (1H, m); 2.14-2.03 (1H, m); 1.97 (1H, dd, *J* 23.3, 3.7 Hz); 1.44 (9H, d, *J* 18.9 Hz)

10 **ii) 1-(*tert*-Butyl) 2-methyl (2*S*, 4*S*)-4-(4-chlorophenoxy)-1,2 pyrrolidinedicarboxylate**

In a flask was dissolved the compound obtained in i) (2.54 g, 10.3 mmole), triphenylphosphine (2.71 g, 10.3 mmole) and 4-Chlorophenol (1.33 g, 10.3 mmole) in THF (50 ml, dried over molecular sieves) under magnetic stirring. The flask was cooled in an ice
15 bath and, to this stirred solution was added diethylazodicarboxylate (DEAD, 1.8 g, 10.3 mmole) dropwise under a few minutes. The reaction was allowed to stand over night, allowing the ice to melt and the reaction to reach room temperature. The solvents were evaporated and the residue was treated with ether (30 ml), allowing the phosphine oxide to precipitate. The solid was removed by filtration and the filtrate concentrated in vacuo. The
20 residue was purified on silica (Heptane:EtOAc 8:1 to 6:1 to 3:1, stepwise gradient. Spots on TLC were visualized by Seebach's reagent). Concentration of pure fractions gave 2.51 g (68%) of the sub-title compound as a colorless viscous oil.

¹H-NMR (400 MHz, CDCl₃) δ: 7.26-7.20 (2H, m); 6.77-6.70 (2H, m); 4.86 (1H, bs); 4.55
25 (1/2 H, dd, *J* 8.6, 2.6 Hz); 4.43 (1/2 H, dd, *J* 7.6, 3.9 Hz); 3.84-3.60 (5H, m); 2.53-2.36 (2H, m); 1.47 (9H, d, *J* 18.2 Hz)

iii) *tert*-Butyl (2*S*,4*S*)-4-(4-chlorophenoxy)-2-(hydroxymethyl)-1-pyrrolidinecarboxylate

In a flask was dissolved the compound obtained in ii) (0.951 g, 2.67 mmole) in THF (10 ml, dried over sieves). The solution was cooled in an ice bath and LiBH₄ (0.09 g, 4.07 mmole) was added. The mixture was stirred over night, allowing the ice to cool, and the solution to reach room temperature. The crude mixture was then partitioned between EtOAc (100 ml) and water (100 ml). The aqueous phase was discarded and the organic solution was washed with 0.5M HCl (aq.), NaHCO₃ (sat, aq) and brine. The solution was evaporated to give an oil which seem to be contaminated with inorganic salts. Dissolution in DCM and filtration through Celite® afforded 0.82 g (94%) of the sub-title compound as a colorless oil.

¹H-NMR (400 MHz, DMSO-*d*₆) δ: 7.33 (2H, d, *J* 9.5 Hz); 6.95 (2H, d, *J* 9.5); 4.96 (1H, bs); 4.71 (1H, bs); 3.84-3.55 (3H, m); 3.32 (2H, bs); 2.29-2.07 (2H, m); 1.41 (9H, s)

iv) *tert*-Butyl (2*S*,4*S*)-4-(4-chlorophenoxy)-2-[[[(methylsulfonyl)oxy]methyl]-1-pyrrolidine carboxylate

In a flask was dissolved the compound obtained in iii) (0.82 g, 2.5 mmole) in dichloromethane (10 ml, dried over molecular sieves). The flask was cooled on ice, and triethylamine (0.69 ml, 5.0 mmole) was added from a syringe. Methanesulfonylchloride (0.30 ml, 3.86 mmole) was added dropwise over a few minutes, and the obtained mixture was stirred over night, allowing the ice to melt. To the mixture was added DCM (60 ml), and the solution was washed with 1M HCl (aq), NaHCO₃ (sat, aq), and brine. The solution was evaporated giving 0.876 g (86%) of the sub-title compound as a yellow oil, which was used in the next step without any further purification.

¹H-NMR (400 MHz, DMSO-*d*₆) δ: 7.35 (2H, d, *J* 9.4 Hz); 6.99 (2H, d, *J* 9.4 Hz); 5.07-5.01 (1H, m); 4.37 (1H, dd, *J* 8.9, 4.2 Hz); 4.20-4.05 (2H, m); 3.71 (1H, dd, *J* 11.8, 5.0 Hz); 3.32 (2H, s); 3.15 (3H, s); 2.07 (1H, d, *J* 14.4 Hz); 1.41 (9H, s)

v) *tert*-Butyl (2*R*,4*S*)-4-(4-chlorophenoxy)-2-methyl-1-pyrrolidinecarboxylate

In a flask was dissolved the compound obtained in iv) (0.876 g, 2.16 mmole) in THF (10 ml, dried over molecular sieves). The reaction mixture was kept under an inert atmosphere and was then cooled in an ice bath. LiB(Et)₃H (1M Lithium triethylborohydride in THF, 9 ml, 9 mmole) was added with a syringe over 15 minutes. The ice bath was removed and the mixture was stirred over night. The crude mixture was partitioned between EtOAc (100 ml) and water (100 ml). The aqueous phase was removed, and the organic phase was washed with 1M HCl (aq.), NaHCO₃ (sat, aq), and brine. The solution was evaporated and the residue was purified on silica (Heptane:EtOAc 10:1 to 5:1 to 4:1 to 2:1 gradient. TLC spots were visualized by Seebach's reagent), giving 0.401 g (60%) of the sub-title compound as a colorless oil.

¹H-NMR (400 MHz, DMSO-*d*₆) δ: 7.33 (2H, d, *J* 8.7 Hz); 6.96 (2H, d, *J* 8.7 Hz); 5.00-4.94 (1H, m); 3.89 (1H, bs); 3.64 (1H, dd, *J* 12.5, 5.2 Hz); 3.38 (1H, d, *J* 12.2 Hz); 2.41-2.28 (1H, m); 1.79 (1H, d, *J* 13.7 Hz); 1.40 (9H, s); 1.23 (3H, d, *J* 6.6 Hz)

vi) (2*R*,4*S*)-4-(4-Chlorophenoxy)-2-methylpyrrolidine trifluoroacetic acid salt

In a flask was dissolved the compound obtained in v) (0.390 g, 1.25 mmole) in dichloromethane (DCM, 15 ml). To this solution was added TFA (trifluoroacetic acid, 6 ml) and the mixture was allowed to stand for 3 hours, after which the volatiles were removed in vacuo. The residue was co-evaporated twice with DCM, giving the sub-title compound as an oil.

APCI-MS (*m/z*): 212 [M+H]

vii) *N*-[2-({(2*S*)-3-[(2*R*,4*S*)-4-(4-Chlorophenoxy)-2-methylpyrrolidinyl]-2-hydroxypropyl}oxy) phenyl]acetamide trifluoroacetic acid salt

The title compound was prepared according to the method outlined in Example 255 starting from the material obtained in vi) and (2*S*)-2-[(2-nitrophenoxy)methyl]oxirane. The compound was obtained in 25% yield.

¹H-NMR (400 MHz, DMSO-*d*₆) δ: 9.88 (1H, bs); 9.02 (1H, s); 7.89 (1H, d, *J* 7.7 Hz); 7.37 (2H, d, *J* 7.7 Hz); 7.09-6.88 (5H, m); 6.02 (1H, bs); 5.18-5.11 (1H, m); 4.34-4.22 (1H, m); 4.02 (1H, dd, *J* 10.2, 4.3 Hz); 3.94 (1H, dd, *J* 9.8, 5.7 Hz); 3.77-3.30 (4H, m); 3.19 (1H, t, *J* 10.7 Hz); 2.84 (1H, p, *J* 6.7 Hz); 2.09 (3H, s); 1.91-1.81 (1H, m); 1.43 (3H, d, *J* 6.4 Hz)

APCI-MS (*m/z*): 419.2 [M+H]

Example 261

N-{2-[(2S)-(3-[(3R)-3-[(4-Chlorophenyl)oxy]-1-pyrrolidinyl)-2-hydroxypropyl]oxy]-4-fluorophenyl}acetamide

APCI-MS: *m/z* 423.1 [M+]

Example 262

N'-(2-{3-[3-(4-Chlorophenoxy)-1-pyrrolidinyl]-2-hydroxypropoxy}-4-methylphenyl)-N,N-dimethylurea trifluoroacetate

The title compound was prepared by analogy to the methods described in Example 253 starting from compound iii) (75mg, 0.2mmol) and dimethylamine (2M in THF, 200μL, 0.4mmol). The substance was obtained as a white amorphous solid (73mg, 65%).

¹H-NMR (400MHz, MeOH-*d*₄): δ 7.54 (dd, 1H); 7.51 (d, 2H); 7.16 and 7.15 (d, 2H); 7.05 (bs, 1H); 6.96 (bd, 1H); 5.40-5.35 (m, 1H); 4.54-4.46 (m, 1H); 4.27 (d, 2H); 4.16-3.56 (m, 6H); 3.20 (bs, 6H); 2.84-2.59 (m, 1H); 2.59-2.44 (m, 1H); 2.50 (s, 3H).

APCI-MS: *m/z* 448, 450 [MH⁺]

Example 263

N-(2-{3-[3-(4-Chloroanilino)-1-pyrrolidinyl]-2-hydroxypropoxy}phenyl)acetamide

i) tert-Butyl-3-hydroxy-1-pyrrolidinecarboxylate

To a solution of 3-hydroxy-1-pyrrolidine (871 mg, 10 mmol) in THF (30 mL) was added di-(tert.butyl) dicarbonate (2.18 g, 10 mmol) in THF (2 mL) and the reaction mixture kept on stirring at room temperature for overnight. The solvent was removed in vacuo and the residue was purified by flash chromatography (0-2% MeOH in CHCl₃) to give the subtitled compound (1.7g).

¹H-NMR (CDCl₃, 400 MHz): δ 4.45 (m, 1H); 3.55-3.25 (m, 4H); 2.18-1.85 (m, 3H); 1.45 (s, 9H).

APCI-MS: m/z 166 (M-Boc).

ii) tert-Butyl-2-oxo-1-pyrrolidinecarboxylate

Chromium (vi) oxide (800 mg, 8.0 mmol) was added to pyridine (1.6 mL) in CH₂Cl₂ (10 mL) and the resulting solution was stirred for 15 min at room temperature. A solution of tert.butyl-3-hydroxy-1-pyrrolidinecarboxylate (374.5 mg, 2.0 mmol) in CH₂Cl₂ (5 mL) was added, immediately followed by acetic anhydride and the reaction mixture kept at room temperature for 15 min. After addition of ethyl acetate, decanted and filtered through a short column of silica gel. The filtrate was concentrated to give the subtitled product (193 mg) and was used directly in the next step.

iii) tert-Butyl-3-(4-chloroanilino)-1-pyrrolidinecarboxylate

Tert.butyl-2-oxo-1-pyrrolidinecarboxylate (190 mg, 1.02 mmol), 4-chloroaniline (64 mg, 0.5 mmol) and acetic acid (184 mg) were mixed in dichloroethane (5 mL). Sodium triacetoxyborohydride (326.5 mg) was added and the reaction mixture kept on stirring at room temperature for overnight. After addition of aq. NaHCO₃ the reaction mixture was diluted by addition of ethyl acetate. Two layers were separated. The organic layer was dried over Na₂SO₄, filtered, concentrated. The residue was purified by flash chromatography (0-15% ethyl acetate in petroleum spirit, 40-60) to give the subtitled product (140 mg).

¹H-NMR (CDCl₃, 400 MHz): δ 7.18 (m, 2H); 6.50 (m, 2H); 3.99 (m, 1H); 3.70 (m, 2H); 3.46 (m, 2H); 3.20 (m, 1H); 2.18 (m, 1H); 1.87 (m, 1H); 1.45 (s, 9H).
APCI-MS: m/z 197 (M-Boc).

5 **iv) N-(4-Chlorophenyl)-3-pyrrolidinamine (2xCF₃COOH)**

To a solution of tert.butyl-3-(4-chloroanilino)-1-pyrrolidinecarboxylate (130 mg, 0.438 mmol) in CH₂Cl₂ (5 mL) was added CF₃COOH (1 mL). After 30 min the volatiles were removed in vacuo to give the subtitled product (186 mg) and was used directly in the next step.

10 **v) N-(2-{3-[3-(4-Chloroanilino)-1-pyrrolidinyl]-2-hydroxypropoxy}phenyl)acetamide**

A mixture of N-(4-chlorophenyl)-3-pyrrolidinamine (2xCF₃COOH), (186 mg, 0.438 mmol), N-[2-(2-oxiranylmethoxy)phenyl]acetamide (91 mg, 0.438 mmol) and K₂CO₃ (200 mg) in ethanol (6 mL) was kept on stirring at 65 °C for 2.5 h. The volatiles were removed in vacuo. The residue was partitioned between ethyl acetate and aq. NH₄Cl solution. The organic layer was washed with water, dried over Na₂SO₄, filtered and concentrated. The residue was purified by flash chromatography (0-3% MeOH in CHCl₃) to give the titled product (70 mg).

15 ¹H-NMR (CDCl₃, 400 MHz): δ 8.35 (m, 1H); 8.21 (br.s, 1H); 7.12 (m, 2H); 7.01 (m, 2H); 6.92 (m, 1H); 6.48 (m, 2H); 4.13-3.92 (m, 4H); 3.84 (br. s, 1H); 2.99 (m, 1H); 2.87-2.30 (m, 6H); 2.18 (s, 3H); 1.66 (m, 1H).

APCI-MS: m/z 446 (MH⁺).

25 **Example 264**

***N*-{2-[(3-{3-[(4-Chlorophenyl)oxy]-1-pyrrolidinyl}-2-hydroxy-1-methylpropyl)oxy]phenyl}acetamide hydrochloride**

30 **i) N-{2-[(1-Methyl-2-propenyl)oxy]phenyl}acetamide**

The compound (557 mg, 40%) was prepared from 3-chloro-1-butene (747 ml, 7.42 mmol) and 2-acetamidophenol (1.02 g, 6.75 mmol) analogously to that described in Example 8 i).

¹H-NMR (400 MHz, CDCl₃): δ 8.37 (m, 1H), 7.80 (bs, 1H), 6.94 (m, 3H), 5.93 (m, 1H),
5.25 (m, 2H), 4.84 (m, 1H), 2.21 (s, 3H), 1.49 (d, *J* 6.3 Hz, 3H).

ii) *N*-{2-[1-(2-Oxiranyl)ethoxy]phenyl}acetamide

The compound was prepared from *N*-{2-[(1-Methyl-2-propenyl)oxy]phenyl}acetamide (549 mg, 2.67 mmol) and *m*-chloroperbenzoic acid (80 %, 923 mg, 4.28 mmol)

analogously to that described in Example 8 (ii). Purification was done on silica gel with petroleum ether/ethyl acetate 10/15 as eluent. This gave separation of the two diastereomeric pairs.

Diastereomer 1: (53 mg, 9%), R_f=0.27

Diastereomer 2: (406 mg, 69 %), R_f=0.20

¹H-NMR (400 MHz, CDCl₃): δ 8.39 (m, 1H), 8.01 (bs, 1H), 7.00 (m, 3H), 3.98 (m, 1H), 3.24 (m, 1H), 2.94 (t, *J* 4.5 Hz, 1H), 2.71 (dd, *J* 2.6 Hz, *J* 4.5 Hz, 1 H), 2.23 (s, 3H), 1.47 (d, *J* 6.3 Hz, 3 H).

iii) *N*-{2-[(3-{3-[(4-Chlorophenyl)oxy]-1-pyrrolidinyl}-2-hydroxy-1-methylpropyl)oxy]phenyl}acetamide hydrochloride

The title compound (230 mg, 100 %) was prepared from *N*-[2-(1-oxiranylethoxy)phenyl]acetamide diastereomer 2 (123 mg, 0.557 mmol) and 3-(4-chlorophenoxy)pyrrolidine (100 mg, 0.506 mmol) analogously to that described in Example 1 (iii).

¹H-NMR (400 MHz, MeOD): δ 7.86 (m, 1H), 7.30 (m, 2H), 7.09 (m, 2H), 6.97 (m, 3H), 5.21 (m, 1H), 4.51 (m, 1H), 3.83-4.22 (m, 3H), 3.37-3.62 (m, 4H), 2.68 (m, ½ H), 2.38 (m, 1H), 2.27 (m, ½ H), 2.19 (m, 3H), 1.32 (m, 3H).

MS-APCI+: m/z 419 [MH⁺]

Example 265

***N*-(2-{3-[3-(4-Chlorophenoxy)-1-pyrrolidinyl]-2-hydroxypropoxy}-4-methoxyphenyl)acetamide hydrochloride**

i) 1-[3-(4-Chlorophenoxy)-1-pyrrolidinyl]-3-(5-methoxy-2-nitrophenoxy)-2-propanol

The subtitle compound was prepared in analogy of Example 253 ii) from

2-[(5-methoxy-2-nitrophenoxy)methyl]oxirane (320mg, 1.6mmol) and

3-(4-chlorophenoxy)pyrrolidine (365mg, 1.6mmol). The crude product was obtained as a yellow oil (580mg) and was used without further purification.

APCI-MS: m/z 423, 425 [MH⁺]

ii) 1-(2-Amino-5-methoxyphenoxy)-3-[3-(4-chlorophenoxy)-1-pyrrolidinyl]-2-propanol

The subtitle compound was prepared in analogy of Example 253 iii) from

compound i) (290mg, 0.7mmol). The crude compound was obtained as a colourless oil (233mg, 85%) and was used without further purification.

APCI-MS: m/z 393, 395 [MH⁺]

iii) *N*-(2-{3-[3-(4-Chlorophenoxy)-1-pyrrolidinyl]-2-hydroxypropoxy}-4-methoxyphenyl)acetamide hydrochloride

To a solution of compound (ii) (157mg, 0.4mmol) in pyridine (3mL) acetic anhydride (1mL) was added. The mixture was stirred for 1h at ambient temperature. After evaporation the residue was dissolved in methanol (mL) and 1.5M sodium methoxide in methanol (1mL) was added. The mixture was left overnight at ambient temperature. After evaporation the residue was taken up in ether and water. The free base of the title compound was obtained from the organic phase as a colourless oil (171mg, 98%). The free

base (43mg) was dissolved in methanol (5mL), acidified with 1M hydrochloric acid till pH<2, diluted with water (50mL) and lyophilized. The title compound was obtained as a white amorphous solid (30mg, 64%).

5 APCI-MS: m/z 435, 437 [MH⁺]

Example 266

N-(2-[3-(4-Chloro-benzyloxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy)-phenyl)-acetamide trifluoroacetic acid salt.

10

i) 3-(4-Chloro-benzyloxy)-pyrrolidine-1-carboxylic acid tert-butyl ester

A solution of tert-butyl 3-hydroxy-1-pyrrolidinecarboxylic acid (0.27 g, 1.44 mmol) in dry THF (4 mL) was added dropwise to a cold (0° C), stirred suspension of sodium hydride (0.078 g, 2.17 mmol, ca. 50% suspension in oil) in THF (10 mL). After 30 min. a solution of 4-chlorobenzyl bromide (0.36 g, 1.74 mmol) in THF (2 mL) was added and the resulting suspension was stirred at R.T. overnight. The reaction mixture was partitioned between water and ethyl acetate. The aqueous phase was extracted with ethyl acetate and the combined organic phase was washed with saturated aqueous sodium chloride, dried and concentrated. The residue was subjected to flash chromatography (heptane-ethyl acetate, 6:1) to afford the subtitle compound 3-(4-Chloro-benzyloxy)-pyrrolidine-1-carboxylic acid tert-butyl ester as an oil (0.30 g, 66.8%).

15
20

¹H-NMR (CDCl₃): δ 7.30 (m, 4H), 4.49 (bs, 2H), 4.11 (m, 1H), 3.45 (m, 4H), 1.90-2.08 (m, 2H), 1.46 (s, 9H).

25

ii) 3-(4-Chloro-benzyloxy)-pyrrolidine

A solution of 3-(4-Chloro-benzyloxy)-pyrrolidine-1-carboxylic acid tert-butyl ester (0.28 g, 0.9 mmol) in aqueous 90% formic acid (7.5 mL) was stirred at (0° C) for 30 min. then at room temperature overnight. The solvents were removed under reduced pressure and the residue was treated with saturated aqueous potassium carbonate and extracted twice with n-

30

butanol. The combined organic extracts was concentrated and the residue was purified by flash chromatography (SiO₂, dichloromethane-methanol-ammonium hydroxide, 8:8:1 then 50:10:1) to afford the subtitle compound 3-(4-Chloro-benzyloxy)-pyrrolidine (0.13 g, 70%).

¹H-NMR (DMSO-d₆): δ 7.32-7.41 (m, 4H), 4.42 (s, 2H), 4.02 (m, 1H), 3.18 (bs, 3H), 2.75-2.86 (m, 3H), 2.68 (m, 1H), 1.66-1.81 (m, 2H).

APCI-MS: m/z 212 [MH⁺].

iii) N-(2-[3-(4-Chloro-benzyloxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy)-phenyl)-acetamide trifluoroacetic acid salt

A solution of 3-(4-chloro-benzyloxy)-pyrrolidine (0.050 g, 0.24 mmol) and N-(2-oxiranylmethoxy-phenyl)-acetamide (0.049 g, 0.24 mmol) in absolute ethanol (3 mL) was heated in a closed vial at 70° C for 2h. The product was purified by HPLC to afford the title compound (0.60 g, 47%).

¹H-NMR (CD₃OD): δ 7.85 (m, 1H), 7.35 (m, 4H), 7.12 (m, 1H), 7.02 (d, 1H, J = 8 Hz), 6.96 (m, 1H), 4.56 (m, 2H), 4.39 (m, 2H), 4.05 (d, 2H, J = 5.9 Hz), 3.85 (m, 2H), 3.48 (m, 4H), 2.10-2.55 (m, 5H).

APCI-MS: m/z 419 [MH⁺] and 421 [MH+2⁺].

Example 267

N-(2-[3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-2-methyl-propoxy}-phenyl)-acetamide

The compound was prepared by a method analogous to that of Example 270 following.

APCI-MS: m/z 419 [MH⁺]

Example 268

N-(2-((1*S*,2*R*,3*S*)*-3-[(3*S*)-3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-cyclopentyloxy)-5-chloro-phenyl)-acetamide (diastereomeric mixture)

Was prepared by analogy to Example 271 following from *N*-(5-chloro-2-[(1*R*,2*S*,5*R*)*-6-oxabicyclo[3.1.0]hex-2-yloxy]phenyl)acetamide (5.3 mg, 20 μ mol) and (3*S*)-3-(3,4-difluoro-phenoxy)-pyrrolidine (4.0 mg, 20 μ mol).

MS-APCI+ : m/z 467 [M+].

Example 269

N-(2-((2*R*,3*S*)*-3-[(3*S*)-3-(4-Chlorophenoxy)pyrrolidinyl]-2-hydroxybutyl)oxy)-4-methylphenyl]acetamide (diastereomeric mixture)

Was prepared analogies to Example 271 following from *N*-(4-methyl-2-[(2*S*,3*R*)*-3-methyloxiranyl]methoxy)phenyl)acetamide (4.7 mg, 20 μ mol) and (3*S*)-3-(4-Chloro-phenoxy)-pyrrolidine (4.0 mg, 20 μ mol).

MS-APCI+ : m/z 433 [M+].

Example 270

N-(2-[(3-{4-[(3,4-Dichlorophenyl)oxy]-1-piperidinyl}-2-hydroxy-2-methylpropyl)oxy]-4-fluorophenyl)acetamide hydrochloride

i) N-[4-Fluoro-2-(2-methyl-allyloxy)-phenyl]-acetamide

3-Chloro-2-methylpropene (1.36g, 15 mmol) was added to a mixture of 5-fluoro-2-nitro-phenol (1.57 g, 10 mmol), potassium carbonate (2.76 g, 20 mmol), tetrabutylammonium hydrogen sulfate (0.068 g, 0.2 mmol) and acetonitrile (30 ml), and the mixture was heated under reflux for 18 h. The cold reaction mixture was diluted with toluene and washed with

5 % aqueous potassium carbonate, dried and evaporated. A part of the residue (0.631 g, 3 mmol), sodium dithionite (1.04 g, 6 mmol) in EtOH-THF-H₂O (2:1:1, 3 ml) was heated at 75 °C for 4 h. The mixture was portioned between dichloromethane and 15 % aqueous potassium carbonate and the organic solution dried and concentrated. The obtained residue was diluted with methanol (1.5 ml) and reacted with acetic anhydride (1.5 ml) at 50 °C for 2 min and allowed to attend room temperature during 20 min, then pyridine (4 ml) was added and the solution heated again at 50 °C for 3 min, cold and concentrated. The material was purified by silica gel chromatography (light petroleum-ethyl acetate 2:1) to give 95 mg of the subtitle compound.

¹H-NMR (300MHz, CDCl₃): δ 8.28 (dd, 1H), 7.62 (bs, 1H), 6.70-6.60 (m, 2H), 5.07 (dd, 2H), 4.49 (s, 2H), 2.20 (s, 3H), 1.84 (s, 3H).

ii) N-(4-Fluoro-2-(((2-methyl-2-oxiranyl)methyl)oxy)phenyl)acetamide

The subtitle compound was prepared from N-[4-fluoro-2-(2-methyl-allyloxy)-phenyl]-acetamide analogously as described in Example 8 ii).

¹H-NMR (300MHz, CDCl₃): δ 8.31-8.26 (dd, 1H), 7.79 (bs, 1H), 6.75-6.65 (m, 2H), 4.14 (d, 1H), 3.97 (d, 1H), 2.93 (d, 1H), 2.80 (d, 1H), 2.21 (s, 3H), 1.50 (s, 3H).

APCI-MS: m/z 240 [MH⁺]

iii) N-{2-[(3-{4-[(3,4-Dichlorophenyl)oxy]-1-piperidinyl}-2-hydroxy-2-methylpropyl)oxy]-4-fluorophenyl}acetamide hydrochloride

A solution of 4-(3,4-dichloro-phenoxy)-piperidine (36 mg, 0.146 mmol), N-(4-fluoro-2-(((2-methyl-2-oxiranyl)methyl)oxy)phenyl)acetamide (35 g, 0.146 mmol) in EtOH (1 ml, 95 %) was stirred for 2.5 hours at 77 °C in a sealed vial. The solvent was evaporated and the residue was purified on silica (dichloromethane-methanol, 15:1, containing 1% of NH₄OH (25%)) to give 45 mg of the corresponding free amine of the title compound.

¹H-NMR of the corresponding free amine of the title compound.

(400MHz, CDCl₃): δ 8.26-8.22 (dd, 1H), 7.89 (bs, 1H), 7.31 (d, 1H), 7.01 (d, 1H), 6.77-6.65 (m, 3H), 4.30 (m, 1H), 3.80 (dd, 2H), 2.93-2.81 (m, 2H), 2.67 (d, 1H), 2.63-2.51 (m, 2H), 2.45 (d, 1H), 2.19 (s, 3H), 1.96 (m, 2H), 1.83 (m, 2H), 1.62 (bs, 1H), 1.31 (s, 3H).

5

iv) N-{2-[(3-{4-[(3,4-Dichlorophenyl)oxy]-1-piperidinyl}-2-hydroxy-2-methylpropyl)oxy]-4-fluorophenyl}acetamide hydrochloride

A solution of the free amine in methanol (10 ml) was acidified with HCl (conc., 0.020 ml) to pH 3 and concentrated. The residue was coevaporated three times with toluene to give
10 the title hydrochloride compound as a white powder.

APCI-MS: m/z 485, 487 [MH⁺]

Example 271

15 N-(2-[(1*S*,2*R*,3*S*)*-3-[(3*S*)-3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-cyclopentyloxy]-4-fluoro-phenyl)-acetamide (diastereomeric mixture)

N-{4-fluoro-2-[(1*R*,2*S*,5*R*)*-6-oxabicyclo[3.1.0]hex-2-yloxy]phenyl}acetamide
(5.0 mg, 20 μmol) and (3*S*)-3-(4-Chloro-phenoxy)-pyrrolidine (3.9 mg, 20 μmol) were
20 dissolved in a 2M solution of LiClO₄ in acetonitrile (0.2 ml) and heated in a sealed tube to 100°C. Dilution by ethyl acetate, neutral aqueous workup and evaporation of the solvent gave a crude product which was used without further purification.

MS-APCI⁺ : m/z 449 [M⁺].

25

Example 272

N-(5-Chloro-2-{3-[3-(3,4-difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide

30 APCI-MS: m/z 441.1 [MH⁺]

Example 273

N-(5-Chloro-2-{3-[3-(4-fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide

APCI-MS: m/z 423.1 [MH⁺]

Example 274

N-(4-Cyano-2-{3-[4-(3,4-dichloroanilino)-1-piperidinyl]-2-hydroxypropoxy}phenyl)acetamide

APCI-MS: m/z 477 [MH⁺]

Example 275

N-(4-Hydroxy-2-[(1*S*,2*R*,3*S*)*-3-[(3*S*)-3-(4-chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-cyclopentyloxy]-phenyl)-acetamide (diastereomeric mixture)

i) *N*-{4-methoxy-2-[(1*R*,2*S*,5*R*)*-6-oxabicyclo[3.1.0]hex-2-yloxy]phenyl}acetamide (32 mg, 122 μmol) and (3*S*)-3-(4-Chloro-phenoxy)-pyrrolidine (24 mg, 122 μmol) were dissolved in a 2M solution of LiClO₄ in acetonitrile (1 ml) and heated in a sealed tube to 100°C. Dilution by ethylacetate, neutral aqueous workup and evaporation of the solvents gave 62 mg (110%) of the crude addition product which was reacted with bortribromide (1M in CH₂Cl₂, 0.37 mL, 371 μmol) in dichloromethane (1 mL) at room temperature over night. The reaction was quenched with methanol (1.0 mL) and all volatile components were evaporated. The remaining crude was subjected to a reversed phase HPLC giving 30 mg (54 %) of the title compound as a diastereomeric mixture.

MS-APCI⁺ : m/z 447.1 [MH⁺].

ii) Separation of the diastereomers

The above under i) described diastereomeric mixture was subjected to chiral phase HPLC (stationary phase: Chiralpak AD; mobile phase: *iso*-hexane/*iso*-propanol/methanol/diethylamine = 80:16:4:0.1) with the compound of Example 276 as the first and the compound of Example 277 as the second eluted stereoisomer. The assignment of the absolute configuration of the respective stereoisomer beneath is set by will and exchangeable.

Example 276

10 *N*-(4-Hydroxy-2-{(1*S*,2*R*,3*S*)-3-[(3*S*)-3-(4-chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-cyclopentyloxy}-phenyl)-acetamide

¹H-NMR (400MHz, CDCl₃; OH-protons are neglected): δ 8.02 (1H, s), 7.49 (1H, d, J 8.4Hz), 7.17 (2H, d, J 8.9Hz), 6.71 (2H, d, J 8.8Hz), 6.43 (1H, s), 6.34 (1H, d, J 7.2Hz),
15 4.76 (1H, m), 4.39 (1H, m), 4.09 (1H, m), 3.10-2.95 (3H, m), 2.89 (1H, m), 2.77 (1H, m), 2.24 (1H, m), 2.08 (3H, s), 2.10-1.84 (3H, m), 1.75 (1H, m), 1.59 (1H, m).

MS-APCI+ : m/z 447.1 [MH⁺].

[α]²² = + 49.5 (CH₂C₂).

20 Example 277

N-(4-Hydroxy-2-{(1*R*,2*S*,3*R*)-3-[(3*S*)-3-(4-chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-cyclopentyloxy}-phenyl)-acetamide

¹H-NMR (400MHz, CDCl₃; OH-protons are neglected): δ 7.75 (1H, s), 7.60 (1H, d, J 8.4Hz), 7.19 (2H, d, J 8.3Hz), 6.73 (2H, d, J 8.6Hz), 6.57 (1H, s), 6.38 (1H, d, J 8.4Hz),
25 4.77(1H, m), 4.43 (1H, m), 4.21 (1H, m), 3.09-2.94 (3H, m), 2.79 (1H, m), 2.68 (1H, m), 2.28 (1H, m), 2.08 (3H, s), 2.05-1.90 (3H, m), 1.86 (1H, m), 1.53 (1H, m).

MS-APCI+ : m/z 447.1 [MH⁺].

[α]²² = - 45.2 (CH₂C₂).

The diastereomers of Examples 278 and 279 were prepared by methods analogous to those used to prepare the compounds of Examples 221-230 and separated as described in Example 275 above. The absolute configuration of the respective isomers is assigned by will as mentioned above and therefore exchangeable.

5 **Example 278**

***N*-[2-({(1*S*,2*R*,3*S*)-3-[(3*S*)-3-(4-Chlorophenoxy)pyrrolidinyl]-2-hydroxycyclopentyl}oxy)phenyl]acetamide**

10 First eluted isomer.

MS-APCI+ : m/z 431.1 [MH+].

$[\alpha]^{22} = + 72.2$ (CH₂C₂).

15 **Example 279**

***N*-[2-({(1*R*,2*S*,3*R*)-3-[(3*S*)-3-(4-Chlorophenoxy)pyrrolidinyl]-2-hydroxycyclopentyl}oxy)phenyl]acetamide**

Second eluted isomer.

20 MS-APCI+ : m/z 431.1 [MH+].

$[\alpha]^{22} = - 51.4$ (CH₂C₂).

The diastereomers of Examples 280 and 249 were prepared by methods analogous to those used to prepare the compounds of Examples 221-230 and separated as described in Example 275 above. The compound of Example 280 is the first eluted isomer whilst the compound of Example 249 is the second eluted isomer. The absolute configuration of the respective isomers is assigned by will as mentioned above and therefore exchangeable.

30 **Example 280**

N-[5-Chloro-2-((1*S*,2*R*,3*S*)-3-[(3*S*)-3-(4-chlorophenoxy)pyrrolidinyl]-2-hydroxycyclopentyl)oxy]phenyl]acetamide

MS-APCI+ : m/z 464.9 [MH+].

5 $[\alpha]^{22} = + 53.0$ (CH₂C₂).

Example 281

N-{5-Chloro-2-(((1*S*,2*R*,3*S*)*-3-{[1-(4-chlorobenzyl)-4-piperidinyl]amino}-2-hydroxycyclopentyl)oxy]phenyl}acetamide(racemic mixture)

10 Was prepared by analogy to Example 271 from *N*-{5-chloro-2-[(1*R*,2*S*,5*R*)*-6-oxabicyclo[3.1.0]hex-2-yloxy]phenyl}acetamide (5.3 mg, 20 μ mol) and 1-(4-chlorobenzyl)-4-piperidinamine (4.5 mg, 20 μ mol).

15 MS-APCI+ : m/z 492 [M+].

Example 282

N-[2-((2*S*)-3-[(3*S*)-3-(4-Chlorophenoxy)pyrrolidinyl]-2-hydroxypropyl)oxy)-4-hydroxyphenyl]acetamide

i) (2*S*)-2-[(5-Methoxy-2-nitrophenoxy)methyl]oxirane

20 The subtitle compound was prepared under Mitsunobu conditions from R-(+)-glycidole (198mg, 1mmol), 5-methoxy-2-nitrophenol (169mg, 1mmol), triphenylphosphine (263mg, 1mmol) and DEAD (157 μ L, 1mmol) using dry THF as solvent. The crude material was
25 purified by flashchromatography on silica using mixtures of ethylacetate and heptane as mobile phase. The appropriate fractions were pooled to give impure product as white crystals (175mg). The product was contaminated with reduced DEAD in molar ratio 1:1, which is equal to a yield of the desired product of 100mg, 44%.

30 ¹H-NMR (400MHz, CDCl₃): δ 8.00 (d, 1H); 6.60 (d, 1H); 6.55 (dd, 1H);

6.4 (bs, 1H, *red.DEAD*); 4.41 (dd, 1H); 4.22 (q, 4H, *red.DEAD*); 4.13 (dd, 1H);
3.89 (s, 3H); 3.44-3.39 (m, 1H); 2.95 (dd, 1H); 2.92 (dd, 1H); 1.29 (t, 6H, *red.DEAD*)

APCI-MS: m/z 226 [MH⁺]

5
ii) (2S)-1-[(3S)-3-(4-Chlorophenoxy)pyrrolidinyl]-3-(5-methoxy-2-nitrophenoxy)-2-propanol

The subtitle compound was prepared by analogy to Example 1(ii) from (i)
(169mg, 0.43mmol) and (3S)-3-(4-chlorophenoxy)pyrrolidine (85mg, 0.43mmol). The
10 product was obtained as a yellow oil and was used without further purification.

APCI-MS: m/z 423, 425 [MH⁺]

15
iii) (2S)-1-(2-Amino-5-methoxyphenoxy)-3-[(3S)-3-(4-chlorophenoxy)pyrrolidinyl]-2-propanol

The subtitle compound was prepared in analogy of Example 253 iii) from (ii)
(0.43mmol). The product obtained (colourless oil, 163mg) was a mixture of the desired
product and reduced DEAD in molar ratio 5:1. The substance was used as it was.

20 ¹H-NMR (400MHz, CDCl₃): δ 7.23 (d, 2H); 6.77 (d, 2H); 6.67 (d, 1H); 6.49 (d, 1H);
6.41 (bs, *red.DEAD*); 6.39 (dd, 1H); 4.83-4.77 (m, 1H); 4.22 (q, *red.DEAD*);
4.14-4.07 (m, 1H); 4.01 (d, 2H); 3.75 (s, 3H); 3.01-2.91 (m, 2H); 2.88-2.72 (m, 3H); 2.62
(dd, 1H); 2.29 (hex, 1H); 2.06-1.96 (m, 1H); 1.29 (t, *red.DEAD*)

25 APCI-MS: m/z 393, 395 [MH⁺]

iv) N-[2-(((2S)-3-[(3S)-3-(4-Chlorophenoxy)pyrrolidinyl]-2-hydroxypropyl)oxy)-4-methoxyphenyl]acetamide

To a solution of compound iii) (157mg) in a mixture of acetonitrile (10mL) and water
30 (2mL) acetic anhydride (1mL) was added and the mixture was stirred at ambient

temperature overnight. 1.5M sodiummethoxid in methanol (1mL) was added and the stirring continued for 1h. After evaporation the residue was taken up in ether and water. The subtitle product was obtained from the organic phase as a colourless oil (155mg).

5 ¹H-NMR (400MHz, CDCl₃): δ 8.18 (d, 1H); 7.95 (bs, 1H); 7.24 (d, 2H); 6.78 (d, 2H); 6.56-6.52 (m, 2H); 4.85-4.78 (m, 1H); 4.22 (q, *red.DEAD*); 4.10-4.02 (m, 2H); 4.00-3.92 (m, 1H); 3.78 (s, 3H); 3.00-2.91 (m, 2H); 2.87-2.73 (m, 3H); 2.53 (dd, 1H); 2.36-2.25 (m, 1H); 2.17 (s, 3H); 2.07-1.99 (m, 1H); 1.29 (t, *red.DEAD*)

10 APCI-MS: m/z 435, 437 [MH⁺]

v) 1-[2-((2*S*)-3-[(3*S*)-3-(4-Chlorophenoxy)pyrrolidinyl]-2-hydroxypropyl)oxy]-4-hydroxyphenyl]acetamide

The title compound was prepared by analogy to Example 254 from iv) (150mg).

15 The product obtained after lyophilization was a white amorphous solid (101mg, 57%).

¹H-NMR (400MHz, CDCl₃+1 drop DMSO-*d*₆): δ 8.7 (bs, 1H); 8.34 (s, 1H); 8.73 (d, 1H); 7.18 (d, 2H); 6.73 (d, 2H); 6.40-6.31 (m, 2H); 4.99-4.93 (m, 1H); 4.4-1.9 (bm, 6H); 4.31-4.23 (m, 1H); 3.88-3.78 (m, 2H); 3.39-3.25 (m, 2H);
20 2.4-2.2 (m, 2H); 2.07 (s, 3H)

APCI-MS: m/z 421, 423 [MH⁺]

THP-1 Chemotaxis Assay

25

Introduction

The assay measured the chemotactic response elicited by MIP-1α chemokine in the human monocytic cell line THP-1. The compounds of the Examples were evaluated by their ability to depress the chemotactic response to a standard concentration of MIP-1α

30 chemokine.

Methods

Culture of THP-1 cells

Cells were thawed rapidly at 37°C from frozen aliquots and resuspended in a 25 cm flask containing 5 ml of RPMI-1640 medium supplemented with Glutamax and 10% heat inactivated fetal calf serum without antibiotics (RPMI+10%HIFCS). At day 3 the medium is discarded and replaced with fresh medium.

THP-1 cells are routinely cultured in RPMI-1640 medium supplemented with 10% heat inactivated fetal calf serum and glutamax but without antibiotics. Optimal growth of the cells requires that they are passaged every 3 days and that the minimum subculture density is 4×10^5 cells/ml.

Chemotaxis assay


Cells were removed from the flask and washed by centrifugation in RPMI+10%HIFCS+glutamax. The cells were then resuspended at 2×10^7 cells/ml in fresh medium (RPMI+10%HIFCS+glutamax) to which was added calcein-AM (5 µl of stock solution to 1 ml to give a final concentration of 5×10^{-6} M). After gentle mixing the cells were incubated at 37°C in a CO₂ incubator for 30 minutes. The cells were then diluted to 50 ml with medium and washed twice by centrifugation at 400xg. Labelled cells were then resuspended at a cell concentration of 1×10^7 cells/ml and incubated with an equal volume of MIP-1α antagonist (10^{-10} M to 10^{-6} M final concentration) for 30 minutes at 37°C in a humidified CO₂ incubator.

Chemotaxis was performed using Neuroprobe 96-well chemotaxis plates employing 8 µm filters (cat no. 101-8). Thirty microlitres of chemoattractant supplemented with various concentrations of antagonists or vehicle were added to the lower wells of the plate in triplicate. The filter was then carefully positioned on top and then 25 µl of cells preincubated with the corresponding concentration of antagonist or vehicle were added to the surface of the filter. The plate was then incubated for 2 hours at 37°C in a humidified

CO₂ incubator. The cells remaining on the surface were then removed by adsorption and the whole plate was centrifuged at 2000 rpm for 10 minutes. The filter was then removed and the cells that had migrated to the lower wells were quantified by the fluorescence of cell associated calcein-AM. Cell migration was then expressed in fluorescence units after subtraction of the reagent blank and values were standardized to % migration by comparing the fluorescence values with that of a known number of labelled cells. The effect of antagonists was calculated as % inhibition when the number of migrated cells were compared with vehicle.


CLAIMS

1. A compound of general formula



wherein,

R represents a group



m is 0, 1, 2 or 3;

each R^1 independently represents halogen, cyano, nitro, carboxyl, hydroxyl, C_3 - C_6 cycloalkyl, C_1 - C_6 alkoxy, C_1 - C_6 alkoxycarbonyl, C_1 - C_6 haloalkyl, C_1 - C_6 haloalkoxy, $-NR^9R^{10}$, C_3 - C_6 cycloalkylamino, C_1 - C_6 alkylthio, C_1 - C_6 alkylcarbonyl, C_1 - C_6 alkylcarbonylamino, sulphonamido, C_1 - C_6 alkylsulphonyl, $-C(O)NR^{11}R^{12}$, $-NR^{13}C(O)-(NH)_pR^{14}$, phenyl, or C_1 - C_6 alkyl optionally substituted by carboxyl or C_1 - C_6 alkoxycarbonyl;

p is 0 or 1;

X represents an oxygen or sulphur atom or a CH_2 , $CH(CH_3)$, OCH_2 , CH_2O , CH_2NH , NH or carbonyl group and Y represents a nitrogen atom or a CH or $C(OH)$ group, provided that when X represents an oxygen or sulphur atom or a CH_2O , CH_2NH or NH group, then


Y represents a CH group;

Z^1 represents a bond or a group $(CH_2)_q$ where q is 1 or 2;

Z^2 represents a bond or a group CH_2 , with the proviso that Z^1 and Z^2 do not both simultaneously represent a bond;

Q represents an oxygen or sulphur atom or a group CH_2 or NH ;

R^2 represents a group



n is 0, 1 or 2;

5 each R^3 independently represents a C_1 - C_6 alkyl, C_1 - C_6 alkoxycarbonyl, $-CH_2OH$ or carboxyl group;

R^4 , R^5 , R^6 and R^7 each independently represent a hydrogen atom or a C_1 - C_6 alkyl group, or R^4 , R^5 , R^6 and R^7 together represent a C_1 - C_4 alkylene chain linking the two carbon atoms to which they are attached to form a 4- to 7-membered saturated carbocycle, or R^5 , R^6 and R^7 each represent a hydrogen atom and R^4 and R^8 together with the carbon atoms to which they are attached form a 5- to 6-membered saturated carbocycle;

R^8 represents a hydrogen atom, a C_1 - C_6 alkyl group or is linked to R^4 as defined above;

R^9 and R^{10} each independently represent a hydrogen atom or a C_1 - C_6 alkyl group, or R^9 and R^{10} together with the nitrogen atom to which they are attached form a 4- to 7-membered saturated heterocycle;

R^{11} and R^{12} each independently represent a hydrogen atom or a C_1 - C_6 alkyl group optionally substituted by C_1 - C_6 alkoxycarbonyl;

R^{13} represents a hydrogen atom or a C_1 - C_6 alkyl group;

20 R^{14} represents a hydrogen atom, or a C_1 - C_6 alkyl group optionally substituted by carboxyl, C_1 - C_6 alkoxy or C_1 - C_6 alkoxycarbonyl;

R^{15} represents carboxyl, C_1 - C_6 alkylcarbonyl, C_1 - C_6 alkoxycarbonyl, C_1 - C_6 alkoxycarbonyl C_1 - C_6 alkyl or a group $-NR^{17}R^{18}$, $-NHSO_2CH_3$, $-NHC(O)CH_3$, $-C(O)NR^{17}R^{18}$, $-NHC(O)NR^{17}R^{18}$, $-OC(O)NR^{17}R^{18}$, $-OCH_2C(O)NR^{17}R^{18}$, $-NHC(O)OR^{17'}$ or $-OR^{17''}$;

t is 0, 1, 2 or 3;

each R^{16} independently represents halogen, cyano, nitro, carboxyl, hydroxyl, C_3 - C_6 cycloalkyl, C_1 - C_6 alkoxy, C_1 - C_6 alkoxycarbonyl, C_1 - C_6 haloalkyl, C_1 - C_6 haloalkoxy, $-NR^{19}R^{20}$, C_3 - C_6 cycloalkylamino, C_1 - C_6 alkylthio, C_1 - C_6 alkylcarbonyl, C_1 - C_6 alkylcarbonylamino, sulphonamido ($-SO_2NH_2$), C_1 - C_6 alkylsulphonyl, $-C(O)NR^{21}R^{22}$, $-NR^{23}C(O)(NH)_vR^{24}$, phenyl, or C_1 - C_6 alkyl optionally substituted by carboxyl or C_1 - C_6 alkoxycarbonyl;

R^{17} and R^{18} each independently represent (i) a hydrogen atom, (ii) a 5- to 6-membered saturated or unsaturated ring which may comprise at least one heteroatom chosen from nitrogen, oxygen and sulphur, the ring being optionally substituted with at least one substituent selected from halogen, methyl and trifluoromethyl, or (iii) a C_1 - C_6 alkyl group optionally substituted by at least one substituent selected from halogen, trifluoromethyl, carboxyl, C_1 - C_6 alkoxycarbonyl and a 5- to 6-membered saturated or unsaturated ring which may comprise at least one heteroatom chosen from nitrogen, oxygen and sulphur, the ring being optionally substituted with at least one substituent selected from halogen, methyl and trifluoromethyl, or

R^{17} and R^{18} together with the nitrogen atom to which they are attached form a 4- to 7-membered saturated heterocycle;

R^{17} represents a hydrogen atom, or a C_1 - C_6 alkyl group optionally substituted by carboxyl or C_1 - C_6 alkoxycarbonyl;

$R^{17''}$ is defined as for R^{17} above except that $R^{17''}$ does not represent a hydrogen atom;

R^{19} and R^{20} each independently represent a hydrogen atom or a C_1 - C_6 alkyl group, or R^{19} and R^{20} together with the nitrogen atom to which they are attached form a 4- to 7-membered saturated heterocycle;

R^{21} and R^{22} each independently represent a hydrogen atom or a C_1 - C_6 alkyl group optionally substituted by C_1 - C_6 alkoxycarbonyl;

v is 0 or 1;

R^{23} represents a hydrogen atom or a C_1 - C_6 alkyl group; and

R^{24} represents a hydrogen atom, or a C_1 - C_6 alkyl group optionally substituted by carboxyl, C_1 - C_6 alkoxy or C_1 - C_6 alkoxycarbonyl;

or a pharmaceutically acceptable salt or solvate thereof.

2. A compound according to claim 1, wherein X represents an oxygen atom or a CH₂, OCH₂, CH₂O, NH or carbonyl group.

5 3. A compound according to claim 1 or 2, wherein Y represents a nitrogen atom or CH group.

4. A compound according to any one of claims 1 to 3, wherein Q represents an oxygen atom.

10

5. A compound according to any one of claims 1 to 4, wherein R¹⁵ represents C₁-C₄ alkoxy, C₁-C₄ alkylcarbonyl, C₁-C₄ alkoxy carbonyl C₁-C₄ alkyl, -NHC(O)CH₃, -C(O)NR¹⁷R¹⁸, -NHSO₂CH₃ or -NHC(O)NR¹⁷R¹⁸.

15 6. A compound according to any one of claims 1 to 5, wherein each R¹⁶ independently represents halogen, cyano, hydroxyl, C₁-C₄ alkoxy, C₁-C₄ alkoxy carbonyl, C₁-C₄ haloalkyl, C₁-C₄ alkylcarbonyl, phenyl or C₁-C₄ alkyl.

7. A compound of formula (I'), or a pharmaceutically acceptable salt or solvate thereof,
20 as defined in claim 1 being selected from:

N-(2-{3-[3R,S-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2R,S-hydroxy-propoxy}-phenyl)-acetamide hydrochloride,

N-(5-Chloro-2-{3-[3R,S-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2R,S-hydroxy-propoxy}-phenyl)-acetamide hydrochloride,

25 *N*-(2-{3-[4-(3,4-dichlorophenoxy)-1-piperidiny]-2-hydroxypropoxy} phenyl)-acetamide,

1-(2-aminophenoxy)-3-[4-(3,4-dichlorophenoxy)-1-piperidiny]-2-propanol dihydrochloride,

N-(2-{3-[3-(3,4-dichlorophenoxy)-1-pyrrolidiny]-2-hydroxypropoxy} phenyl)-
30 acetamide hydrochloride,

2-{3-[4-(4-Fluoro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-benzoic acid methyl ester,

2-(2-{3-[4-(3,4-Difluoro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-benzoylamino)-2-methyl-propionic acid methyl ester,

5 N-[2-({(1*R*,2*S*,3*R*)*-3-[4-(3,4-dichlorophenoxy)-1-piperidiny]-2-hydroxycyclopentyl}oxy)phenyl]acetamide,

N-[2-({(1*S*,2*S*,3*R*)*-3-[4-(3,4-dichlorophenoxy)-1-piperidiny]-2-hydroxycyclopentyl}oxy)phenyl]acetamide,

10 N-[2-({(2,3-*trans*)-3-[4-(3,4-dichlorophenoxy)-1-piperidiny]-2-hydroxycyclohexyl}oxy)phenyl]acetamide,

N-(5-Chloro-2-{3-[3-(3,4-dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

N-(3-Acetyl-2-{3-[3-(3,4-dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-5-methyl-phenyl)-acetamide,

15 N-(2-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-4-methyl-phenyl)-acetamide,

N-(2-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-5-fluoro-phenyl)-acetamide,

20 N-(4-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-biphenyl-3-yl)-acetamide,

N-(2-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-4-fluoro-phenyl)-acetamide,

N-(2-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-5-methyl-phenyl)-acetamide,

25 N-(2-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

N-(5-Chloro-2-{3-[3-(4-chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

30 N-(3-Acetyl-2-{3-[3-(4-chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-5-methyl-phenyl)-acetamide,

N-(2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-4-methyl-phenyl)-acetamide,

N-(2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-5-fluoro-phenyl)-acetamide,

5 N-(4-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-biphenyl-3-yl)-acetamide,

N-(2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-4-fluoro-phenyl)-acetamide,

10 N-(2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-5-methyl-phenyl)-acetamide,

N-(2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide

N-(5-Chloro-2-{3-[3-(4-fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

15 N-(3-Acetyl-2-{3-[3-(4-fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-5-methyl-phenyl)-acetamide,

N-(2-{3-[3-(4-Fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-4-methyl-phenyl)-acetamide,

20 N-(5-Fluoro-2-{3-[3-(4-fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

N-(4-{3-[3-(4-Fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-biphenyl-3-yl)-acetamide,

N-(4-Fluoro-2-{3-[3-(4-fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

25 N-(2-{3-[3-(4-Fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-5-methyl-phenyl)-acetamide,

N-(2-{3-[3-(4-Fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

30 N-(5-Chloro-2-{3-[3-(3,4-difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

N-(3-Acetyl-2-{3-[3-(3,4-difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-5-methyl-phenyl)-acetamide,

N-(2-{3-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-4-methyl-phenyl)-acetamide,

5 N-(2-{3-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-5-fluoro-phenyl)-acetamide,

N-(4-{3-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-biphenyl-3-yl)-acetamide,

10 N-(2-{3-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-4-fluoro-phenyl)-acetamide,

N-(2-{3-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-5-methyl-phenyl)-acetamide,

N-(2-{3-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

15 N-(5-Chloro-2-{3-[4-(3,4-dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

N-(3-Acetyl-2-{3-[4-(3,4-dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-5-methyl-phenyl)-acetamide,

20 N-(2-{3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-4-methyl-phenyl)-acetamide,

N-(2-{3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-5-fluoro-phenyl)-acetamide,

N-(4-{3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-biphenyl-3-yl)-acetamide,

25 N-(2-{3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-4-fluoro-phenyl)-acetamide,

N-(2-{3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-5-methyl-phenyl)-acetamide,

30 N-(2-{3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

N-(5-Chloro-2-{3-[4-(4-chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

N-(3-Acetyl-2-{3-[4-(4-chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-5-methyl-phenyl)-acetamide,

5 N-(2-{3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-4-methyl-phenyl)-acetamide,

N-(2-{3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-5-fluoro-phenyl)-acetamide,

10 N-(4-{3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-biphenyl-3-yl)-acetamide,

N-(2-{3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-4-fluoro-phenyl)-acetamide,

N-(2-{3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-5-methyl-phenyl)-acetamide,

15 N-(2-{3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

N-(2-{3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

20 3-(2-{3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propionic acid methyl ester,

1-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-3-(2,6-dimethoxy-phenoxy)-propan-2-ol,

1-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-3-(2-methoxy-phenoxy)-propan-2-ol,

2-{3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-N,N-dimethylbenzamide,

25 1-(2-{3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propan-1-one,

1-(2-{3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-ethanone,

30 3-(2-{3-[3-(4-Fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propionic acid methyl ester,

1-(2,6-Dimethoxy-phenoxy)-3-[3-(4-fluoro-phenoxy)-pyrrolidin-1-yl]-propan-2-ol,
 1-[3-(4-Fluoro-phenoxy)-pyrrolidin-1-yl]-3-(2-methoxy-phenoxy)-propan-2-ol,
 (2-{3-[3-(4-Fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-benzoylamino)-
 acetic acid methyl ester,

5 (2-{3-[3-(4-Chloro-phenoxy)methyl]-piperidin-1-yl]-2-hydroxy-propoxy}-
 benzoylamino)-acetic acid methyl ester,

2-(2-{3-[3-(4-Fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-benzoylamino)-
 2-methyl-propionic acid methyl ester,

2-{3-[3-(4-Fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-N,N-dimethyl-
 10 benzamide,

1-(2-{3-[3-(4-Fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-6-methoxy-
 phenyl)-ethanone,

1-(2-{3-[3-(4-Fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propan-
 1-one,

15 1-(2-{3-[3-(4-Fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-
 ethanone,

N-[2-(3-{[1-(3,4-dichlorobenzyl)-4-piperidinyl]amino}-2-hydroxypropoxy)-4-
 methylphenyl]acetamide,

3-(2-{3-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-
 20 propionic acid methyl ester,

1-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-3-(2-methoxy-phenoxy)-propan-2-ol,
 (2-{3-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-
 benzoylamino)-acetic acid methyl ester,

2-{3-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-N,N-dimethyl-
 25 benzamide,

1-(2-{3-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-6-methoxy-
 phenyl)-ethanone,

1-(2-{3-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-
 propan-1-one,

1-(2-{3-[3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-ethanone,

N-(2-{3-[4-(3,4-dichloroanilino)-1-piperidinyl]-2-hydroxypropoxy}phenyl)acetamide,

3-(2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-

5 propionic acid methyl ester,

1-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-3-(2,6-dimethoxy-phenoxy)-propan-2-ol,

1-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-3-(2-methoxy-phenoxy)-propan-2-ol,

(2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-benzoylamino)-acetic acid, methyl ester,

10 2-(2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-benzoylamino)-2-methyl-propionic acid methyl ester,

2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-*N,N*-dimethylbenzamide,

1-(2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-6-methoxy-phenyl)-ethanone,

1-(2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propan-1-one,

1-(2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-ethanone,

20 *N*-(2-{3-[3-(4-Cyano-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

3-(2-{3-[3-(4-Cyano-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propionic acid methyl ester,

(2-{3-[3-(4-Cyano-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-benzoylamino)-acetic acid methyl ester,

25 2-{3-[3-(4-Cyano-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-*N,N*-dimethylbenzamide,

4-{1-[2-Hydroxy-3-(2-propionyl-phenoxy)-propyl]-pyrrolidin-3-yloxy}-benzonitrile,

N-(2-{2-Hydroxy-3-[3-(4-methoxy-phenoxy)-pyrrolidin-1-yl]-propoxy}-phenyl)-acetamide,

30

N-(4-chloro-2-{3-[4-(3,4-dichloroanilino)-1-piperidinyl]-2-hydroxypropoxy}phenyl)acetamide,

3-(2-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propionic acid methyl ester,

5 1-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-3-(2-methoxy-phenoxy)-propan-2-ol,
(2-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-

benzoylamino)-acetic acid methyl ester,

2-(2-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-benzoylamino)-2-methyl-propionic acid methyl ester,

10 2-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-*N,N*-dimethylbenzamide,

1-(2-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-6-methoxyphenyl)-ethanone,

15 1-(2-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propan-1-one,

1-(2-{3-[3-(3,4-Dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-ethanone,

N-(2-{3-[4-(3,4-Difluoro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

20 3-(2-{3-[4-(3,4-Difluoro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propionic acid methyl ester,

2-{3-[4-(3,4-Difluoro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-*N,N*-dimethylbenzamide,

1-(2-{3-[4-(3,4-Difluoro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propan-1-one,

(2-{3-[4-(3,4-Difluoro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-benzoylamino)-acetic acid methyl ester,

N-(2-{3-[3-(3,4-Difluoro-phoxymethyl)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

N-(2-{3-[4-(4-Fluoro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

3-(2-{3-[4-(4-Fluoro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propionic acid methyl ester,

5 1-(2,6-Dimethoxy-phenoxy)-3-[4-(4-fluoro-phenoxy)-piperidin-1-yl]-propan-2-ol,
1-[4-(4-Fluoro-phenoxy)-piperidin-1-yl]-3-(2-methoxy-phenoxy)-propan-2-ol,
1-(2-{3-[4-(4-Fluoro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-ethanone,
2-{3-[4-(4-Fluoro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-N,N-dimethyl-benzamide,

10 1-(2-{3-[4-(4-Fluoro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propan-1-one,

(2-{3-[4-(4-Fluoro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-benzoylamino)-acetic acid methyl ester,

N-(2-{3-[3-(4-Fluoro-phenoxy)methyl]-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

3-(2-{3-[3-(4-Fluoro-phenoxy)methyl]-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propionic acid methyl ester,

1-[3-(4-Fluoro-phenoxy)methyl]-piperidin-1-yl]-3-(2-methoxy-phenoxy)-propan-2-ol,
1-(2-{3-[3-(4-Fluoro-phenoxy)methyl]-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-ethanone,

20 2-(2-{3-[3-(4-Fluoro-phenoxy)methyl]-piperidin-1-yl]-2-hydroxy-propoxy}-benzoylamino)-2-methyl-propionic acid methyl ester,

2-{3-[3-(4-Fluoro-phenoxy)methyl]-piperidin-1-yl]-2-hydroxy-propoxy}-N,N-dimethyl-benzamide,

25 1-(2-{3-[3-(4-Fluoro-phenoxy)methyl]-piperidin-1-yl]-2-hydroxy-propoxy}-6-methoxy-phenyl)-ethanone,

N-(2-{3-[4-(4-Acetylamino-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

N-(4-{1-[3-(2-Acetyl-phenoxy)-2-hydroxy-propyl]-piperidin-4-yloxy}-phenyl)-acetamide,

N-(4-cyano-2-{3-[4-(3,4-dichloroanilino)-1-piperidinyl]-2-hydroxypropoxy}phenyl)acetamide,

3-(2-{3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propionic acid methyl ester,

5 1-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-3-(2-methoxy-phenoxy)-propan-2-ol,

1-(2-{3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-ethanone,

2-(2-{3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-benzoylamino)-2-methyl-propionic acid methyl ester,

10 2-{3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-*N,N*-dimethylbenzamide,

1-(2-{3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-6-methoxyphenyl)-ethanone,

1-(2-{3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propan-1-one,

(2-{3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-benzoylamino)-acetic acid methyl ester,

N-(2-{3-[3-(4-Chloro-phenoxy)methyl]-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)acetamide,

20 3-(2-{3-[3-(4-Chloro-phenoxy)methyl]-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propionic acid methyl ester,

1-(2-{3-[3-(4-Chloro-phenoxy)methyl]-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-ethanone,

2-{3-[3-(4-Chloro-phenoxy)methyl]-piperidin-1-yl]-2-hydroxy-propoxy}-*N,N*-dimethyl-benzamide,

25 1-(2-{3-[3-(4-Chloro-phenoxy)methyl]-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-propan-1-one,

N-[2-((1*R*,2*R*)-2-[4-(3,4-dichlorophenoxy)-1-piperidinyl]-1-hydroxycyclopentyl)methoxy]phenyl]acetamide,

Methyl (2*S*,4*R*)-1-{3-[2-(acetylamino)phenoxy]-2-hydroxypropyl}-4-[(4-chlorobenzyl)oxy]-2-pyrrolidinecarboxylate hydrochloride,

N-(2-{3-[4-(3,4-Dichloroanilino)-1-piperidinyl]-2-hydroxypropoxy}-4-methylphenyl)acetamide,

5 N-(2-{3-[4-(4-Chloroanilino)-1-piperidinyl]-2-hydroxypropoxy}phenyl)acetamide,

N-(4-Chloro-2-{3-[4-(4-chloroanilino)-1-piperidinyl]-2-hydroxypropoxy}phenyl)-acetamide,

N-(2-{3-[4-(4-Chloroanilino)-1-piperidinyl]-2-hydroxypropoxy}-4-cyanophenyl)acetamide,

10 N-(2-{3-[4-(4-Chloroanilino)-1-piperidinyl]-2-hydroxypropoxy}-4-methylphenyl)acetamide,

N-(5-Chloro-2-{3-[4-(4-fluoroanilino)-1-piperidinyl]-2-hydroxypropoxy}phenyl)acetamide,

15 N-(5-Chloro-2-{3-[4-(3,4-difluoroanilino)-1-piperidinyl]-2-hydroxypropoxy}phenyl)acetamide,

N-(5-Cyano-2-{3-[4-(4-fluoroanilino)-1-piperidinyl]-2-hydroxypropoxy}phenyl)acetamide,

N-(5-Cyano-2-{3-[4-(3,4-difluoroanilino)-1-piperidinyl]-2-hydroxypropoxy}phenyl)acetamide,

20 N-(2-{3-[4-(4-Fluoroanilino)-1-piperidinyl]-2-hydroxypropoxy}-4-methylphenyl)acetamide,

N-(2-{3-[4-(3,4-Difluoroanilino)-1-piperidinyl]-2-hydroxypropoxy}-4-methylphenyl)acetamide,

25 N-(2-{3-[3(*S*)-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-(*R*)-hydroxy-propoxy}phenyl)acetamide,

N-(2-{3-[3*S*-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2*S*-hydroxy-propoxy}-phenyl)-acetamide hydrochloride,

N-(2-{3-[3(*R*)-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-(*S*)-hydroxy-propoxy}phenyl)acetamide,

N-[5-Chloro-2-({(2S)-3-[(3S)-3-(4-chloro-phenoxy)pyrrolidinyl]-2-hydroxypropyl}oxy)phenyl]acetamide,

N-[5-Chloro-2-({(2R)-3-[(3R)-3-(4-chloro-phenoxy)pyrrolidinyl]-2-hydroxypropyl}oxy)phenyl]acetamide,

5 N-[5-Chloro-2-({(2S)-3-[(3R)-3-(4-chloro-phenoxy)pyrrolidinyl]-2-hydroxypropyl}oxy)phenyl]acetamide,

N-[5-Chloro-2-({(2R)-3-[(3S)-3-(4-chloro-phenoxy)pyrrolidinyl]-2-hydroxypropyl}oxy)phenyl]acetamide,

10 N-(2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-4,5-difluorophenyl)-acetamide,

N-{5-Chloro-2-[2-hydroxy-3-(3-phenoxy-pyrrolidin-1-yl)-propoxy]-phenyl}-acetamide,

N-(5-Chloro-2-{2-hydroxy-3-[3-(4-nitro-phenoxy)-pyrrolidin-1-yl]-propoxy}-phenyl)-acetamide,

15 N-(5-Acetyl-2-{3-[3-(3,4-dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

4-Acetylamino-3-{3-[3-(3,4-dichloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-benzoic acid methyl ester,

20 N-(2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-5-cyanophenyl)-acetamide,

4-Acetylamino-3-{3-[3-(4-chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-benzoic acid methyl ester,

N-(5-Cyano-2-{3-[4-(3,4-dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

25 N-(2-{3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-propoxy}-5-trifluoromethyl-phenyl)-acetamide,

N-(5-Chloro-2-{3-[3-(4-fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide trifluoroacetate,

30 N-(5-Acetyl-2-{3-[3-(4-chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide trifluoroacetate,

N-(2-{3-[3-(4-Chlorophenoxy)-1-pyrrolidinyl]-2-hydroxypropoxy}phenyl)-methanesulfonamide,

N-(5-Chloro-2-[3-[3,4-dichlorophenoxy)-1-pyrrodinyl]-2-hydroxypropoxy)-phenyl)urea,

5 1-(3-{2-[(Aminocarbonyl)amino]phenoxy}-2-hydroxypropyl)-3-(4-chlorophenoxy)pyrrolidinium 2,2,2-trifluoroacetate,

1-(3-{2-[(Aminocarbonyl)amino]phenoxy}-2-hydroxypropyl)-3-(3,4-dichlorophenoxy)pyrrolidinium 2,2,2-trifluoroacetate,

10 1-(3-{2-[(Aminocarbonyl)amino]-4-chlorophenoxy}-2-hydroxypropyl)-3-(4-chlorophenoxy)pyrrolidinium 2,2,2-trifluoroacetate,

N-(2-{3-[3-(4-Chlorophenoxy)-1-pyrrolidinyl]-2-hydroxypropoxy}phenyl)-*N*'-ethylurea hydrochloride,

N-(2-{3-[3-(4-Chlorophenoxy)-1-pyrrolidinyl]-2-hydroxypropoxy}phenyl)-*N*'-methylurea hydrochloride,

15 (2*S*,4*S*)-1-{3-[2-(Acetylamino)phenoxy]-2-hydroxypropyl}-4-(4-chlorophenoxy)-2-pyrrolidinecarboxylic acid; compound with trifluoroacetic acid,

Ethyl (2*S*,4*S*)-1-{3-[2-(acetylamino)phenoxy]-2-hydroxypropyl}-4-(3,4-dichlorophenoxy)-2-pyrrolidinecarboxylate; trifluoroacetic acid salt,

20 *N*-[2-({(2*S*)-3-[(2*S*,4*S*)-4-(4-Chlorophenoxy)-2-(hydroxymethyl)pyrrolidinyl]-2-hydroxypropyl}oxy)phenyl]acetamide; trifluoroacetic acid salt,

N-[2-({(2*R*)-3-[(2*S*,4*S*)-4-(4-Chlorophenoxy)-2-(hydroxymethyl)pyrrolidinyl]-2-hydroxypropyl}oxy)phenyl]acetamide; trifluoroacetic acid salt,

N-(2-{3-[3-(4-Chlorophenoxy)-1-pyrrolidinyl]-2-hydroxy-2-methylpropoxy}phenyl)acetamide hydrochloride,

25 *N*-(2-{(1*S*^{*},2*R*^{*},3*S*^{*})-3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-cyclopentyloxy}-phenyl)-acetamide,

N-(2-{(1*R*^{*},2*R*^{*},3*S*^{*})-3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-cyclopentyloxy}-phenyl)-acetamide,

30 *N*-(2-{(2*R*^{*},3*R*^{*})-3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-butoxy}-phenyl)-acetamide,

N-(2-((1*S*,2*R*,3*S*)-3-[4-(4-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-cyclopentyloxy)-phenyl)-acetamide,

N-(2-((2*R*,3*S*)-3-[4-(3,4-Dichloro-phenoxy)-piperidin-1-yl]-2-hydroxy-butoxy)-phenyl)-acetamide,

5 N-(2-((2*R*,3*R*)-3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-butoxy)-phenyl)-acetamide,

N-(2-((2*R*,3*S*)-3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-butoxy)-phenyl)-acetamide,

10 N-(2-((1*S*,2*R*,3*S*)-3-[4-(3-Chloro-phenoxy)-piperidin-1-yl]-2-hydroxy-cyclopentyloxy)-phenyl)-acetamide,

N-[5-Chloro-2-((1*S*,2*R*,3*S*)-3-[4-(3,4-dichlorophenoxy)-1-piperidinyl]-2-hydroxycyclopentyl)oxy]phenyl]acetamide,

N-[4-Fluoro-2-((1*S*,2*R*,3*S*)-3-[4-(3,4-dichlorophenoxy)-1-piperidinyl]-2-hydroxycyclopentyl)oxy]phenyl]acetamide,

15 N-(2-{3-[4-(3,4-Dichlorobenzyl)-1-piperazinyl]-2-hydroxypropoxy}-phenyl)acetamide dihydrochloride,

N-(2-{3-[4-(3,4-Dichlorobenzyl)-1-piperazinyl]-2-hydroxypropoxy}-4-fluorophenyl)acetamide,

N-(2-{3-[4-(3,4-Dichlorobenzyl)-2,5-dimethyl-1-piperazinyl]-2-hydroxypropoxy}phenyl)acetamide,

20 N-(5-Chloro-2-{3-[4-(3,4-dichlorobenzyl)-1-piperazinyl]-2-hydroxypropoxy}phenyl)acetamide,

N-(5-Chloro-2-{3-[4-(3,4-dichlorobenzyl)-2,5-dimethyl-1-piperazinyl]-2-hydroxypropoxy}phenyl)acetamide,

25 N-(2-{3-[4-(3,4-Dichlorobenzyl)-2,5-dimethyl-1-piperazinyl]-2-hydroxypropoxy}-4-methylphenyl)acetamide,

N-(2-{3-[4-(3,4-Dichlorobenzyl)-2,5-dimethyl-1-piperazinyl]-2-hydroxypropoxy}-4-fluorophenyl)acetamide,

30 N-(2-{3 [(*S***R*)-4-(3,4-Dichlorobenzyl)-2,5-dimethyl-1-piperazinyl]-2-hydroxypropoxy}phenyl)acetamide,

N-(2-{3 [(*S***R**)-4-(4-Chlorobenzyl)-2,5-dimethyl-1-piperazinyl]-2-hydroxypropoxy}phenyl)acetamide,

N-(5-Chloro-2-{3-[(*S***R**)-4-(3,4-dichlorobenzyl)-2,5-dimethylpiperazinyl]-2-hydroxypropoxy}phenyl)acetamide,

5 *N*-(5-Chloro-2-{3-[(*S***R**)-4-(4-chlorobenzyl)-2,5-dimethylpiperazinyl]-2-hydroxypropoxy}phenyl)acetamide,

1-(5-Chloro-2-{3-[4-(4-chlorobenzoyl)-1-piperazinyl]-2-hydroxypropoxy}phenyl)-1-ethanone,

N-(5-Cyano-2-{3-[(*S***R**)-4-(3,4-dichlorobenzyl)-2,5-dimethylpiperazinyl]-2-hydroxypropoxy}phenyl)acetamide,

N-(2-{3-[(*S***R**)-4-(4-Chlorobenzyl)-2,5-dimethylpiperazinyl]-2-hydroxypropoxy}-5-cyanophenyl)acetamide,

N-(5-Chloro-2-{3-[4-(4-chlorobenzyl)-1-piperazinyl]-2-hydroxypropoxy}-phenyl)acetamide,

15 *N*-(4-Chloro-2-{3-[4-(4-chlorobenzyl)-2,5-dimethyl-1-piperazinyl]-2-hydroxypropoxy}phenyl)acetamide,

N-(2-{3-[4-(4-Chlorobenzoyl)-1-piperazinyl]-2-hydroxypropoxy}-5-cyanophenyl)acetamide,

20 *N*-(2-{3-[4-(4-Chlorobenzoyl)-1-piperazinyl]-2-hydroxypropoxy}-4-methylphenyl)acetamide,

N-[5-Chloro-2-((1*R*,2*S*,3*R*)-3-[(3*S*)-3-(4-chlorophenoxy)pyrrolidinyl]-2-hydroxycyclopentyl}oxy)phenyl]acetamide,

N-(2-[(2*S*)-(3-[(3*S*)-3-[(4-Chlorophenyl)oxy]-1-pyrrolidinyl]-2-hydroxypropyl)oxy]-4-fluorophenyl)acetamide,

25 *N*-(2-[(2*S*)-3-[(3*S*)-3-(4-Chlorobenzyl)pyrrolidinyl]-2-hydroxypropyl}oxy)phenyl]acetamide hydrochloride,

N-(5-Chloro-2-{3-[3-(4-chloro-benzyl)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide trifluoroacetic acid salt,

30 *N*-(2-{3-[3-(4-Chlorophenoxy)-1-pyrrolidinyl]-2-hydroxypropoxy}-4-methylphenyl)-1-pyrrolidinecarboxamide trifluoroacetate,

N-(2-{3-[3-(4-Chlorophenoxy)-1-pyrrolidinyl]-2-hydroxypropoxy}-4-hydroxyphenyl)acetamide trifluoroacetate,

N-[2-({(2*S*)-3-[4-(3,4-Dichlorophenoxy)-1-piperidinyl]-2-hydroxypropyl}oxy)-4-fluorophenyl]acetamide trifluoroacetic acid salt,

5 *N*-(2-(3-(4-Chloro-phenoxy)-pyrrolidin-1-yl)-2-hydroxy-propoxy)-4,6-difluorophenyl)-acetamide hydrochloride,

N-[2-({(2*S*)-3-[(2*S*,4*S*)-4-(4-Chlorophenoxy)-2-methylpyrrolidinyl]-2-hydroxypropyl}oxy)-4-fluorophenyl]acetamide trifluoroacetic acid salt,

N-[2-({(2*S*)-3-[(3*R*)-3-(4-Chlorobenzyl)pyrrolidinyl]-2-hydroxypropyl}oxy)phenyl]acetamide hydrochloride,

10 *N*-{2-[(2*R*)- (3-{(3*S*)-3-[(4-Chlorophenyl)oxy]-1-pyrrolidinyl}-2-hydroxypropyl)oxy]-4-fluorophenyl} acetamide,

N-[2-({(2*S*)-3-[(2*R*,4*S*)-4-(4-Chlorophenoxy)-2-methylpyrrolidinyl]-2-hydroxypropyl}oxy) phenyl]acetamide trifluoroacetic acid salt,

15 *N*-{2-[(2*S*)-(3-{(3*R*)-3-[(4-Chlorophenyl)oxy]-1-pyrrolidinyl}-2-hydroxypropyl)oxy]-4-fluorophenyl} acetamide,

N'-(2-{3-[3-(4-Chlorophenoxy)-1-pyrrolidinyl]-2-hydroxypropoxy}-4-methylphenyl)-*N,N*-dimethylurea trifluoroacetate,

N-(2-{3-[3-(4-Chloroanilino)-1-pyrrolidinyl]-2-hydroxypropoxy} phenyl)acetamide,

20 *N*-{2-[(3-{3-[(4-Chlorophenyl)oxy]-1-pyrrolidinyl}-2-hydroxy-1-methylpropyl)oxy]phenyl} acetamide hydrochloride,

N-(2-{3-[3-(4-Chlorophenoxy)-1-pyrrolidinyl]-2-hydroxypropoxy}-4-methoxyphenyl)acetamide hydrochloride,

N-(2-[3-(4-Chloro-benzyloxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy)-phenyl)-acetamide trifluoroacetic acid salt,

25 *N*-(2-{3-[3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-2-methyl-propoxy}-phenyl)-acetamide,

N-(2-{{(1*S*,2*R*,3*S*)*-3-[(3*S*)-3-(3,4-Difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-cyclopentyloxy}-5-chloro-phenyl)-acetamide (diastereomeric mixture),

N-[2-({(2*R*,3*S*)*-3-[(3*S*)-3-(4-Chlorophenoxy)pyrrolidinyl]-2-hydroxybutyl}oxy)-4-methylphenyl]acetamide (diastereomeric mixture),

N-{2-[(3-{4-[(3,4-Dichlorophenyl)oxy]-1-piperidinyl}-2-hydroxy-2-methylpropyl)oxy]-4-fluorophenyl}acetamide hydrochloride,

5 *N*-(2-({(1*S*,2*R*,3*S*)*-3-[(3*S*)-3-(4-Chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-cyclopentyloxy}-4-fluoro-phenyl)-acetamide (diastereomeric mixture),

N-(5-Chloro-2-{3-[3-(3,4-difluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

10 *N*-(5-Chloro-2-{3-[3-(4-fluoro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-propoxy}-phenyl)-acetamide,

N-(4-Cyano-2-{3-[4-(3,4-dichloroanilino)-1-piperidinyl]-2-hydroxypropoxy} phenyl)acetamide,

N-(4-Hydroxy-2-({(1*S*,2*R*,3*S*)*-3-[(3*S*)-3-(4-chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-cyclopentyloxy}-phenyl)-acetamide (diastereomeric mixture),

15 *N*-(4-Hydroxy-2-({(1*S*,2*R*,3*S*)-3-[(3*S*)-3-(4-chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-cyclopentyloxy}-phenyl)-acetamide,

N-(4-Hydroxy-2-({(1*R*,2*S*,3*R*)-3-[(3*S*)-3-(4-chloro-phenoxy)-pyrrolidin-1-yl]-2-hydroxy-cyclopentyloxy}-phenyl)-acetamide,

20 *N*-[2-({(1*S*,2*R*,3*S*)-3-[(3*S*)-3-(4-Chlorophenoxy)pyrrolidinyl]-2-hydroxycyclopentyl}oxy)phenyl]acetamide,

N-[2-({(1*R*,2*S*,3*R*)-3-[(3*S*)-3-(4-Chlorophenoxy)pyrrolidinyl]-2-hydroxycyclopentyl}oxy)phenyl]acetamide,

N-[5-Chloro-2-({(1*S*,2*R*,3*S*)-3-[(3*S*)-3-(4-chlorophenoxy)pyrrolidinyl]-2-hydroxycyclopentyl}oxy)phenyl]acetamide,

25 *N*-{5-Chloro-2-[(1*S*,2*R*,3*S*)*-3-{[1-(4-chlorobenzyl)-4-piperidinyl]amino}-2-hydroxycyclopentyl]oxy}phenyl}acetamide (racemic mixture), and


N-[2-({(2*S*)-3-[(3*S*)-3-(4-Chlorophenoxy)pyrrolidinyl]-2-hydroxypropyl}oxy)-4-hydroxyphenyl]acetamide.

8. A process for the preparation of a compound of formula (I') as defined in claim 1 which comprises,

(a) reacting a compound of general formula




5 wherein R is as defined in formula (I'), with a compound of general formula




wherein Q, R², R⁴, R⁵, R⁶, R⁷ and R⁸ are as defined in formula (I'); or

(b) reacting a compound of general formula



wherein R, R⁴, R⁵, R⁶, R⁷ and R⁸ are as defined in formula (I'), with a compound of general formula



15 wherein L¹ represents a hydrogen atom or an activating group and Q and R² are as defined in formula (I');

and optionally thereafter converting the compound of formula (I') to a further compound of formula (I'); and, if desired, forming a pharmaceutically acceptable salt or solvate of the compound of formula (I').

9. A pharmaceutical composition comprising a compound of formula (I'), or a pharmaceutically acceptable salt or solvate thereof, as claimed in any one of claims 1 to 7 in association with a pharmaceutically acceptable adjuvant, diluent or carrier.

10. A process for the preparation of a pharmaceutical composition as claimed in claim 9 which comprises mixing a compound of formula (I'), or a pharmaceutically acceptable salt or solvate thereof, as claimed in any one of claims 1 to 7 with a pharmaceutically acceptable adjuvant, diluent or carrier.

5

11. A compound of formula (I'), or a pharmaceutically-acceptable salt or solvate thereof, as claimed in any one of claims 1 to 7 for use in therapy.

10

12. Use of a compound of formula (I'), or a pharmaceutically acceptable salt or solvate thereof, as claimed in any one of claims 1 to 7 in the manufacture of a medicament for use in therapy.

15

13. Use of a compound of formula (I'), or a pharmaceutically acceptable salt or solvate thereof, as claimed in any one of claims 1 to 7 in the manufacture of a medicament for the treatment of human diseases or conditions in which modulation of chemokine receptor activity is beneficial.

20

14. Use of a compound of formula (I'), or a pharmaceutically acceptable salt or solvate thereof, as claimed in any one of claims 1 to 7 in the manufacture of a medicament for use in treating rheumatoid arthritis.

25

15. Use of a compound of formula (I'), or a pharmaceutically acceptable salt or solvate thereof, as claimed in any one of claims 1 to 7 in the manufacture of a medicament for use in treating chronic obstructive pulmonary disease.

16. Use of a compound of formula (I'), or a pharmaceutically acceptable salt or solvate thereof, as claimed in any one of claims 1 to 7 in the manufacture of a medicament for use in treating asthma.

17. Use of a compound of formula (I'), or a pharmaceutically acceptable salt or solvate thereof, as claimed in any one of claims 1 to 7 in the manufacture of a medicament for use in treating multiple sclerosis.

5 18. A method of treating an inflammatory disease in a patient suffering from, or at risk of, said disease, which comprises administering to the patient a therapeutically effective amount of a compound of formula (I'), or a pharmaceutically acceptable salt or solvate thereof, as claimed in any one of claims 1 to 7.

10 19. A method of treating an airways disease in a patient suffering from, or at risk of, said disease, which comprises administering to the patient a therapeutically effective amount of a compound of formula (I'), or a pharmaceutically acceptable salt or solvate thereof, as claimed in any one of claims 1 to 7.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 01/00403

A. CLASSIFICATION OF SUBJECT MATTER

IPC7: C07D 207/04, C07D 211/06, C07D 295/08, A61K 31/40, A61K 31/435, A61P 11/00,
A61P 37/00, A61P 19/00 A61P 31/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7: C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 0058305 A1 (ASTRAZENECA UK LIMITED), 5 October 2000 (05.10.00), page 49, line 20 - page 50, line 9; page 88, line 11 - page 89, line 9; page 97, line 1 - page 98, line 5, page 99, line 1 - line 6; page 104, line 5 - page 105, line 10; page 107, line 1 - line 5; the claims --	1-19
X	Bioorganic & Medicinal Chemistry Letters, Volume 7, No 11, 1997, Jon L. Wright et al, "Discovery of selective dopamine D4 receptor antagonists: 1-ARYLOXY-3-(4-ARYLOXYPIPERIDINYL)-2-PROPANOLS" page 1377 - page 1380 --	1-12

☒ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

18 May 2001

Date of mailing of the international search report

31 -05- 2001

Name and mailing address of the ISA/

Swedish Patent Office

Box 5055, S-102 42 STOCKHOLM

Facsimile No. +46 8 666 02 86

Authorized officer

Gerd Strandell/Eö

Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 01/00403

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 3577432 A (GROVER CLEVELAND HELSLEY), 4 May 1971 (04.05.71), column 13, line 1 - line 15; column 13 - column 14, Table 4; the claims --	1-12
X	US 5789402 A (JAMES E. AUDIA ET AL), 4 August 1998 (04.08.98), column 38, line 36 - line 54; the claims --	1-12
A	EP 0903349 A2 (F. HOFFMANN-LA ROCHE AG), 24 March 1999 (24.03.99) -- -----	1-19

INTERNATIONAL SEARCH REPORT

International application No.
PCT/SE01/00403

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: **18, 19**
because they relate to subject matter not required to be searched by this Authority, namely:
see next sheet
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

Claims 18, 19 relate to methods of treatment of the human or animal body by surgery or by therapy/ diagnostic methods practised on the human or animal body/Rule 39.1.(iv). Nevertheless, a search has been executed for these claims. The search has been based on the alleged effects of the compounds/compositions.

INTERNATIONAL SEARCH REPORT
Information on patent family members

02/04/01

International application No.

PCT/SE 01/00403

Patent document cited in search report			Publication date	Patent family member(s)		Publication date
WO	0058305	A1	05/10/00	AU	4157500 A	16/10/00
				AU	4942599 A	17/01/00
				SE	9901117 D	00/00/00
				AU	2013500 A	13/06/00
				SE	9902194 D	00/00/00

US	3577432	A	04/05/71	BR	6915185 D	00/00/00
				CA	918663 A	09/01/73
				CH	538473 A	30/06/73
				DE	1964511 A	09/07/70
				FR	2026921 A,B	25/09/70
				GB	1289867 A	20/09/72
				JP	49014225 B	05/04/74
				JP	50034026 B	05/11/75
				JP	50034027 B	05/11/75
JP	50034028 B	05/11/75				

US	5789402	A	04/08/98	AU	718875 B	20/04/00
				AU	4651696 A	07/08/96
				BR	9607077 A	18/11/97
				CA	2210220 A	25/07/96
				CZ	9702207 A	18/02/98
				EP	0722941 A	24/07/96
				FI	973024 A	16/07/97
				HU	9901099 A	28/07/99
				JP	10512861 T	08/12/98
				NO	973281 A	08/09/97
				NZ	301161 A	28/10/99
				PL	321851 A	22/12/97
				TR	9700644 T	00/00/00
				US	5576321 A	19/11/96
				US	5614523 A	25/03/97
				US	5627196 A	06/05/97
				US	5741789 A	21/04/98
				US	6172073 B	09/01/01
				WO	9622290 A	25/07/96

INTERNATIONAL SEARCH REPORT

Information on patent family members

02/04/01

International application No.

PCT/SE 01/00403

Patent document cited in search report			Publication date	Patent family member(s)		Publication date
EP	0903349	A2	24/03/99	AU	8080098 A	25/02/99
				BR	9803179 A	28/03/00
				CA	2245043 A	18/02/99
				CN	1211572 A	24/03/99
				CZ	9802566 A	17/03/99
				DE	19837386 A	25/02/99
				FR	2767826 A	05/03/99
				GB	2330580 A	28/04/99
				GB	9817910 D	00/00/00
				HR	980450 A	30/06/99
				HU	9801861 D	00/00/00
				HU	9801887 A	28/06/99
				IL	125658 D	00/00/00
				IT	MI981902 A	18/02/00
				JP	3014367 B	28/02/00
				JP	11147872 A	02/06/99
				NO	983749 A	19/02/99
				PL	328049 A	01/03/99
				SG	70110 A	25/01/00
				TR	9801594 A	00/00/00
				ZA	9807448 A	22/01/99

THIS PAGE BLANK (USPTO)